



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 05:05 pm GMT

PDB ID : 3DNF
Title : Structure of (E)-4-Hydroxy-3-methyl-but-2-enyl Diphosphate Reductase, the
Terminal Enzyme of the Non-Mevalonate Pathway
Authors : Rekittke, I.
Deposited on : 2008-07-02
Resolution : 1.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	1368 (1.66-1.66)
Clashscore	112137	1468 (1.66-1.66)
Ramachandran outliers	110173	1438 (1.66-1.66)
Sidechain outliers	110143	1438 (1.66-1.66)
RSRZ outliers	101464	1371 (1.66-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	297	<div> <div>17%</div> <div> <div></div> <div>80%</div> <div>13%</div> <div>• 5%</div> </div> </div>
1	B	297	<div> <div>13%</div> <div> <div></div> <div>82%</div> <div>12%</div> <div>• 5%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	B	3732	-	-	-	X

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5076 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called 4-hydroxy-3-methylbut-2-enyl diphosphate reductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	281	Total	C	N	O	S	0	3	0
			2215	1408	383	415	9			
1	B	282	Total	C	N	O	S	0	12	0
			2274	1455	390	420	9			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	290	ARG	-	EXPRESSION TAG	UNP O67625
A	291	SER	-	EXPRESSION TAG	UNP O67625
A	292	HIS	-	EXPRESSION TAG	UNP O67625
A	293	HIS	-	EXPRESSION TAG	UNP O67625
A	294	HIS	-	EXPRESSION TAG	UNP O67625
A	295	HIS	-	EXPRESSION TAG	UNP O67625
A	296	HIS	-	EXPRESSION TAG	UNP O67625
A	297	HIS	-	EXPRESSION TAG	UNP O67625
B	290	ARG	-	EXPRESSION TAG	UNP O67625
B	291	SER	-	EXPRESSION TAG	UNP O67625
B	292	HIS	-	EXPRESSION TAG	UNP O67625
B	293	HIS	-	EXPRESSION TAG	UNP O67625
B	294	HIS	-	EXPRESSION TAG	UNP O67625
B	295	HIS	-	EXPRESSION TAG	UNP O67625
B	296	HIS	-	EXPRESSION TAG	UNP O67625
B	297	HIS	-	EXPRESSION TAG	UNP O67625

- Molecule 2 is FE3-S4 CLUSTER (three-letter code: F3S) (formula: Fe₃S₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	Fe	S	0	0
			7	3	4		
2	B	1	Total	Fe	S	0	0
			7	3	4		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	243	Total 243	O 243	0	0
4	B	318	Total 318	O 318	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 4-hydroxy-3-methylbut-2-enyl diphosphate reductase



- Molecule 1: 4-hydroxy-3-methylbut-2-enyl diphosphate reductase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	60.37Å 87.76Å 72.40Å 90.00° 95.13° 90.00°	Depositor
Resolution (Å)	9.90 – 1.65 9.86 – 1.65	Depositor EDS
% Data completeness (in resolution range)	91.3 (9.90-1.65) 91.4 (9.86-1.65)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.04	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.18 (at 1.65Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.193 , 0.226 0.201 , 0.231	Depositor DCC
R_{free} test set	4120 reflections (5.28%)	DCC
Wilson B-factor (Å ²)	22.5	Xtriage
Anisotropy	0.077	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.45 , 59.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5076	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.49% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, F3S

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.83	5/2259 (0.2%)	0.73	2/3052 (0.1%)
1	B	0.71	2/2348 (0.1%)	0.69	0/3170
All	All	0.77	7/4607 (0.2%)	0.71	2/6222 (0.0%)

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	237	ASN	CB-CG	12.81	1.80	1.51
1	A	237	ASN	CG-OD1	9.14	1.44	1.24
1	B	211	ASP	CG-OD2	6.90	1.41	1.25
1	A	246	GLU	C-O	6.51	1.35	1.23
1	A	246	GLU	C-N	5.62	1.47	1.34
1	B	211	ASP	CG-OD1	5.52	1.38	1.25
1	A	250	GLU	CB-CG	5.42	1.62	1.52

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	239	TYR	CZ-CE2-CD2	-5.79	114.59	119.80
1	A	239	TYR	CB-CG-CD1	-5.77	117.54	121.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2215	0	2261	25	0
1	B	2274	0	2365	39	0
2	A	7	0	0	1	0
2	B	7	0	0	0	0
3	A	6	0	8	2	0
3	B	6	0	8	1	0
4	A	243	0	0	1	0
4	B	318	0	0	9	0
All	All	5076	0	4642	62	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (62) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:237:ASN:CB	1:A:237:ASN:CG	1.80	1.50
1:B:37[B]:LEU:HD12	1:B:69[B]:VAL:CG1	1.72	1.18
1:B:37[B]:LEU:CD1	1:B:69[B]:VAL:CG1	2.26	1.13
1:B:37[B]:LEU:CD1	1:B:69[B]:VAL:HG12	1.87	1.02
1:B:37[B]:LEU:HD12	1:B:69[B]:VAL:HG13	1.47	0.95
1:B:37[B]:LEU:HD11	1:B:69[B]:VAL:CG1	2.03	0.88
1:B:37[B]:LEU:HD12	1:B:69[B]:VAL:HG12	1.48	0.86
1:B:124:HIS:HD2	1:B:126:GLU:H	1.22	0.86
1:A:124:HIS:HD2	1:A:126:GLU:H	1.32	0.75
1:B:37[B]:LEU:HD11	1:B:69[B]:VAL:HG12	1.67	0.71
1:B:17:LYS:NZ	4:B:4008:HOH:O	2.21	0.70
1:B:104:GLU:OE1	4:B:4136:HOH:O	2.11	0.68
1:B:124:HIS:CD2	1:B:126:GLU:H	2.08	0.68
1:A:247:LEU:HD13	1:A:278:ILE:HG13	1.79	0.64
1:A:124:HIS:CD2	1:A:126:GLU:H	2.14	0.64
1:B:243:THR:HG22	1:B:245:GLU:H	1.63	0.63
1:A:247:LEU:HD11	1:A:274:VAL:HG13	1.81	0.62
1:B:37[B]:LEU:HD11	1:B:69[B]:VAL:HG11	1.80	0.62
1:B:211:ASP:HB2	1:B:255:VAL:HG13	1.82	0.61
1:A:40:ILE:HG23	1:A:41:ILE:HD12	1.83	0.59
1:B:48:ASN:OD1	4:B:4175:HOH:O	2.15	0.58
1:B:59:GLN:OE1	4:B:4176:HOH:O	2.16	0.58
1:B:223:ASN:HB2	4:B:4165:HOH:O	2.03	0.58
1:B:277:ARG:HD3	1:B:277:ARG:O	2.05	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:33:LYS:HE3	1:B:56:PHE:CD2	2.41	0.55
1:B:247:LEU:HD11	1:B:274:VAL:HG13	1.89	0.55
1:B:104:GLU:HG2	4:B:4101:HOH:O	2.06	0.54
1:B:204:LYS:HG3	1:B:230:ILE:HD11	1.89	0.54
1:A:248:GLN:HB2	1:A:251:TRP:CE2	2.43	0.53
1:B:196:THR:HG23	4:B:3891:HOH:O	2.08	0.53
1:B:156:HIS:HD2	3:B:3732:GOL:O2	1.92	0.53
1:A:4:ILE:HG13	1:A:4:ILE:O	2.10	0.51
1:B:217:GLY:HA3	1:B:224:THR:HG21	1.93	0.50
1:B:203:VAL:HG21	1:B:227:LEU:HD22	1.94	0.50
1:B:204:LYS:CG	1:B:230:ILE:HD11	2.42	0.50
1:A:237:ASN:CG	1:A:237:ASN:CA	2.72	0.49
1:B:156:HIS:HE1	4:B:3908:HOH:O	1.95	0.48
1:B:4:ILE:H	1:B:4:ILE:HD12	1.78	0.47
1:A:40:ILE:O	1:A:41:ILE:HG23	2.15	0.47
1:A:156:HIS:HD2	3:A:3731:GOL:O2	1.97	0.47
1:A:142:ILE:HD13	1:A:152:GLU:HB3	1.95	0.47
1:A:28:LYS:HG2	1:A:29:GLU:HG3	1.96	0.47
1:A:256:LYS:HE3	1:A:257:ARG:HG3	1.97	0.47
1:B:37[B]:LEU:CD1	1:B:69[B]:VAL:HG11	2.31	0.46
1:A:72:ARG:NH2	1:A:74:HIS:HE1	2.14	0.46
1:B:267:PRO:HD2	1:B:270:ILE:HD12	1.98	0.46
1:A:74:HIS:HD2	1:A:75:GLY:O	1.98	0.45
1:A:137[B]:CYS:SG	4:A:3800:HOH:O	2.61	0.45
1:A:217:GLY:HA3	1:A:224:THR:HG21	2.00	0.44
1:B:210:VAL:HA	1:B:257:ARG:HE	1.83	0.43
1:A:3:ASP:HB3	1:A:5:ILE:HG12	1.99	0.43
1:A:15:GLY:HA3	2:A:2302:F3S:S4	2.58	0.43
1:A:248:GLN:HE21	1:A:251:TRP:HZ2	1.65	0.43
1:A:187:LYS:HE3	1:B:185[A]:GLU:CD	2.40	0.42
1:B:6:ILE:HD11	1:B:275:LYS:HD2	2.02	0.42
1:A:187:LYS:HE3	1:B:185[A]:GLU:OE2	2.20	0.42
1:A:43:ASN:C	1:A:43:ASN:HD22	2.22	0.42
1:B:74:HIS:HE1	4:B:4021:HOH:O	2.02	0.41
1:B:124:HIS:HD2	1:B:126:GLU:N	2.03	0.41
1:A:113:GLY:O	3:A:3731:GOL:H31	2.21	0.41
1:B:107:CYS:SG	1:B:137[A]:CYS:SG	3.14	0.40
1:B:227:LEU:O	1:B:230:ILE:HG22	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	282/297 (95%)	278 (99%)	3 (1%)	1 (0%)	38	17
1	B	292/297 (98%)	287 (98%)	5 (2%)	0	100	100
All	All	574/594 (97%)	565 (98%)	8 (1%)	1 (0%)	51	28

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	247	LEU

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	243/255 (95%)	228 (94%)	15 (6%)	21	4
1	B	253/255 (99%)	249 (98%)	4 (2%)	68	45
All	All	496/510 (97%)	477 (96%)	19 (4%)	41	11

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MET
1	A	2	VAL
1	A	4	ILE
1	A	41	ILE
1	A	43	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	63	PHE
1	A	193[A]	CYS
1	A	193[C]	CYS
1	A	197[A]	SER
1	A	197[B]	SER
1	A	198	LEU
1	A	233	GLU
1	A	234	LEU
1	A	235	ASN
1	A	277	ARG
1	B	1	MET
1	B	4	ILE
1	B	277	ARG
1	B	282	CYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	43	ASN
1	A	74	HIS
1	A	123	ASN
1	A	124	HIS
1	A	156	HIS
1	A	235	ASN
1	B	9	HIS
1	B	59	GLN
1	B	124	HIS
1	B	156	HIS
1	B	223	ASN
1	B	273	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

There are no carbohydrates in this entry.

5.6 Ligand geometry

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	F3S	A	2302	1	0,9,9	0.00	-	0,15,15	0.00	-
3	GOL	A	3731	-	5,5,5	0.31	0	5,5,5	0.46	0
2	F3S	B	3302	1	0,9,9	0.00	-	0,15,15	0.00	-
3	GOL	B	3732	-	5,5,5	0.20	0	5,5,5	0.66	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	F3S	A	2302	1	-	0/0/24/24	0/0/3/3
3	GOL	A	3731	-	-	0/4/4/4	0/0/0/0
2	F3S	B	3302	1	-	0/0/24/24	0/0/3/3
3	GOL	B	3732	-	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	2302	F3S	1	0
3	A	3731	GOL	2	0
3	B	3732	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	281/297 (94%)	1.14	51 (18%)	1 1	28, 33, 41, 75	0
1	B	282/297 (94%)	0.93	38 (13%)	3 3	27, 33, 40, 71	0
All	All	563/594 (94%)	1.03	89 (15%)	2 2	27, 33, 40, 75	0

All (89) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	2	VAL	12.4
1	A	1	MET	12.2
1	B	1	MET	10.8
1	B	2	VAL	9.0
1	A	42	HIS	8.4
1	B	250	GLU	6.7
1	B	255	VAL	6.5
1	A	4	ILE	6.5
1	A	3	ASP	6.3
1	B	281	ILE	6.3
1	A	279	GLN	6.2
1	A	253	ARG	6.1
1	A	280	GLU	6.0
1	B	253	ARG	6.0
1	B	3	ASP	5.9
1	B	254	GLY	5.9
1	A	281	ILE	5.9
1	B	282	CYS	5.4
1	A	257	ARG	5.4
1	A	256	LYS	5.2
1	A	276	SER	4.9
1	A	5	ILE	4.6
1	B	233	GLU	4.4
1	A	254	GLY	4.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	245	GLU	4.4
1	A	251	TRP	4.3
1	A	250	GLU	4.3
1	A	245	GLU	4.2
1	B	223	ASN	4.0
1	B	236	PRO	4.0
1	A	255	VAL	4.0
1	B	251	TRP	3.9
1	A	249	PRO	3.8
1	A	238	THR	3.8
1	A	205	LYS	3.8
1	A	226	ARG	3.6
1	A	244	ALA	3.5
1	A	222	GLY	3.5
1	A	210	VAL	3.5
1	A	278	ILE	3.5
1	A	220	ASN	3.4
1	B	123[A]	ASN	3.4
1	A	248	GLN	3.4
1	B	280	GLU	3.4
1	B	249	PRO	3.4
1	A	193[A]	CYS	3.4
1	A	8	GLU	3.3
1	B	239	TYR	3.1
1	B	42	HIS	3.1
1	A	229	TYR	3.1
1	A	233	GLU	3.1
1	A	31	GLN	3.1
1	B	220	ASN	3.0
1	B	248	GLN	3.0
1	A	273	GLN	3.0
1	B	226	ARG	2.9
1	A	148	GLU	2.9
1	B	279	GLN	2.9
1	B	4	ILE	2.9
1	A	221	SER	2.8
1	A	277	ARG	2.8
1	B	257	ARG	2.8
1	B	148	GLU	2.7
1	B	277	ARG	2.7
1	B	229	TYR	2.7
1	A	197[A]	SER	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	228	TYR	2.6
1	A	237	ASN	2.6
1	A	209	GLU	2.6
1	B	238	THR	2.6
1	B	252	PHE	2.6
1	B	182	TRP	2.5
1	B	98	TYR	2.4
1	B	5	ILE	2.4
1	A	243	THR	2.4
1	B	208	PRO	2.4
1	A	269	TRP	2.4
1	B	96	CYS	2.3
1	A	123	ASN	2.3
1	A	211	ASP	2.2
1	B	212	VAL	2.2
1	A	30	SER	2.2
1	A	241	ILE	2.2
1	B	127	VAL	2.2
1	A	239	TYR	2.1
1	B	114	TYR	2.1
1	A	91	VAL	2.1
1	A	107	CYS	2.1
1	A	207	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	GOL	B	3732	6/6	0.89	0.22	2.88	32,37,41,45	0
3	GOL	A	3731	6/6	0.91	0.18	0.75	24,33,37,39	0
2	F3S	B	3302	7/7	0.95	0.15	0.55	24,27,33,37	0
2	F3S	A	2302	7/7	0.95	0.12	-0.13	21,25,29,35	0

6.5 Other polymers [i](#)

There are no such residues in this entry.