



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 14, 2017 – 06:18 pm GMT

PDB ID : 5DOI
Title : Crystal structure of Tetrahymena p45N and p19
Authors : Wan, B.; Tang, T.; Wu, J.; Lei, M.
Deposited on : 2015-09-11
Resolution : 2.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	4002 (2.20-2.20)
Clashscore	112137	4730 (2.20-2.20)
Ramachandran outliers	110173	4656 (2.20-2.20)
Sidechain outliers	110143	4657 (2.20-2.20)
RSRZ outliers	101464	4033 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	163	<div> <div>0%</div> <div> <div>78%</div> <div>13%</div> <div>6%</div> </div> </div>
1	B	163	<div> <div>4%</div> <div> <div>69%</div> <div>23%</div> <div>6%</div> </div> </div>
1	C	163	<div> <div>3%</div> <div> <div>73%</div> <div>18%</div> <div>6%</div> </div> </div>
1	D	163	<div> <div>2%</div> <div> <div>77%</div> <div>13%</div> <div>6%</div> </div> </div>
2	E	128	<div> <div>2%</div> <div> <div>48%</div> <div>24%</div> <div>27%</div> </div> </div>
2	F	128	<div> <div>3%</div> <div> <div>55%</div> <div>18%</div> <div>27%</div> </div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	G	128	<div><div><div></div><div></div><div></div></div><div><div>2%</div><div>52%</div><div>19%</div><div>•</div><div>27%</div></div></div>
2	H	128	<div><div><div></div><div></div><div></div></div><div><div>2%</div><div>55%</div><div>18%</div><div></div><div>27%</div></div></div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 8613 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Telomerase-associated protein 19.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	153	Total	C	N	O	S	0	0	0
			1281	826	213	240	2			
1	B	153	Total	C	N	O	S	0	0	0
			1281	826	213	240	2			
1	C	153	Total	C	N	O	S	0	0	0
			1281	826	213	240	2			
1	D	153	Total	C	N	O	S	0	0	0
			1281	826	213	240	2			

- Molecule 2 is a protein called Telomerase associated protein p45.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	E	93	Total	C	N	O	S	0	0	0
			745	494	115	133	3			
2	F	93	Total	C	N	O	S	0	0	0
			745	494	115	133	3			
2	G	93	Total	C	N	O	S	0	0	0
			745	494	115	133	3			
2	H	93	Total	C	N	O	S	0	0	0
			745	494	115	133	3			

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	113	Total	O	0	0
			113	113		
3	B	66	Total	O	0	0
			66	66		
3	C	108	Total	O	0	0
			108	108		
3	D	107	Total	O	0	0
			107	107		


Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	E	29	Total	O	0	0
			29	29		
3	F	27	Total	O	0	0
			27	27		
3	G	30	Total	O	0	0
			30	30		
3	H	29	Total	O	0	0
			29	29		

- Molecule 1: Telomerase-associated protein 19







• Molecule 2: Telomerase associated protein p45




• Molecule 2: Telomerase associated protein p45



• Molecule 2: Telomerase associated protein p45



• Molecule 2: Telomerase associated protein p45



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	44.19Å 125.83Å 180.17Å 90.00° 90.04° 90.00°	Depositor
Resolution (Å)	44.19 – 2.20 90.08 – 2.20	Depositor EDS
% Data completeness (in resolution range)	96.8 (44.19-2.20) 96.6 (90.08-2.20)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.18 (at 2.20Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.226 , 0.264 0.239 , 0.268	Depositor DCC
R_{free} test set	4828 reflections (5.00%)	DCC
Wilson B-factor (Å ²)	27.3	Xtriage
Anisotropy	0.284	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 36.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.468 for h,-k,-l	Xtriage
Reported twinning fraction	0.490 for h,-k,-l	Depositor
Outliers	4 of 96759 reflections (0.004%)	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	8613	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 53.49 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 4.1861e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.58	1/1306 (0.1%)	0.62	0/1756
1	B	0.60	0/1306	0.64	0/1756
1	C	0.52	0/1306	0.64	1/1756 (0.1%)
1	D	0.53	1/1306 (0.1%)	0.62	0/1756
2	E	0.38	0/754	0.57	0/1008
2	F	0.37	0/754	0.57	0/1008
2	G	0.36	0/754	0.55	0/1008
2	H	0.39	0/754	0.57	0/1008
All	All	0.50	2/8240 (0.0%)	0.61	1/11056 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	145	TYR	CZ-OH	-5.25	1.28	1.37
1	D	145	TYR	CZ-OH	-5.12	1.29	1.37

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	98	LYS	CD-CE-NZ	7.47	128.89	111.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1281	0	1266	16	0
1	B	1281	0	1266	22	0
1	C	1281	0	1266	23	1
1	D	1281	0	1266	17	0
2	E	745	0	787	20	0
2	F	745	0	787	12	0
2	G	745	0	787	16	0
2	H	745	0	787	18	0
3	A	113	0	0	5	1
3	B	66	0	0	1	1
3	C	108	0	0	4	0
3	D	107	0	0	4	1
3	E	29	0	0	5	0
3	F	27	0	0	1	0
3	G	30	0	0	0	0
3	H	29	0	0	2	0
All	All	8613	0	8212	135	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 135 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:E:94:GLN:HG2	3:E:216:HOH:O	1.33	1.29
2:E:94:GLN:CG	3:E:216:HOH:O	2.03	0.84
1:A:46:PHE:O	3:A:201:HOH:O	1.94	0.84
1:A:31:GLN:O	3:A:202:HOH:O	2.05	0.74
1:C:121:ASP:OD2	3:C:201:HOH:O	2.05	0.74

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:210:HOH:O	3:B:254:HOH:O[2_655]	1.68	0.52
1:C:116:SER:OG	3:D:296:HOH:O[2_656]	2.17	0.03

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	149/163 (91%)	141 (95%)	7 (5%)	1 (1%)	25	24
1	B	149/163 (91%)	141 (95%)	8 (5%)	0	100	100
1	C	149/163 (91%)	142 (95%)	7 (5%)	0	100	100
1	D	149/163 (91%)	144 (97%)	5 (3%)	0	100	100
2	E	85/128 (66%)	82 (96%)	3 (4%)	0	100	100
2	F	85/128 (66%)	82 (96%)	3 (4%)	0	100	100
2	G	85/128 (66%)	80 (94%)	5 (6%)	0	100	100
2	H	85/128 (66%)	83 (98%)	2 (2%)	0	100	100
All	All	936/1164 (80%)	895 (96%)	40 (4%)	1 (0%)	55	63

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	80	ASN

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	142/152 (93%)	138 (97%)	4 (3%)	49	61
1	B	142/152 (93%)	137 (96%)	5 (4%)	41	51
1	C	142/152 (93%)	135 (95%)	7 (5%)	29	35
1	D	142/152 (93%)	133 (94%)	9 (6%)	21	23

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	E	85/117 (73%)	80 (94%)	5 (6%)	23	26
2	F	85/117 (73%)	80 (94%)	5 (6%)	23	26
2	G	85/117 (73%)	80 (94%)	5 (6%)	23	26
2	H	85/117 (73%)	81 (95%)	4 (5%)	30	37
All	All	908/1076 (84%)	864 (95%)	44 (5%)	30	36

5 of 44 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	63	SER
1	D	161	ILE
2	H	29	LEU
1	D	106	LEU
1	D	149	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	80	ASN
1	A	101	GLN
1	C	80	ASN
1	C	153	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	153/163 (93%)	-0.08	2 (1%) 77 75	20, 29, 48, 93	0
1	B	153/163 (93%)	0.07	6 (3%) 40 38	20, 32, 60, 107	0
1	C	153/163 (93%)	-0.01	5 (3%) 47 44	19, 29, 54, 106	0
1	D	153/163 (93%)	-0.01	3 (1%) 65 63	19, 32, 65, 107	0
2	E	93/128 (72%)	0.15	2 (2%) 62 60	23, 38, 67, 101	0
2	F	93/128 (72%)	0.12	4 (4%) 36 34	22, 36, 70, 84	0
2	G	93/128 (72%)	0.10	3 (3%) 48 46	20, 39, 73, 117	0
2	H	93/128 (72%)	0.10	2 (2%) 62 60	23, 36, 69, 89	0
All	All	984/1164 (84%)	0.04	27 (2%) 55 52	19, 33, 66, 117	0

The worst 5 of 27 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	85	LYS	5.2
1	B	7	ASN	4.1
2	E	64	ASP	4.1
2	G	64	ASP	4.0
1	B	85	LYS	3.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.