



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 15, 2017 – 01:45 am GMT

PDB ID : 1DV6  
Title : PHOTOSYNTHETIC REACTION CENTER FROM RHODOBACTER SPHAEROIDES IN THE CHARGE-NEUTRAL DQAQB STATE WITH THE PROTON TRANSFER INHIBITOR ZN2+  
Authors : Axelrod, H.L.; Abresch, E.C.; Paddock, M.L.; Okamura, M.Y.; Feher, G.  
Deposited on : 2000-01-19  
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	3846 (2.50-2.50)
Clashscore	112137	4554 (2.50-2.50)
Ramachandran outliers	110173	4463 (2.50-2.50)
Sidechain outliers	110143	4465 (2.50-2.50)
RSRZ outliers	101464	3876 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	L	281	<div> <div>2%</div> <div> <div></div> <div>77%</div> <div>21%</div> <div>•</div> </div> </div>
1	R	281	<div> <div>5%</div> <div> <div></div> <div>74%</div> <div>25%</div> <div>•</div> </div> </div>
2	M	307	<div> <div>%</div> <div> <div></div> <div>77%</div> <div>19%</div> <div>• •</div> </div> </div>
2	S	307	<div> <div>%</div> <div> <div></div> <div>75%</div> <div>21%</div> <div>• •</div> </div> </div>
3	H	260	<div> <div>4%</div> <div> <div></div> <div>73%</div> <div>21%</div> <div>5%</div> </div> </div>
3	T	260	<div> <div>8%</div> <div> <div></div> <div>70%</div> <div>23%</div> <div>• 5%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
10	LDA	M	1012	-	-	-	X
10	LDA	M	1013	-	-	-	X
10	LDA	M	1014	-	-	-	X
10	LDA	S	2012	-	-	-	X
7	BCL	S	2003	-	-	-	X
9	U10	L	1009	-	-	-	X
9	U10	R	2009	-	-	-	X

## 2 Entry composition

There are 11 unique types of molecules in this entry. The entry contains 14349 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PHOTOSYNTHETIC REACTION CENTER.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	L	281	Total	C	N	O	S	0	0	0
			2232	1507	355	362	8			
1	R	281	Total	C	N	O	S	0	0	0
			2232	1507	355	362	8			

- Molecule 2 is a protein called PHOTOSYNTHETIC REACTION CENTER.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	M	299	Total	C	N	O	S	0	0	0
			2390	1597	391	392	10			
2	S	299	Total	C	N	O	S	0	0	0
			2390	1597	391	392	10			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
M	307	ALA	ASN	CONFLICT	UNP P02953
S	307	ALA	ASN	CONFLICT	UNP P02953

- Molecule 3 is a protein called PHOTOSYNTHETIC REACTION CENTER.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	H	246	Total	C	N	O	S	0	0	0
			1869	1196	320	343	10			
3	T	246	Total	C	N	O	S	0	0	0
			1869	1196	320	343	10			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
H	8	GLN	GLY	CONFLICT	UNP P11846
T	8	GLN	GLY	CONFLICT	UNP P11846

- Molecule 4 is FE (II) ION (three-letter code: FE2) (formula: Fe).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	S	1	Total	Fe	0	0
			1	1		
4	M	1	Total	Fe	0	0
			1	1		


- Molecule 5 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	H	1	Total	Zn	0	0
			1	1		
5	T	1	Total	Zn	0	0
			1	1		

- Molecule 6 is CHLORIDE ION (three-letter code: CL) (formula: Cl).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	S	1	Total	Cl	0	0
			1	1		
6	M	1	Total	Cl	0	0
			1	1		

- Molecule 7 is BACTERIOCHLOROPHYLL A (three-letter code: BCL) (formula: C<sub>55</sub>H<sub>74</sub>MgN<sub>4</sub>O<sub>6</sub>).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	M	1	Total	C	Mg	N	O	
			51	40	1	4	6	
7	L	1	Total	C	Mg	N	O	
			66	55	1	4	6	
7	M	1	Total	C	Mg	N	O	
			66	55	1	4	6	
7	M	1	Total	C	Mg	N	O	
			66	55	1	4	6	
7	S	1	Total	C	Mg	N	O	
			51	40	1	4	6	
7	R	1	Total	C	Mg	N	O	
			66	55	1	4	6	
7	S	1	Total	C	Mg	N	O	
			66	55	1	4	6	
7	R	1	Total	C	Mg	N	O	
			66	55	1	4	6	

- Molecule 8 is BACTERIOPHEOPHYTIN A (three-letter code: BPH) (formula:  $C_{55}H_{76}N_4O_6$ ).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
8	M	1	Total	C	N	O		
			51	41	4	6		
8	L	1	Total	C	N	O		
			65	55	4	6		
8	S	1	Total	C	N	O		
			52	42	4	6		
8	R	1	Total	C	N	O		
			65	55	4	6		

- Molecule 9 is UBIQUINONE-10 (three-letter code: U10) (formula:  $C_{59}H_{90}O_4$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	M	1	Total	C	O	0	0
			38	34	4		
9	L	1	Total	C	O	0	0
			44	40	4		
9	S	1	Total	C	O	0	0
			32	28	4		
9	R	1	Total	C	O	0	0
			18	14	4		

- Molecule 10 is LAURYL DIMETHYLAMINE-N-OXIDE (three-letter code: LDA) (formula:  $C_{14}H_{31}NO$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
10	M	1	Total	C	N	O	0	0
			16	14	1	1		
10	M	1	Total	C	N	O	0	0
			16	14	1	1		
10	M	1	Total	C	N	O	0	0
			10	8	1	1		
10	S	1	Total	C	N	O	0	0
			16	14	1	1		

- Molecule 11 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
11	H	82	Total	O	0	0
			82	82		
11	L	75	Total	O	0	0
			75	75		
11	M	92	Total	O	0	0
			92	92		
11	R	55	Total	O	0	0
			55	55		
11	S	75	Total	O	0	0
			75	75		
11	T	61	Total	O	0	0
			61	61		




### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: PHOTOSYNTHETIC REACTION CENTER



- Molecule 1: PHOTOSYNTHETIC REACTION CENTER




- Molecule 2: PHOTOSYNTHETIC REACTION CENTER




- Molecule 2: PHOTOSYNTHETIC REACTION CENTER






### • Molecule 3: PHOTOSYNTHETIC REACTION CENTER



### • Molecule 3: PHOTOSYNTHETIC REACTION CENTER



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	141.52Å 141.52Å 278.51Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	27.52 – 2.50 27.48 – 2.50	Depositor EDS
% Data completeness (in resolution range)	96.4 (27.52-2.50) 96.7 (27.48-2.50)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.09 (at 2.51Å)	Xtriage
Refinement program	CNS 0.9	Depositor
R, $R_{free}$	0.238 , 0.265 0.233 , 0.261	Depositor DCC
$R_{free}$ test set	4424 reflections (4.90%)	DCC
Wilson B-factor (Å <sup>2</sup> )	49.8	Xtriage
Anisotropy	0.275	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 55.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	14349	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.80% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BCL, ZN, CL, BPH, LDA, FE2, U10

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	L	0.41	0/2320	0.56	0/3175
1	R	0.41	0/2320	0.56	0/3175
2	M	0.42	0/2482	0.53	0/3389
2	S	0.43	0/2482	0.54	0/3389
3	H	0.35	0/1917	0.59	0/2608
3	T	0.36	0/1917	0.59	0/2608
All	All	0.40	0/13438	0.56	0/18344

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	2232	0	2187	55	0
1	R	2232	0	2187	61	0
2	M	2390	0	2304	54	0
2	S	2390	0	2304	60	0
3	H	1869	0	1884	38	0
3	T	1869	0	1884	57	0
4	M	1	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	S	1	0	0	0	0
5	H	1	0	0	0	0
5	T	1	0	0	0	0
6	M	1	0	0	1	0
6	S	1	0	0	0	0
7	L	66	0	74	9	0
7	M	183	0	188	21	0
7	R	132	0	148	12	0
7	S	117	0	115	12	0
8	L	65	0	76	6	0
8	M	51	0	45	2	0
8	R	65	0	76	7	0
8	S	52	0	47	4	0
9	L	44	0	57	3	0
9	M	38	0	47	0	0
9	R	18	0	15	2	0
9	S	32	0	39	2	0
10	M	42	0	78	8	0
10	S	16	0	31	1	0
11	H	82	0	0	2	0
11	L	75	0	0	5	0
11	M	92	0	0	3	0
11	R	55	0	0	7	0
11	S	75	0	0	14	0
11	T	61	0	0	17	0
All	All	14349	0	13786	335	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 335 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:T:239:GLY:HA2	11:T:2042:HOH:O	1.54	1.05
2:S:267:ARG:HA	11:S:2068:HOH:O	1.66	0.94
1:R:195:LEU:HD11	11:S:2068:HOH:O	1.70	0.91
1:L:217:ARG:HD2	11:M:1077:HOH:O	1.77	0.83
3:T:178:PHE:HB3	11:T:2036:HOH:O	1.78	0.82

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	L	279/281 (99%)	262 (94%)	17 (6%)	0	100	100
1	R	279/281 (99%)	262 (94%)	17 (6%)	0	100	100
2	M	297/307 (97%)	288 (97%)	9 (3%)	0	100	100
2	S	297/307 (97%)	288 (97%)	9 (3%)	0	100	100
3	H	244/260 (94%)	234 (96%)	10 (4%)	0	100	100
3	T	244/260 (94%)	234 (96%)	10 (4%)	0	100	100
All	All	1640/1696 (97%)	1568 (96%)	72 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	220/220 (100%)	212 (96%)	8 (4%)	40	67
1	R	220/220 (100%)	211 (96%)	9 (4%)	35	61
2	M	235/239 (98%)	225 (96%)	10 (4%)	33	58
2	S	235/239 (98%)	225 (96%)	10 (4%)	33	58
3	H	199/209 (95%)	192 (96%)	7 (4%)	41	68
3	T	199/209 (95%)	192 (96%)	7 (4%)	41	68
All	All	1308/1336 (98%)	1257 (96%)	51 (4%)	37	63

5 of 51 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	H	228	LEU
1	R	129	LEU
3	T	135	LYS
1	R	7	ARG
1	R	189	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 18 such sidechains are listed below:

Mol	Chain	Res	Type
3	H	206	ASN
1	R	87	GLN
2	S	300	ASN
3	H	194	GLN
3	H	199	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 26 ligands modelled in this entry, 6 are monoatomic - leaving 20 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
7	BCL	L	1002	1	55,74,74	1.33	5 (9%)	65,115,115	1.29	9 (13%)
8	BPH	L	1006	-	65,70,70	1.25	9 (13%)	75,101,101	1.61	13 (17%)
9	U10	L	1009	-	44,44,63	1.66	9 (20%)	53,56,79	1.25	5 (9%)
7	BCL	M	1001	2	40,59,74	1.60	8 (20%)	47,97,115	1.72	15 (31%)
7	BCL	M	1003	2	55,74,74	1.26	7 (12%)	65,115,115	1.35	12 (18%)
7	BCL	M	1004	1	55,74,74	1.25	7 (12%)	65,115,115	1.46	14 (21%)
8	BPH	M	1005	-	51,56,70	1.17	5 (9%)	58,84,101	1.97	12 (20%)
9	U10	M	1008	-	38,38,63	1.77	9 (23%)	46,49,79	1.08	4 (8%)
10	LDA	M	1012	-	13,15,15	2.63	1 (7%)	14,17,17	2.53	3 (21%)
10	LDA	M	1013	-	13,15,15	2.86	1 (7%)	14,17,17	2.30	4 (28%)
10	LDA	M	1014	-	7,9,15	3.88	1 (14%)	8,11,17	3.02	2 (25%)
7	BCL	R	2002	1	55,74,74	1.38	5 (9%)	65,115,115	1.54	13 (20%)
7	BCL	R	2004	1	55,74,74	1.24	7 (12%)	65,115,115	1.40	10 (15%)
8	BPH	R	2006	-	65,70,70	0.99	5 (7%)	75,101,101	1.72	12 (16%)
9	U10	R	2009	-	18,18,63	1.84	3 (16%)	22,25,79	1.07	1 (4%)
7	BCL	S	2001	2	40,59,74	1.54	8 (20%)	47,97,115	1.56	11 (23%)
7	BCL	S	2003	2	55,74,74	1.33	6 (10%)	65,115,115	1.37	10 (15%)
8	BPH	S	2005	-	52,57,70	1.18	8 (15%)	59,85,101	1.85	10 (16%)
9	U10	S	2008	-	32,32,63	1.61	6 (18%)	38,41,79	1.03	2 (5%)
10	LDA	S	2012	-	13,15,15	2.81	1 (7%)	14,17,17	2.43	2 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	BCL	L	1002	1	-	0/37/137/137	0/0/9/9
8	BPH	L	1006	-	-	0/54/105/105	0/1/6/6
9	U10	L	1009	-	-	0/41/65/87	0/1/1/1
7	BCL	M	1001	2	-	0/19/119/137	0/0/9/9
7	BCL	M	1003	2	-	0/37/137/137	0/0/9/9
7	BCL	M	1004	1	-	0/37/137/137	0/0/9/9
8	BPH	M	1005	-	-	0/38/89/105	0/1/6/6
9	U10	M	1008	-	-	0/33/57/87	0/1/1/1
10	LDA	M	1012	-	-	0/13/13/13	0/0/0/0
10	LDA	M	1013	-	-	0/13/13/13	0/0/0/0
10	LDA	M	1014	-	-	0/7/7/13	0/0/0/0

*Continued on next page...*



*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	BCL	R	2002	1	-	0/37/137/137	0/0/9/9
7	BCL	R	2004	1	-	0/37/137/137	0/0/9/9
8	BPH	R	2006	-	-	0/54/105/105	0/1/6/6
9	U10	R	2009	-	-	0/9/33/87	0/1/1/1
7	BCL	S	2001	2	-	0/19/119/137	0/0/9/9
7	BCL	S	2003	2	-	0/37/137/137	0/0/9/9
8	BPH	S	2005	-	-	0/39/90/105	0/1/6/6
9	U10	S	2008	-	-	0/26/50/87	0/1/1/1
10	LDA	S	2012	-	-	0/13/13/13	0/0/0/0

The worst 5 of 111 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
10	M	1014	LDA	O1-N1	-10.16	1.22	1.42
10	M	1013	LDA	O1-N1	-10.03	1.22	1.42
10	S	2012	LDA	O1-N1	-9.84	1.22	1.42
10	M	1012	LDA	O1-N1	-9.24	1.24	1.42
9	L	1009	U10	C7-C8	-3.95	1.44	1.50

The worst 5 of 164 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	R	2006	BPH	C4D-C3D-CAD	-8.46	102.94	107.78
8	S	2005	BPH	C4D-C3D-CAD	-8.07	103.16	107.78
10	M	1012	LDA	CM2-N1-CM1	-8.03	95.62	110.99
10	S	2012	LDA	CM2-N1-CM1	-7.84	96.00	110.99
10	M	1014	LDA	CM2-N1-CM1	-7.76	96.14	110.99

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

19 monomers are involved in 76 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	L	1002	BCL	9	0
8	L	1006	BPH	6	0
9	L	1009	U10	3	0
7	M	1001	BCL	6	0
7	M	1003	BCL	8	0
7	M	1004	BCL	8	0
8	M	1005	BPH	2	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Clashes	Symm-Clashes
10	M	1012	LDA	1	0
10	M	1013	LDA	2	0
10	M	1014	LDA	5	0
7	R	2002	BCL	5	0
7	R	2004	BCL	8	0
8	R	2006	BPH	7	0
9	R	2009	U10	2	0
7	S	2001	BCL	6	0
7	S	2003	BCL	7	0
8	S	2005	BPH	4	0
9	S	2008	U10	2	0
10	S	2012	LDA	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	L	281/281 (100%)	0.11	5 (1%) 69 70	30, 49, 74, 85	0
1	R	281/281 (100%)	0.00	14 (4%) 30 31	33, 50, 73, 85	0
2	M	299/307 (97%)	0.09	4 (1%) 77 78	34, 44, 59, 76	0
2	S	299/307 (97%)	-0.12	4 (1%) 77 78	36, 46, 59, 75	0
3	H	246/260 (94%)	0.11	11 (4%) 34 36	40, 52, 73, 90	0
3	T	246/260 (94%)	0.33	20 (8%) 13 12	42, 53, 74, 90	0
All	All	1652/1696 (97%)	0.08	58 (3%) 44 47	30, 49, 71, 90	0

The worst 5 of 58 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	T	92	VAL	4.8
3	T	55	PRO	4.4
1	L	59	TRP	4.2
3	H	255	MET	4.0
3	T	255	MET	3.9

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
10	LDA	S	2012	16/16	0.57	0.46	7.61	51,61,69,70	0
10	LDA	M	1014	10/16	0.76	0.40	7.20	51,53,57,58	0
9	U10	L	1009	44/63	0.76	0.30	4.30	70,83,97,97	0
10	LDA	M	1012	16/16	0.80	0.26	3.56	57,61,69,69	0
10	LDA	M	1013	16/16	0.85	0.26	3.20	53,56,58,58	0
9	U10	R	2009	18/63	0.86	0.23	3.12	73,75,76,76	0
7	BCL	S	2003	66/66	0.93	0.20	2.16	41,42,55,56	0
9	U10	S	2008	32/63	0.94	0.17	1.57	52,54,56,56	0
8	BPH	R	2006	65/65	0.93	0.16	1.50	50,56,60,62	0
7	BCL	L	1002	66/66	0.95	0.20	1.29	34,38,41,43	0
7	BCL	R	2002	66/66	0.93	0.16	1.09	40,44,51,54	0
8	BPH	L	1006	65/65	0.92	0.17	0.61	31,40,43,44	0
7	BCL	S	2001	51/66	0.93	0.14	0.49	38,42,49,51	0
7	BCL	M	1003	66/66	0.95	0.18	0.34	29,35,44,49	0
8	BPH	S	2005	52/65	0.94	0.14	0.28	41,44,59,62	0
9	U10	M	1008	38/63	0.91	0.18	0.07	31,37,54,54	0
7	BCL	M	1004	66/66	0.95	0.18	-0.11	29,33,53,57	0
7	BCL	M	1001	51/66	0.96	0.14	-0.25	33,35,45,47	0
7	BCL	R	2004	66/66	0.95	0.13	-0.27	35,42,66,68	0
8	BPH	M	1005	51/65	0.96	0.14	-0.48	31,33,40,42	0
4	FE2	M	1007	1/1	0.98	0.13	-1.65	35,35,35,35	0
4	FE2	S	2007	1/1	1.00	0.06	-2.54	42,42,42,42	0
5	ZN	T	2010	1/1	0.95	0.04	-	60,60,60,60	0
6	CL	M	1011	1/1	0.96	0.15	-	45,45,45,45	0
6	CL	S	2011	1/1	0.95	0.18	-	66,66,66,66	0
5	ZN	H	1010	1/1	0.97	0.04	-	57,57,57,57	0

## 6.5 Other polymers ⓘ

There are no such residues in this entry.