



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 03:02 am GMT

PDB ID : 2DWI
Title : Crystal structure of the complex formed between C-terminal half of bovine lactoferrin and cellobiose at 2.2 Å resolution
Authors : Prem Kumar, R.; Mir, R.; Sinha, M.; Singh, N.; Sharma, S.; Kaur, P.; Bhushan, A.; Singh, T.P.
Deposited on : 2006-08-13
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	112137	4730 (2.20-2.20)
Ramachandran outliers	110173	4656 (2.20-2.20)
Sidechain outliers	110143	4657 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	345	79%17%..

2 Entry composition

There are 10 unique types of molecules in this entry. The entry contains 3143 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Lactotransferrin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	341	Total	C	N	O	S	0	5	0
			2647	1645	461	520	21			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	565	LYS	ASN	SEE REMARK 999	UNP P24627
A	608	GLU	LYS	SEE REMARK 999	UNP P24627

- Molecule 2 is a polymer of unknown type called SUGAR (5-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	5	Total	C	N	O	0	0
			61	34	2	25		


- Molecule 3 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	2	Total	C	N	O	0	0
			28	16	2	10		

- Molecule 4 is a polymer of unknown type called SUGAR (6-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	6	Total	C	N	O	0	0
			72	40	2	30		

- Molecule 5 is SUGAR (CELLOBIOSE) (three-letter code: CBI) (formula: C₁₂H₂₂O₁₁).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			23	12	11		

- Molecule 6 is FE (III) ION (three-letter code: FE) (formula: Fe).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Fe	0	0
			1	1		

- Molecule 7 is CARBONATE ION (three-letter code: CO3) (formula: CO₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			4	1	3		

- Molecule 8 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	2	Total	Zn	0	0
			2	2		

- Molecule 9 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 10 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	300	Total	O	0	0
			300	300		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: Lactotransferrin



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	63.47Å 50.44Å 65.93Å 90.00° 107.78° 90.00°	Depositor
Resolution (Å)	20.00 – 2.20	Depositor
% Data completeness (in resolution range)	99.5 (20.00-2.20)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	0.14	Depositor
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.184 , 0.221	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	3143	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, BMA, NAG, CBI, CO3, SO4, MAN, FE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.37	0/2695	0.65	0/3647

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	342	TYR	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2647	0	2547	52	0
2	A	61	0	52	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	28	0	25	0	0
4	A	72	0	61	2	0
5	A	23	0	22	1	0
6	A	1	0	0	0	0
7	A	4	0	0	0	0
8	A	2	0	0	0	0
9	A	5	0	0	0	0
10	A	300	0	0	10	0
All	All	3143	0	2707	55	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (55) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:419:LYS:HD3	1:A:420:HIS:H	1.17	1.07
4:A:691:BMA:H3	4:A:692:BMA:O5	1.56	1.06
1:A:343:THR:HA	1:A:606:HIS:NE2	1.70	1.05
1:A:416:LYS:HG2	10:A:1226:HOH:O	1.66	0.94
1:A:565:LYS:HD2	1:A:567:GLU:H	1.35	0.91
1:A:565:LYS:CD	1:A:567:GLU:H	1.91	0.84
1:A:419:LYS:HD3	1:A:420:HIS:N	1.96	0.76
1:A:565:LYS:HD3	1:A:566:ARG:N	2.03	0.74
1:A:625:CYS:O	1:A:630:CYS:SG	2.48	0.71
2:A:2:NAG:O3	2:A:3:BMA:H2	1.92	0.69
1:A:428:ARG:HD2	10:A:1162:HOH:O	1.92	0.69
1:A:635:GLU:O	1:A:636:THR:HG22	1.97	0.64
1:A:638:ASN:HD22	1:A:643:ASP:H	1.48	0.60
1:A:663:THR:OG1	5:A:695:CBI:H1	2.02	0.59
1:A:625:CYS:C	1:A:630:CYS:SG	2.82	0.58
1:A:635:GLU:HG2	1:A:637[B]:LYS:NZ	2.19	0.58
1:A:565:LYS:HD2	1:A:567:GLU:N	2.11	0.58
1:A:460:ALA:HB3	1:A:463:ARG:HD3	1.86	0.57
1:A:471:MET:HE2	1:A:474:ILE:HD12	1.86	0.57
4:A:691:BMA:C3	4:A:692:BMA:O5	2.44	0.56
1:A:565:LYS:HD2	1:A:567:GLU:HB2	1.89	0.54
1:A:585[B]:GLN:NE2	10:A:1225:HOH:O	2.42	0.53
1:A:625:CYS:HA	1:A:629:PHE:O	2.10	0.51
1:A:430:THR:HB	1:A:594:ASN:ND2	2.26	0.51
1:A:638:ASN:ND2	1:A:643:ASP:H	2.08	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:344:ARG:NH1	10:A:1075:HOH:O	2.45	0.49
1:A:471:MET:CE	1:A:474:ILE:HD12	2.43	0.49
1:A:565:LYS:C	1:A:565:LYS:HD3	2.34	0.48
1:A:461:VAL:O	1:A:462:ASP:HB2	2.15	0.47
1:A:452:LYS:O	1:A:453:ASP:HB2	2.14	0.47
1:A:658:GLU:CD	10:A:1158:HOH:O	2.53	0.47
1:A:635:GLU:CG	1:A:637[B]:LYS:HZ2	2.28	0.46
1:A:415:ARG:HD3	10:A:1080:HOH:O	2.17	0.45
1:A:636:THR:HG23	1:A:636:THR:O	2.16	0.45
1:A:395:ASP:HA	1:A:595:HIS:CD2	2.52	0.45
1:A:669:ILE:O	1:A:673:LYS:HG2	2.16	0.45
1:A:446:LEU:HD11	1:A:451:LEU:HD23	1.99	0.44
1:A:565:LYS:HE3	1:A:567:GLU:HB2	2.00	0.44
1:A:471:MET:HE1	1:A:487:PHE:HE2	1.82	0.44
1:A:682:GLU:O	1:A:683:ALA:HB2	2.19	0.43
1:A:665:TYR:CZ	1:A:669:ILE:HD11	2.54	0.43
1:A:603:ARG:HD2	10:A:1123:HOH:O	2.18	0.43
1:A:402:ALA:HB1	1:A:407:LEU:HD12	2.01	0.42
1:A:514:LYS:HD2	10:A:1129:HOH:O	2.19	0.42
1:A:653:GLY:O	1:A:654:ARG:C	2.58	0.42
1:A:416:LYS:HG2	1:A:416:LYS:H	1.72	0.42
1:A:633:LYS:HA	1:A:633:LYS:HD3	1.71	0.42
1:A:636:THR:HG21	10:A:1072:HOH:O	2.20	0.41
1:A:471:MET:HA	1:A:471:MET:HE2	2.02	0.41
1:A:585[B]:GLN:HE21	1:A:585[B]:GLN:HB2	1.71	0.41
1:A:635:GLU:HG3	10:A:1247:HOH:O	2.21	0.41
1:A:665:TYR:CE2	1:A:669:ILE:HD11	2.56	0.41
1:A:663:THR:O	1:A:667:THR:HG23	2.21	0.41
1:A:455:LYS:HB3	1:A:504:LEU:HD21	2.03	0.40
1:A:531:ARG:HG3	1:A:560:TRP:CD2	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	342/345 (99%)	324 (95%)	16 (5%)	2 (1%)	28	29

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	634	SER
1	A	652	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	287/286 (100%)	269 (94%)	18 (6%)	21	23

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	343	THR
1	A	414	ASN
1	A	416	LYS
1	A	419	LYS
1	A	498	LYS
1	A	500	ARG
1	A	504	LEU
1	A	515	CYS
1	A	559[A]	ASP
1	A	559[B]	ASP
1	A	565	LYS
1	A	567	GLU
1	A	583	GLU
1	A	591	VAL
1	A	611	LEU
1	A	631	LEU
1	A	636	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	664	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	355	GLN
1	A	414	ASN
1	A	621	ASN
1	A	638	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

13 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	A	1	1,2	14,14,15	0.52	0	15,19,21	0.71	0
2	NAG	A	2	2	14,14,15	0.56	0	15,19,21	1.40	1 (6%)
2	BMA	A	3	2	11,11,12	0.61	0	13,15,17	1.04	2 (15%)
2	MAN	A	4	2	11,11,12	0.57	0	13,15,17	1.25	2 (15%)
2	BMA	A	5	2	11,11,12	0.61	0	13,15,17	0.56	0
3	NAG	A	687	1,3	14,14,15	0.55	0	15,19,21	1.19	2 (13%)
3	NAG	A	688	3	14,14,15	0.48	0	15,19,21	1.25	1 (6%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	NAG	A	689	1,4	14,14,15	0.52	0	15,19,21	1.60	2 (13%)
4	NAG	A	690	4	14,14,15	0.52	0	15,19,21	0.86	0
4	BMA	A	691	4	11,11,12	0.76	0	13,15,17	1.90	2 (15%)
4	BMA	A	692	4	11,11,12	0.54	0	13,15,17	1.08	1 (7%)
4	MAN	A	693	4	11,11,12	0.57	0	13,15,17	1.04	1 (7%)
4	BMA	A	694	4	11,11,12	0.59	0	13,15,17	0.81	1 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	A	2	2	-	0/6/23/26	0/1/1/1
2	BMA	A	3	2	-	0/2/19/22	0/1/1/1
2	MAN	A	4	2	-	0/2/19/22	0/1/1/1
2	BMA	A	5	2	-	0/2/19/22	0/1/1/1
3	NAG	A	687	1,3	-	0/6/23/26	0/1/1/1
3	NAG	A	688	3	-	0/6/23/26	0/1/1/1
4	NAG	A	689	1,4	-	0/6/23/26	0/1/1/1
4	NAG	A	690	4	-	0/6/23/26	0/1/1/1
4	BMA	A	691	4	-	0/2/19/22	0/1/1/1
4	BMA	A	692	4	-	0/2/19/22	0/1/1/1
4	MAN	A	693	4	-	0/2/19/22	0/1/1/1
4	BMA	A	694	4	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	687	NAG	O5-C1-C2	-2.95	107.37	111.47
2	A	3	BMA	O5-C1-C2	-2.27	107.24	110.79
4	A	692	BMA	C3-C4-C5	-2.17	106.40	110.22
4	A	689	NAG	O5-C1-C2	-2.07	108.59	111.47
2	A	3	BMA	C3-C4-C5	2.13	113.97	110.22
3	A	687	NAG	C1-O5-C5	2.15	115.13	112.17
2	A	4	MAN	C1-O5-C5	2.31	115.36	112.17
4	A	694	BMA	C1-C2-C3	2.45	112.76	109.65
4	A	693	MAN	C1-C2-C3	2.71	113.09	109.65
4	A	691	BMA	C2-C3-C4	3.14	116.36	110.88

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	688	NAG	C2-N2-C7	3.19	127.59	122.94
2	A	4	MAN	C1-C2-C3	3.26	113.78	109.65
2	A	2	NAG	C4-C3-C2	3.48	116.12	111.02
4	A	689	NAG	C1-O5-C5	5.25	119.41	112.17
4	A	691	BMA	C1-C2-C3	5.34	116.42	109.65

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	2	NAG	1	0
2	A	3	BMA	1	0
4	A	691	BMA	2	0
4	A	692	BMA	2	0

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 3 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
7	CO3	A	1002	6	0,3,3	0.00	-	0,3,3	0.00	-
9	SO4	A	1005	-	4,4,4	0.10	0	6,6,6	0.21	0
5	CBI	A	695	-	24,24,24	0.40	0	35,35,35	1.26	4 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	CO3	A	1002	6	-	0/0/0/0	0/0/0/0
9	SO4	A	1005	-	-	0/0/0/0	0/0/0/0
5	CBI	A	695	-	-	0/8/48/48	0/2/2/2

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	695	CBI	C1-O4'-C4'	-3.19	110.22	118.00
5	A	695	CBI	C1-O5-C5	-2.34	109.31	113.72
5	A	695	CBI	C4-C3-C2	2.04	114.43	110.84
5	A	695	CBI	O4'-C1-C2	4.43	118.10	108.11

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	695	CBI	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.