



Full wwPDB X-ray Structure Validation Report ⓘ

Jul 19, 2017 – 07:14 PM EDT

PDB ID : 4DXB
Title : 2.29A structure of the engineered MBP TEM-1 fusion protein RG13 in complex with zinc, P1 space group
Authors : van den Akker, F.; Ke, W.
Deposited on : unknown
Resolution : 2.29 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20029824
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20029824


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.29 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	4130 (2.30-2.30)
Clashscore	112137	4751 (2.30-2.30)
Ramachandran outliers	110173	4705 (2.30-2.30)
Sidechain outliers	110143	4704 (2.30-2.30)
RSRZ outliers	101464	4156 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	637	<div> <div style="width: 100%; height: 10px; background-color: red;"></div> <div style="width: 91%; height: 10px; background-color: green;"></div> <div style="width: 8%; height: 10px; background-color: yellow;"></div> <div style="width: 1%; height: 10px; background-color: red;"></div> <div style="width: 0%; height: 10px; background-color: grey;"></div> <div style="display: flex; justify-content: space-between; font-size: 0.8em;"> 0% 91% 8% </div> </div>
1	B	637	<div> <div style="width: 5%; height: 10px; background-color: red;"></div> <div style="width: 87%; height: 10px; background-color: green;"></div> <div style="width: 13%; height: 10px; background-color: yellow;"></div> <div style="width: 0%; height: 10px; background-color: red;"></div> <div style="width: 0%; height: 10px; background-color: grey;"></div> <div style="display: flex; justify-content: space-between; font-size: 0.8em;"> 5% 87% 13% </div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 10345 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Maltose-binding periplasmic protein, Beta-lactamase TEM chimera.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	637	Total	C	N	O	S	0	1	0
			4927	3129	833	948	17			
1	B	636	Total	C	N	O	S	0	2	0
			4924	3129	832	946	17			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	377	GLY	-	LINKER	UNP P62593
A	378	SER	-	LINKER	UNP P62593
A	379	GLY	-	LINKER	UNP P62593
A	380	GLY	-	LINKER	UNP P62593
A	381	GLY	-	LINKER	UNP P62593
A	585	SER	-	LINKER	UNP P0AEX9
B	377	GLY	-	LINKER	UNP P62593
B	378	SER	-	LINKER	UNP P62593
B	379	GLY	-	LINKER	UNP P62593
B	380	GLY	-	LINKER	UNP P62593
B	381	GLY	-	LINKER	UNP P62593
B	585	SER	-	LINKER	UNP P0AEX9

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	2	Total	Zn	0	0
			2	2		
2	A	2	Total	Zn	0	0
			2	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	260	Total 260	O 260	0	0
3	B	230	Total 230	O 230	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Maltose-binding periplasmic protein, Beta-lactamase TEM chimera



- Molecule 1: Maltose-binding periplasmic protein, Beta-lactamase TEM chimera



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	48.32Å 74.18Å 103.13Å 83.54° 77.64° 89.98°	Depositor
Resolution (Å)	31.45 – 2.29 31.45 – 2.29	Depositor EDS
% Data completeness (in resolution range)	95.2 (31.45-2.29) 86.9 (31.45-2.29)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.21 (at 2.29Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.232 , 0.292 0.227 , 0.285	Depositor DCC
R_{free} test set	3008 reflections (5.36%)	DCC
Wilson B-factor (Å ²)	32.2	Xtriage
Anisotropy	0.068	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 12.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.169 for h,-k,h-l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	10345	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.46% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.35	0/5034	0.49	0/6824
1	B	0.35	0/5034	0.49	0/6825
All	All	0.35	0/10068	0.49	0/13649

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4927	0	4910	29	0
1	B	4924	0	4911	48	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	A	260	0	0	0	0
3	B	230	0	0	2	0
All	All	10345	0	9821	77	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (77) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:418:PRO:HB2	1:B:540:VAL:HG23	1.60	0.83
1:B:184:ASP:OD1	1:B:629:LYS:HA	1.96	0.65
1:A:128:THR:HG22	1:A:130:GLU:H	1.65	0.61
1:B:418:PRO:HB2	1:B:540:VAL:CG2	2.31	0.60
1:B:575:PRO:HA	1:B:578:ARG:HG3	1.83	0.60
1:A:518:LEU:HD11	1:A:525:LEU:HD22	1.84	0.59
1:B:62:TRP:HB3	1:B:67:PHE:HE1	1.66	0.58
1:B:79:ILE:HG22	1:B:81:PRO:HD3	1.85	0.58
1:A:333:ILE:HD11	1:A:575:PRO:HD2	1.85	0.57
1:A:474:THR:HG22	1:A:477:GLU:H	1.73	0.53
1:B:148:MET:HG2	1:B:213:ALA:HA	1.92	0.52
1:B:70:TYR:HA	1:B:75:LEU:HD12	1.92	0.52
1:A:192:LEU:HD12	1:A:195:LEU:HD23	1.92	0.52
1:B:8:VAL:HG13	1:B:58:ASP:H	1.75	0.52
1:B:99:TYR:CB	1:B:104:ILE:HD11	2.40	0.51
1:B:625:ASP:O	1:B:629:LYS:HB2	2.10	0.51
1:B:372:LEU:O	1:B:376:TRP:HD1	1.93	0.51
1:A:349:ILE:HG12	1:A:402:TYR:HD1	1.75	0.51
1:A:372:LEU:O	1:A:376:TRP:HD1	1.94	0.50
1:B:474:THR:HG22	1:B:477:GLU:H	1.77	0.50
1:B:99:TYR:HB3	1:B:104:ILE:HD11	1.94	0.50
1:B:262:LEU:HB2	3:B:921:HOH:O	2.11	0.49
1:A:192:LEU:HD23	1:A:624:VAL:HG13	1.94	0.49
1:B:357:THR:HG22	1:B:360:GLU:H	1.77	0.49
1:A:337:LEU:HD23	1:A:571:LYS:HD2	1.94	0.49
1:A:462:SER:HB3	1:A:465:THR:OG1	2.13	0.49
1:A:439:ARG:NH1	1:A:497:THR:O	2.46	0.48
1:A:447:LEU:O	1:A:474:THR:HG23	2.14	0.48
1:B:62:TRP:HB3	1:B:67:PHE:CE1	2.48	0.48
1:A:382:HIS:HD2	1:A:384:GLU:HB2	1.79	0.48
1:B:291:GLU:O	1:B:295:LYS:HB2	2.15	0.47
1:A:333:ILE:HD12	1:A:365:ILE:HG21	1.96	0.47
1:B:287:ASP:O	1:B:291:GLU:HB2	2.14	0.47
1:B:218:ASN:HD22	1:B:218:ASN:N	2.13	0.47
1:B:97:VAL:HG21	1:B:105:ALA:HB3	1.97	0.47
1:B:142[A]:LYS:HA	1:B:142[A]:LYS:NZ	2.31	0.46
1:A:43:LEU:HA	1:A:46:LYS:HB2	1.97	0.46
1:A:46:LYS:O	1:A:50:VAL:HG22	2.16	0.46
1:A:218:ASN:HD21	1:A:235:ILE:HG12	1.79	0.46
1:B:518:LEU:HD11	1:B:525:LEU:HD22	1.98	0.46
1:B:437:LEU:HA	1:B:440:ILE:HG12	1.98	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:552:GLY:O	1:B:560:ARG:NH2	2.49	0.45
1:A:440:ILE:HG12	1:A:447:LEU:HG	1.99	0.45
1:B:151:LEU:HD12	1:B:205:ASN:O	2.17	0.45
1:B:383:PRO:O	1:B:387:VAL:HG23	2.17	0.45
1:B:8:VAL:HG13	1:B:57:PRO:HA	1.98	0.45
1:B:154:PRO:O	1:B:158:TRP:HB2	2.18	0.44
1:B:348:VAL:HB	1:B:403:ILE:HG22	1.98	0.44
1:A:349:ILE:HG12	1:A:402:TYR:CD1	2.52	0.44
1:B:46:LYS:O	1:B:50:VAL:HG22	2.18	0.44
1:A:334:ILE:HG22	1:A:348:VAL:HG22	1.98	0.44
1:A:139:LEU:O	1:A:142:LYS:HG2	2.18	0.44
1:B:110:VAL:HB	1:B:301:ALA:HB3	2.00	0.44
1:B:10:TRP:CE2	1:B:40:PRO:HG2	2.53	0.43
1:B:79:ILE:HG21	1:B:106:TYR:CZ	2.54	0.43
1:B:386:LEU:HD22	1:B:390:LYS:HE2	2.00	0.43
1:B:462:SER:HB3	1:B:465:THR:OG1	2.19	0.43
1:A:327:GLU:HG3	1:A:331:ARG:NH1	2.34	0.42
1:B:432:LEU:HD21	1:B:494:LEU:HB2	2.01	0.42
1:B:338:GLY:HA3	1:B:343:PRO:HA	2.01	0.42
1:B:447:LEU:HB3	1:B:476:ARG:HB3	2.02	0.42
1:B:334:ILE:HG22	1:B:348:VAL:HG22	2.01	0.42
1:A:382:HIS:CD2	1:A:384:GLU:HB2	2.55	0.42
1:B:8:VAL:HG12	1:B:58:ASP:OD2	2.20	0.42
1:A:370:ALA:HA	1:A:580:ALA:HB2	2.02	0.41
1:B:439:ARG:NH1	3:B:929:HOH:O	2.53	0.41
1:A:90:TYR:HA	1:A:91:PRO:HD3	1.94	0.41
1:B:44:GLU:HB3	1:B:66:ARG:HH21	1.85	0.41
1:A:534:ARG:O	1:A:536:THR:HG23	2.21	0.41
1:A:331:ARG:HA	1:A:351:THR:OG1	2.21	0.41
1:B:418:PRO:CB	1:B:540:VAL:HG23	2.41	0.41
1:A:314:ASP:HA	1:A:315:PRO:HD3	1.95	0.41
1:B:597:MET:HA	1:B:598:PRO:HD3	1.91	0.41
1:B:540:VAL:O	1:B:544:THR:HG23	2.20	0.41
1:B:79:ILE:HG21	1:B:106:TYR:CE2	2.56	0.41
1:A:365:ILE:HG22	1:A:576:LEU:HA	2.03	0.40
1:B:43:LEU:HA	1:B:46:LYS:HB2	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	636/637 (100%)	619 (97%)	16 (2%)	1 (0%)	51	63
1	B	636/637 (100%)	615 (97%)	20 (3%)	1 (0%)	51	63
All	All	1272/1274 (100%)	1234 (97%)	36 (3%)	2 (0%)	51	63

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	173	ASN
1	B	530	PRO

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	517/516 (100%)	506 (98%)	11 (2%)	59	76
1	B	517/516 (100%)	502 (97%)	15 (3%)	48	64
All	All	1034/1032 (100%)	1008 (98%)	26 (2%)	54	70

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	LYS
1	A	44	GLU
1	A	189	LYS
1	A	253	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	317	TRP
1	A	333	ILE
1	A	384	GLU
1	A	474	THR
1	A	510	ASN
1	A	546	LEU
1	A	551	THR
1	B	4	GLU
1	B	45	GLU
1	B	49	GLN
1	B	60	ILE
1	B	119	LYS
1	B	142[A]	LYS
1	B	142[B]	LYS
1	B	202	LYS
1	B	317	TRP
1	B	321	ASP
1	B	357	THR
1	B	386	LEU
1	B	474	THR
1	B	485	MET
1	B	502	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	18	ASN
1	A	201	ASN
1	A	218	ASN
1	A	375	HIS
1	A	382	HIS
1	A	592	GLN
1	B	72	GLN
1	B	201	ASN
1	B	218	ASN
1	B	375	HIS
1	B	514	HIS
1	B	592	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	637/637 (100%)	0.07	6 (0%) 84 87	10, 30, 51, 67	0
1	B	636/637 (99%)	0.38	33 (5%) 28 35	12, 35, 61, 74	0
All	All	1273/1274 (99%)	0.23	39 (3%) 49 56	10, 32, 58, 74	0

All (39) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	340	ASP	4.9
1	B	320	ALA	4.4
1	B	275	LEU	4.0
1	A	322	LYS	3.6
1	B	70	TYR	3.5
1	B	2	ILE	3.4
1	B	71	ALA	3.4
1	B	46	LYS	3.3
1	B	47	PHE	3.2
1	B	97	VAL	3.2
1	B	608	TYR	3.2
1	B	340	ASP	3.1
1	B	80	THR	3.1
1	B	322	LYS	3.0
1	B	55	ASP	2.9
1	B	379	GLY	2.9
1	B	75	LEU	2.8
1	B	377	GLY	2.8
1	B	105	ALA	2.7
1	B	633	THR	2.7
1	A	637	LYS	2.7
1	B	4	GLU	2.7
1	B	376	TRP	2.6
1	B	540	VAL	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	580	ALA	2.5
1	A	342	LYS	2.5
1	B	76	LEU	2.4
1	B	265	GLY	2.4
1	B	77	ALA	2.3
1	B	267	ASN	2.3
1	A	376	TRP	2.3
1	B	100	ASN	2.2
1	B	98	ARG	2.2
1	A	143	GLY	2.1
1	B	106	TYR	2.1
1	B	53	THR	2.1
1	B	579	SER	2.1
1	B	56	GLY	2.0
1	B	173	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	ZN	A	701	1/1	0.99	0.11	-0.50	37,37,37,37	0
2	ZN	B	701	1/1	0.99	0.10	-	36,36,36,36	0
2	ZN	A	702	1/1	0.99	0.10	-	24,24,24,24	0
2	ZN	B	702	1/1	1.00	0.07	-	27,27,27,27	0

6.5 Other polymers [i](#)

There are no such residues in this entry.