



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 02:02 pm GMT

PDB ID : 5E19  
Title : Crystal Structure of the ER-alpha Ligand-binding Domain in Complex with the Cyclofenil Derivative methyl {4-[bis(4-hydroxyphenyl)methylidene]cyclohexyl}acetate  
Authors : Nwachukwu, J.C.; Srinivasan, S.; Zheng, Y.; Wang, S.; Min, J.; Dong, C.; Liao, Z.; Cavett, V.; Nowak, J.; Houtman, R.; Carlson, K.E.; Josan, J.S.; Elemento, O.; Katzenellenbogen, J.A.; Zhou, H.B.; Nettles, K.W.  
Deposited on : 2015-09-29  
Resolution : 2.24 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.24 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	1804 (2.26-2.22)
Clashscore	112137	1957 (2.26-2.22)
Ramachandran outliers	110173	1916 (2.26-2.22)
Sidechain outliers	110143	1917 (2.26-2.22)
RSRZ outliers	101464	1809 (2.26-2.22)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	257	<div> <div></div> <div> <div></div> <div>81%</div> <div>9%</div> <div>•</div> <div>9%</div> </div> </div>
1	B	257	<div> <div></div> <div> <div></div> <div>86%</div> <div>7%</div> <div>7%</div> </div> </div>
2	C	14	<div> <div></div> <div> <div></div> <div>50%</div> <div>14%</div> <div>36%</div> </div> </div>
2	D	14	<div> <div></div> <div> <div></div> <div>57%</div> <div>14%</div> <div>29%</div> </div> </div>

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3974 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Estrogen receptor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	233	Total	C	N	O	S	0	2	0
			1817	1162	311	325	19			
1	B	238	Total	C	N	O	S	0	0	0
			1830	1167	312	333	18			


There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	537	SER	TYR	engineered mutation	UNP P03372
B	537	SER	TYR	engineered mutation	UNP P03372

- Molecule 2 is a protein called Nuclear receptor coactivator 2.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	9	Total	C	N	O	0	0	0
			75	48	15	12			
2	D	10	Total	C	N	O	0	0	0
			84	54	17	13			

- Molecule 3 is methyl {4-[bis(4-hydroxyphenyl)methylidene]cyclohexyl}acetate (three-letter code: 5K7) (formula: C<sub>22</sub>H<sub>24</sub>O<sub>4</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			26	22	4		
3	B	1	Total	C	O	0	0
			26	22	4		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	52	Total	O	0	0
			52	52		
4	B	61	Total	O	0	0
			61	61		
4	C	2	Total	O	0	0
			2	2		
4	D	1	Total	O	0	0
			1	1		



- Molecule 1: Estrogen receptor



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	55.37Å 82.00Å 58.48Å 90.00° 110.83° 90.00°	Depositor
Resolution (Å)	46.79 – 2.24 46.79 – 2.23	Depositor EDS
% Data completeness (in resolution range)	90.4 (46.79-2.24) 90.1 (46.79-2.23)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.82 (at 2.22Å)	Xtriage
Refinement program	PHENIX	Depositor
R, $R_{free}$	0.194 , 0.233 0.194 , 0.230	Depositor DCC
$R_{free}$ test set	1840 reflections (8.62%)	DCC
Wilson B-factor (Å <sup>2</sup> )	36.3	Xtriage
Anisotropy	0.554	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 47.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	3974	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	56.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.57% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 5K7

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.21	0/1857	0.37	0/2519
1	B	0.24	0/1863	0.37	0/2528
2	C	0.17	0/75	0.32	0/100
2	D	0.18	0/84	0.35	0/111
All	All	0.22	0/3879	0.37	0/5258

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	332	ASP	Peptide

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1817	0	1821	15	0
1	B	1830	0	1816	10	0
2	C	75	0	77	2	0
2	D	84	0	90	2	0
3	A	26	0	0	0	0
3	B	26	0	0	0	0
4	A	52	0	0	3	0
4	B	61	0	0	1	0
4	C	2	0	0	1	0
4	D	1	0	0	0	0
All	All	3974	0	3804	25	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (25) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:334:THR:O	4:A:1001:HOH:O	2.06	0.73
1:A:385:GLU:OE1	4:A:1002:HOH:O	2.09	0.70
1:B:468:SER:N	4:B:1003:HOH:O	2.28	0.66
1:A:376:VAL:HG13	2:D:690:LEU:HD23	1.81	0.61
1:A:396:MET:O	1:A:436[B]:ARG:NH1	2.26	0.61
1:B:531:LYS:O	1:B:533:VAL:N	2.34	0.59
1:A:501:HIS:ND1	4:A:1007:HOH:O	2.32	0.58
2:C:694:LEU:O	4:C:701:HOH:O	2.18	0.55
1:B:487:ILE:HD11	1:B:504:LEU:HD22	1.89	0.55
1:A:533:VAL:HG22	1:A:534:VAL:HG23	1.90	0.53
1:B:376:VAL:HG13	2:C:690:LEU:HD23	1.91	0.51
1:A:498:GLN:HA	1:A:501:HIS:CE1	2.46	0.50
1:A:351:ASP:OD1	1:A:537:SER:OG	2.24	0.49
1:B:539:LEU:O	1:B:543:MET:HG2	2.15	0.46
1:A:376:VAL:HG21	2:D:691:HIS:CE1	2.51	0.46
1:A:539:LEU:O	1:A:543:MET:HG2	2.19	0.42
1:A:331:TYR:O	1:A:333:PRO:HD3	2.20	0.42
1:A:339:GLU:HG3	1:A:418:VAL:HG22	2.02	0.42
1:B:531:LYS:C	1:B:533:VAL:H	2.21	0.42
1:A:458:VAL:HG13	1:A:459:TYR:CD2	2.56	0.41
1:B:382:ALA:HB2	1:B:456:SER:HB2	2.02	0.41
1:A:337:PHE:HA	1:A:417:CYS:SG	2.60	0.41
1:A:455:ASN:O	1:B:513:HIS:NE2	2.53	0.41
1:B:534:VAL:HA	1:B:535:PRO:HD3	1.86	0.41

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:354:LEU:HA	1:B:354:LEU:HD23	1.94	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	231/257 (90%)	224 (97%)	4 (2%)	3 (1%)	14	9
1	B	234/257 (91%)	230 (98%)	3 (1%)	1 (0%)	38	40
2	C	7/14 (50%)	7 (100%)	0	0	100	100
2	D	8/14 (57%)	8 (100%)	0	0	100	100
All	All	480/542 (89%)	469 (98%)	7 (2%)	4 (1%)	22	19

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	332	ASP
1	A	333	PRO
1	A	336	PRO
1	B	532	ASN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	199/232 (86%)	197 (99%)	2 (1%)	80	86
1	B	199/232 (86%)	197 (99%)	2 (1%)	80	86
2	C	8/14 (57%)	8 (100%)	0	100	100
2	D	9/14 (64%)	9 (100%)	0	100	100
All	All	415/492 (84%)	411 (99%)	4 (1%)	80	86

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	338	SER
1	A	533	VAL
1	B	417	CYS
1	B	469	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	414	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link

column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	5K7	A	901	-	28,28,28	1.73	3 (10%)	38,38,38	1.20	5 (13%)
3	5K7	B	900	-	28,28,28	1.67	4 (14%)	38,38,38	1.57	5 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	5K7	A	901	-	-	0/18/28/28	0/3/3/3
3	5K7	B	900	-	-	0/18/28/28	0/3/3/3

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	900	5K7	O20-C21	-2.09	1.40	1.45
3	B	900	5K7	C18-C19	2.08	1.55	1.50
3	A	901	5K7	O20-C19	2.29	1.40	1.32
3	B	900	5K7	C07-C06	3.88	1.55	1.49
3	A	901	5K7	C07-C06	4.16	1.56	1.49
3	B	900	5K7	C14-C06	4.82	1.38	1.34
3	A	901	5K7	C14-C06	6.50	1.40	1.34

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	900	5K7	C24-C14-C06	-4.99	119.12	124.18
3	B	900	5K7	C07-C06-C14	-3.69	118.29	122.71
3	A	901	5K7	C24-C14-C06	-3.46	120.67	124.18
3	B	900	5K7	C23-C17-C18	-3.07	107.79	111.48
3	A	901	5K7	C07-C06-C14	-2.94	119.19	122.71
3	A	901	5K7	C16-C17-C18	-2.66	108.28	111.48
3	B	900	5K7	O20-C19-O22	-2.37	115.27	123.12
3	A	901	5K7	O22-C19-C18	-2.12	119.92	124.70
3	A	901	5K7	O20-C19-C18	3.12	118.76	111.47
3	B	900	5K7	O20-C19-C18	4.14	121.15	111.47

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	233/257 (90%)	-0.26	3 (1%) 77 78	27, 52, 88, 115	0
1	B	238/257 (92%)	-0.20	2 (0%) 86 86	25, 54, 97, 128	0
2	C	9/14 (64%)	-0.03	0 100 100	58, 68, 90, 96	0
2	D	10/14 (71%)	-0.04	0 100 100	54, 70, 86, 90	0
All	All	490/542 (90%)	-0.22	5 (1%) 82 83	25, 53, 93, 128	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	533	VAL	5.7
1	B	305	SER	3.0
1	A	334	THR	2.9
1	A	337	PHE	2.8
1	A	526	TYR	2.7

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains.

The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	5K7	B	900	26/26	0.91	0.15	0.43	32,39,65,68	0
3	5K7	A	901	26/26	0.96	0.12	-0.08	30,34,50,56	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.