



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 15, 2017 – 11:30 PM EDT

PDB ID : 3E29
Title : X-Ray structure of the protein Q7WE92_BORBR from thioesterase super-family. Northeast Structural Genomics Consortium Target BoR214A.
Authors : Kuzin, A.P.; Chen, Y.; Setharaman, J.; Wang, D.; Mao, L.; Foote, E.L.; Xiao, R.; Nair, R.; Everett, J.K.; Acton, T.B.; Rost, B.; Montelione, G.T.; Tong, L.; Hunt, J.; Northeast Structural Genomics Consortium (NESG)
Deposited on : unknown
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20030345
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20030345


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION





The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	112137	3674 (2.40-2.40)
Ramachandran outliers	110173	3616 (2.40-2.40)
Sidechain outliers	110143	3617 (2.40-2.40)
RSRZ outliers	101464	3195 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	144	 73% 18% • 7%
1	B	144	 75% 17% •• 6%
1	C	144	 77% 15% • 8%
1	D	144	 79% 11% • 8%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 4321 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called uncharacterized protein Q7WE92_BORBR.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	134	Total	C	N	O	Se	8	2	0
			1043	656	195	186	6			
1	B	135	Total	C	N	O	Se	8	1	0
			1040	654	193	187	6			
1	C	133	Total	C	N	O	Se	8	2	0
			1035	651	194	185	5			
1	D	132	Total	C	N	O	Se	0	1	0
			1021	643	190	183	5			

There are 32 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	137	LEU	-	expression tag	UNP Q7WE92
A	138	GLU	-	expression tag	UNP Q7WE92
A	139	HIS	-	expression tag	UNP Q7WE92
A	140	HIS	-	expression tag	UNP Q7WE92
A	141	HIS	-	expression tag	UNP Q7WE92
A	142	HIS	-	expression tag	UNP Q7WE92
A	143	HIS	-	expression tag	UNP Q7WE92
A	144	HIS	-	expression tag	UNP Q7WE92
B	137	LEU	-	expression tag	UNP Q7WE92
B	138	GLU	-	expression tag	UNP Q7WE92
B	139	HIS	-	expression tag	UNP Q7WE92
B	140	HIS	-	expression tag	UNP Q7WE92
B	141	HIS	-	expression tag	UNP Q7WE92
B	142	HIS	-	expression tag	UNP Q7WE92
B	143	HIS	-	expression tag	UNP Q7WE92
B	144	HIS	-	expression tag	UNP Q7WE92
C	137	LEU	-	expression tag	UNP Q7WE92
C	138	GLU	-	expression tag	UNP Q7WE92
C	139	HIS	-	expression tag	UNP Q7WE92
C	140	HIS	-	expression tag	UNP Q7WE92
C	141	HIS	-	expression tag	UNP Q7WE92

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	142	HIS	-	expression tag	UNP Q7WE92
C	143	HIS	-	expression tag	UNP Q7WE92
C	144	HIS	-	expression tag	UNP Q7WE92
D	137	LEU	-	expression tag	UNP Q7WE92
D	138	GLU	-	expression tag	UNP Q7WE92
D	139	HIS	-	expression tag	UNP Q7WE92
D	140	HIS	-	expression tag	UNP Q7WE92
D	141	HIS	-	expression tag	UNP Q7WE92
D	142	HIS	-	expression tag	UNP Q7WE92
D	143	HIS	-	expression tag	UNP Q7WE92
D	144	HIS	-	expression tag	UNP Q7WE92


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	43	Total O 43 43	0	0
2	B	41	Total O 41 41	0	0
2	C	49	Total O 49 49	0	0
2	D	49	Total O 49 49	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: uncharacterized protein Q7WE92_BORBR




- Molecule 1: uncharacterized protein Q7WE92_BORBR



- Molecule 1: uncharacterized protein Q7WE92_BORBR



- Molecule 1: uncharacterized protein Q7WE92_BORBR



4 Data and refinement statistics

Property	Value	Source
Space group	P 61	Depositor
Cell constants a, b, c, α , β , γ	112.52Å 112.52Å 119.66Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	28.59 – 2.40 28.60 – 2.40	Depositor EDS
% Data completeness (in resolution range)	100.0 (28.59-2.40) 99.3 (28.60-2.40)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	6.27 (at 2.39Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.221 , 0.258 0.222 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	28.6	Xtriage
Anisotropy	0.532	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 23.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.480 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	4321	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 57.13 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 2.4741e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.53	0/1065	0.69	1/1431 (0.1%)
1	B	0.54	0/1059	0.67	1/1424 (0.1%)
1	C	0.51	0/1057	0.62	0/1421
1	D	0.48	0/1040	0.64	0/1399
All	All	0.52	0/4221	0.66	2/5675 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	128	ARG	NE-CZ-NH2	-5.41	117.59	120.30
1	A	118	ASP	CB-CG-OD1	5.02	122.82	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1043	0	1054	27	1
1	B	1040	0	1047	19	4
1	C	1035	0	1042	11	0
1	D	1021	0	1025	9	0
2	A	43	0	0	0	0
2	B	41	0	0	2	0
2	C	49	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	D	49	0	0	1	3
All	All	4321	0	4168	62	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (62) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1:MSE:HB3	1:A:30:GLY:HA3	1.32	1.07
1:C:103:ILE:HG13	1:C:113:HIS:CE1	2.23	0.73
1:B:108:ARG:HH21	1:D:108:ARG:NH1	1.88	0.71
1:A:103:ILE:HG13	1:A:113:HIS:CE1	2.26	0.69
1:C:97:ARG:HD3	2:C:159:HOH:O	1.92	0.68
1:C:88:HIS:HE1	1:C:126:SER:OG	1.77	0.66
1:A:60:THR:OG1	1:B:56:GLY:HA3	1.97	0.64
1:D:88:HIS:HE1	1:D:126:SER:OG	1.79	0.64
1:A:108[A]:ARG:H	1:A:108[A]:ARG:HE	1.49	0.59
1:B:103:ILE:HG13	1:B:113:HIS:CE1	2.37	0.59
1:B:115:ARG:NH1	2:B:174:HOH:O	2.30	0.59
1:A:106:GLY:HA3	1:A:108[A]:ARG:NH2	2.19	0.58
1:A:108[A]:ARG:H	1:A:108[A]:ARG:NE	2.02	0.58
1:A:108[A]:ARG:HE	1:A:109:PHE:H	1.53	0.57
1:C:103:ILE:HG13	1:C:113:HIS:ND1	2.19	0.57
1:D:41:ARG:H	1:D:44:LEU:HD22	1.69	0.57
1:B:115:ARG:HD3	2:B:174:HOH:O	2.05	0.56
1:A:113:HIS:CD2	1:A:128:ARG:HB3	2.41	0.56
1:B:134:ARG:HG2	1:B:135:ALA:H	1.70	0.55
1:A:108[A]:ARG:HG2	1:A:109:PHE:CD2	2.42	0.55
1:B:108:ARG:HH21	1:D:108:ARG:HH11	1.56	0.53
1:B:92:THR:HB	1:B:93:PRO:HD2	1.90	0.53
1:C:41:ARG:H	1:C:44:LEU:HD22	1.74	0.53
1:B:88:HIS:HE1	1:B:126:SER:OG	1.91	0.53
1:D:46:SER:HB2	1:D:54:HIS:HA	1.92	0.52
1:A:1:MSE:HB3	1:A:30:GLY:CA	2.23	0.51
1:A:56:GLY:HA3	1:B:60:THR:OG1	2.11	0.50
1:A:83:MSE:HE1	1:A:85:VAL:HB	1.94	0.50
1:B:133:ILE:O	1:B:134:ARG:HB2	2.12	0.50
1:A:108[A]:ARG:CD	1:A:108[A]:ARG:H	2.27	0.48
1:B:74:THR:HB	1:B:76:HIS:CE1	2.50	0.47
1:C:46:SER:HB2	1:C:54:HIS:HA	1.97	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:81:MSE:HG3	1:A:130:LEU:HG	1.97	0.46
1:D:99:GLU:O	1:D:114:ALA:HA	2.15	0.46
1:B:113:HIS:CD2	1:B:128:ARG:HB3	2.50	0.45
1:A:83:MSE:CE	1:A:85:VAL:HB	2.46	0.45
1:B:92:THR:HB	1:B:93:PRO:CD	2.47	0.44
1:B:41:ARG:O	1:B:44:LEU:HB2	2.17	0.44
1:B:46:SER:HB2	1:B:54:HIS:HA	2.00	0.44
1:C:3:SER:O	1:C:7:GLU:HB2	2.18	0.44
1:C:81:MSE:HE3	1:C:132:LEU:HB2	1.99	0.43
1:A:88:HIS:HE1	1:A:126:SER:OG	2.01	0.43
1:A:106:GLY:HA3	1:A:108[A]:ARG:HH21	1.84	0.42
1:D:103:ILE:HG12	1:D:113:HIS:CE1	2.55	0.42
1:C:88:HIS:CE1	1:C:126:SER:OG	2.65	0.42
1:A:103:ILE:HA	1:A:103:ILE:HD13	1.86	0.42
1:A:7:GLU:HG3	1:A:8:MSE:N	2.34	0.42
1:B:24:MSE:HA	1:B:37:GLY:O	2.20	0.42
1:C:118:ASP:OD1	1:C:122:ASN:HB2	2.20	0.41
1:A:113:HIS:HD2	1:A:128:ARG:HB3	1.85	0.41
1:A:88:HIS:HD2	2:D:181:HOH:O	2.02	0.41
1:A:92:THR:HB	1:A:93:PRO:CD	2.50	0.41
1:A:108[A]:ARG:HG2	1:A:109:PHE:CE2	2.55	0.41
1:D:92:THR:HB	1:D:93:PRO:CD	2.51	0.41
1:B:134:ARG:HG2	1:B:135:ALA:N	2.35	0.40
1:A:24:MSE:HA	1:A:37:GLY:O	2.21	0.40
1:C:92:THR:HB	1:C:93:PRO:CD	2.52	0.40
1:D:63:ASP:HB2	1:D:83:MSE:SE	2.71	0.40
1:A:6:LEU:O	1:A:10:SER:HB2	2.22	0.40

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:51[A]:ARG:CZ	2:D:183:HOH:O[6_555]	1.60	0.60
1:B:51[A]:ARG:NH1	2:D:183:HOH:O[6_555]	1.81	0.39
1:B:51[A]:ARG:NH2	2:D:183:HOH:O[6_555]	1.98	0.22
1:A:7:GLU:OE2	1:B:11:ARG:NH1[3_444]	2.15	0.05

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	134/144 (93%)	133 (99%)	1 (1%)	0	100	100
1	B	134/144 (93%)	131 (98%)	2 (2%)	1 (1%)	25	37
1	C	133/144 (92%)	129 (97%)	3 (2%)	1 (1%)	22	33
1	D	131/144 (91%)	129 (98%)	2 (2%)	0	100	100
All	All	532/576 (92%)	522 (98%)	8 (2%)	2 (0%)	38	54

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	134	ARG
1	C	133	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	108/109 (99%)	100 (93%)	8 (7%)	16	25
1	B	107/109 (98%)	102 (95%)	5 (5%)	30	48
1	C	107/109 (98%)	102 (95%)	5 (5%)	30	48
1	D	105/109 (96%)	100 (95%)	5 (5%)	30	47
All	All	427/436 (98%)	404 (95%)	23 (5%)	27	41

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	7	GLU
1	A	32	GLN
1	A	44	LEU
1	A	49	GLU
1	A	107	LYS
1	A	108[A]	ARG
1	A	108[B]	ARG
1	A	128	ARG
1	B	1	MSE
1	B	44	LEU
1	B	107	LYS
1	B	124	VAL
1	B	128	ARG
1	C	6	LEU
1	C	44	LEU
1	C	49	GLU
1	C	117	LEU
1	C	128	ARG
1	D	6	LEU
1	D	41	ARG
1	D	44	LEU
1	D	47	SER
1	D	128	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	19	ASN
1	A	84	HIS
1	A	88	HIS
1	B	19	ASN
1	B	76	HIS
1	B	84	HIS
1	B	88	HIS
1	C	88	HIS
1	D	88	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	128/144 (88%)	-0.34	0 100 100	22, 28, 36, 46	1 (0%)
1	B	129/144 (89%)	-0.30	2 (1%) 72 70	22, 29, 38, 51	1 (0%)
1	C	128/144 (88%)	-0.30	0 100 100	23, 29, 41, 49	1 (0%)
1	D	127/144 (88%)	-0.26	1 (0%) 86 84	22, 29, 40, 47	1 (0%)
All	All	512/576 (88%)	-0.30	3 (0%) 89 87	22, 29, 40, 51	4 (0%)

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	134	ARG	4.0
1	B	2	SER	2.2
1	B	135	ALA	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.