



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 14, 2017 – 01:01 am GMT

PDB ID : 1E3H  
Title : SEMET DERIVATIVE OF STREPTOMYCES ANTIBIOTICUS PN-  
PASE/GPSI ENZYME  
Authors : Symmons, M.F.; Jones, G.H.; Luisi, B.F.  
Deposited on : 2000-06-15  
Resolution : 2.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	2542 (2.60-2.60)
Clashscore	112137	2895 (2.60-2.60)
Ramachandran outliers	110173	2848 (2.60-2.60)
Sidechain outliers	110143	2848 (2.60-2.60)
RSRZ outliers	101464	2550 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	757	<div> <div>4%</div> <div>61%</div> <div>18%</div> <div>•</div> <div>21%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	SO4	A	903	-	-	-	X
2	SO4	A	904	-	-	X	X

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	SO4	A	906	-	-	-	X

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4842 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GUANOSINE PENTAPHOSPHATE SYNTHETASE.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	601	Total	C	N	O	S	Se	0	1	0
			4506	2828	790	868	3	17			

There are 39 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	31	ARG	LYS	SEQUENCING AMBIGUITY	UNP Q53597
A	156	ILE	TYR	SEQUENCING AMBIGUITY	UNP Q53597
A	210	ILE	VAL	SEQUENCING AMBIGUITY	UNP Q53597
A	260	PHE	PRO	SEQUENCING AMBIGUITY	UNP Q53597
A	261	LEU	SER	SEQUENCING AMBIGUITY	UNP Q53597
A	262	ASP	SER	SEQUENCING AMBIGUITY	UNP Q53597
A	263	TYR	THR	SEQUENCING AMBIGUITY	UNP Q53597
A	264	GLN	THR	SEQUENCING AMBIGUITY	UNP Q53597
A	265	ASP	ARG	SEQUENCING AMBIGUITY	UNP Q53597
A	267	VAL	THR	SEQUENCING AMBIGUITY	UNP Q53597
A	268	LEU	THR	SEQUENCING AMBIGUITY	UNP Q53597
A	269	GLU	SER	SEQUENCING AMBIGUITY	UNP Q53597
A	323	ALA	PRO	SEQUENCING AMBIGUITY	UNP Q53597
A	324	LEU	TRP	SEQUENCING AMBIGUITY	UNP Q53597
A	325	THR	PRO	SEQUENCING AMBIGUITY	UNP Q53597
A	326	LYS	SER	SEQUENCING AMBIGUITY	UNP Q53597
A	328	LEU	SER	SEQUENCING AMBIGUITY	UNP Q53597
A	329	VAL	SER	SEQUENCING AMBIGUITY	UNP Q53597
A	330	ARG	ALA	SEQUENCING AMBIGUITY	UNP Q53597
A	335	ALA	LYS	SEQUENCING AMBIGUITY	UNP Q53597
A	409	TYR	ILE	SEQUENCING AMBIGUITY	UNP Q53597
A	-15	MSE	MET	MODIFIED RESIDUE	PDB 1E3H
A	-9	MSE	MET	MODIFIED RESIDUE	PDB 1E3H
A	1	MSE	MET	MODIFIED RESIDUE	UNP Q53597
A	46	MSE	MET	MODIFIED RESIDUE	UNP Q53597
A	74	MSE	MET	MODIFIED RESIDUE	UNP Q53597
A	125	MSE	MET	MODIFIED RESIDUE	UNP Q53597

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	184	MSE	MET	MODIFIED RESIDUE	UNP Q53597
A	199	MSE	MET	MODIFIED RESIDUE	UNP Q53597
A	200	MSE	MET	MODIFIED RESIDUE	UNP Q53597
A	382	MSE	MET	MODIFIED RESIDUE	UNP Q53597
A	385	MSE	MET	MODIFIED RESIDUE	UNP Q53597
A	401	MSE	MET	MODIFIED RESIDUE	UNP Q53597
A	464	MSE	MET	MODIFIED RESIDUE	UNP Q53597
A	472	MSE	MET	MODIFIED RESIDUE	UNP Q53597
A	490	MSE	MET	MODIFIED RESIDUE	UNP Q53597
A	519	MSE	MET	MODIFIED RESIDUE	UNP Q53597
A	565	MSE	MET	MODIFIED RESIDUE	UNP Q53597
A	566	MSE	MET	MODIFIED RESIDUE	UNP Q53597

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total O S 5 4 1	0	0
2	A	1	Total O S 5 4 1	0	0
2	A	1	Total O S 5 4 1	0	0
2	A	1	Total O S 5 4 1	0	0
2	A	1	Total O S 5 4 1	0	0

Continued on next page...

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	291	Total	O	0	0
			291	291		

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

[illegible]

## 4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	133.61Å 133.61Å 344.47Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	38.99 – 2.60 38.99 – 2.60	Depositor EDS
% Data completeness (in resolution range)	99.6 (38.99-2.60) 99.7 (38.99-2.60)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.09	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.71 (at 2.61Å)	Xtriage
Refinement program	CNS 0.9	Depositor
R, $R_{free}$	0.201 , 0.226 0.197 , 0.223	Depositor DCC
$R_{free}$ test set	1790 reflections (4.88%)	DCC
Wilson B-factor (Å <sup>2</sup> )	46.2	Xtriage
Anisotropy	0.040	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 51.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.008 for $-1/3^*h+1/3^*k+1/3^*l,-k,8/3^*h+4/3^*k+1/3^*l$ 0.017 for $-2/3^*h-1/3^*k-1/3^*l,-1/3^*h-2/3^*k+1/3^*l,-4/3^*h+4/3^*k+1/3^*l$ 0.012 for $-h,1/3^*h-1/3^*k-1/3^*l,-4/3^*h-8/3^*k+1/3^*l$	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	4842	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	49.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.01% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.33	0/4563	0.59	0/6159

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4506	0	4520	118	0
2	A	45	0	0	2	0
3	A	291	0	0	9	0
All	All	4842	0	4520	118	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

The worst 5 of 118 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:112:LYS:H	1:A:112:LYS:HD3	1.20	1.03

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:100:ARG:HH21	1:A:423:ARG:HD3	1.32	0.94
1:A:100:ARG:NH2	1:A:423:ARG:HD3	1.84	0.93
1:A:613:GLU:H	1:A:626:ALA:HB3	1.38	0.87
1:A:285:ILE:HB	1:A:291:ARG:HB2	1.61	0.83

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	596/757 (79%)	556 (93%)	35 (6%)	5 (1%)	22	44

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	459	ASN
1	A	630	PRO
1	A	461	SER
1	A	626	ALA
1	A	625	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	462/577 (80%)	451 (98%)	11 (2%)	54 80

5 of 11 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	132	LEU
1	A	152	PHE
1	A	289	GLN
1	A	112	LYS
1	A	276	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	6	HIS
1	A	381	ASN
1	A	495	GLN
1	A	502	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected

value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	SO4	A	901	-	4,4,4	0.34	0	6,6,6	0.10	0
2	SO4	A	902	-	4,4,4	0.35	0	6,6,6	0.07	0
2	SO4	A	903	-	4,4,4	0.43	0	6,6,6	0.11	0
2	SO4	A	904	-	4,4,4	0.37	0	6,6,6	0.09	0
2	SO4	A	905	-	4,4,4	0.35	0	6,6,6	0.09	0
2	SO4	A	906	-	4,4,4	0.34	0	6,6,6	0.08	0
2	SO4	A	907	-	4,4,4	0.35	0	6,6,6	0.07	0
2	SO4	A	908	-	4,4,4	0.35	0	6,6,6	0.10	0
2	SO4	A	909	-	4,4,4	0.34	0	6,6,6	0.13	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SO4	A	901	-	-	0/0/0/0	0/0/0/0
2	SO4	A	902	-	-	0/0/0/0	0/0/0/0
2	SO4	A	903	-	-	0/0/0/0	0/0/0/0
2	SO4	A	904	-	-	0/0/0/0	0/0/0/0
2	SO4	A	905	-	-	0/0/0/0	0/0/0/0
2	SO4	A	906	-	-	0/0/0/0	0/0/0/0
2	SO4	A	907	-	-	0/0/0/0	0/0/0/0
2	SO4	A	908	-	-	0/0/0/0	0/0/0/0
2	SO4	A	909	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	904	SO4	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	585/757 (77%)	-0.09	29 (4%) 30 23	24, 43, 98, 100	0

The worst 5 of 29 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	626	ALA	8.4
1	A	623	TYR	5.7
1	A	612	ALA	5.4
1	A	624	ILE	5.3
1	A	610	THR	4.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	SO4	A	906	5/5	0.94	0.44	9.24	72,73,73,73	5

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	SO4	A	903	5/5	0.75	0.35	7.18	62,63,70,70	5
2	SO4	A	904	5/5	0.94	0.44	4.87	83,83,85,86	5
2	SO4	A	907	5/5	0.83	0.26	0.02	69,69,71,71	5
2	SO4	A	901	5/5	0.87	0.19	-	66,66,68,69	5
2	SO4	A	908	5/5	0.78	0.35	-	61,61,63,64	5
2	SO4	A	905	5/5	0.91	0.21	-	65,66,67,67	5
2	SO4	A	902	5/5	0.96	0.13	-	59,60,61,61	5
2	SO4	A	909	5/5	0.93	0.23	-	43,46,48,48	5

## 6.5 Other polymers [i](#)

There are no such residues in this entry.