



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 01:01 am GMT

PDB ID : 2E33
Title : Structural basis for selection of glycosylated substrate by SCFFbs1 ubiquitin ligase
Authors : Mizushima, T.; Yoshida, Y.; Kumanomidou, T.; Hasegawa, Y.; Yamane, T.; Tanaka, K.
Deposited on : 2006-11-20
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	2259 (2.70-2.70)
Clashscore	112137	2590 (2.70-2.70)
Ramachandran outliers	110173	2550 (2.70-2.70)
Sidechain outliers	110143	2550 (2.70-2.70)
RSRZ outliers	101464	2275 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	197	
2	B	124	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2423 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called F-box only protein 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	175	Total	C	N	O	S	0	0	0
			1412	893	237	279	3			

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	101	GLY	-	CLONING ARTIFACT	UNP Q80UW2
A	102	SER	-	CLONING ARTIFACT	UNP Q80UW2
A	103	HIS	-	CLONING ARTIFACT	UNP Q80UW2
A	104	MET	-	INITIATING METHIONINE	UNP Q80UW2
A	151	LYS	ARG	SEE REMARK 999	UNP Q80UW2

- Molecule 2 is a protein called Ribonuclease pancreatic.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	124	Total	C	N	O	S	0	0	0
			950	575	171	192	12			


- Molecule 3 is a polymer of unknown type called SUGAR (5-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	B	5	Total	C	N	O	0	0
			61	34	2	25		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: F-box only protein 2



- Molecule 2: Ribonuclease pancreatic



4 Data and refinement statistics

Property	Value	Source
Space group	P 4 3 2	Depositor
Cell constants a, b, c, α , β , γ	148.31Å 148.31Å 148.31Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.70 46.90 – 2.70	Depositor EDS
% Data completeness (in resolution range)	99.4 (50.00-2.70) 99.4 (46.90-2.70)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.24 (at 2.69Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.216 , 0.288 0.215 , 0.287	Depositor DCC
R_{free} test set	790 reflections (5.25%)	DCC
Wilson B-factor (Å ²)	56.3	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 32.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	2423	wwPDB-VP
Average B, all atoms (Å ²)	54.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.58% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, NAG, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.26	6/1452 (0.4%)	1.21	8/1973 (0.4%)
2	B	1.27	2/966 (0.2%)	1.20	2/1304 (0.2%)
All	All	1.27	8/2418 (0.3%)	1.21	10/3277 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	0	1

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	190	ALA	CA-CB	7.65	1.68	1.52
2	B	65	CYS	CB-SG	-7.62	1.69	1.82
1	A	211	TYR	CD2-CE2	6.02	1.48	1.39
1	A	237	GLU	CG-CD	5.70	1.60	1.51
2	B	37	LYS	CD-CE	5.67	1.65	1.51
1	A	162	GLU	CB-CG	5.66	1.62	1.52
1	A	144	GLU	CG-CD	5.44	1.60	1.51
1	A	194	TRP	CG-CD1	5.14	1.44	1.36

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	226	ARG	NE-CZ-NH2	-8.26	116.17	120.30
1	A	123	ARG	NE-CZ-NH1	-7.54	116.53	120.30
1	A	123	ARG	NE-CZ-NH2	6.33	123.46	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	14	ASP	CB-CG-OD1	-6.13	112.78	118.30
1	A	226	ARG	NE-CZ-NH1	5.85	123.23	120.30
2	B	26	CYS	CA-CB-SG	-5.44	104.20	114.00
1	A	184	GLN	N-CA-C	-5.30	96.70	111.00
1	A	148	ASP	CB-CG-OD1	-5.18	113.64	118.30
1	A	287	ARG	NE-CZ-NH1	-5.15	117.73	120.30
1	A	137	LEU	CA-CB-CG	-5.12	103.53	115.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	69	GLN	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1412	0	1297	42	0
2	B	950	0	904	37	0
3	B	61	0	52	7	0
All	All	2423	0	2253	79	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

All (79) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:179:TRP:HB3	1:A:181:ARG:NH1	1.58	1.17
1:A:179:TRP:CB	1:A:181:ARG:HH12	1.60	1.12
1:A:179:TRP:HB3	1:A:181:ARG:HH12	0.99	1.10
2:B:49:GLU:HG3	2:B:79:MET:HG2	1.52	0.92
2:B:39:ARG:HH11	2:B:39:ARG:HB2	1.32	0.91
2:B:39:ARG:HH11	2:B:39:ARG:CB	1.88	0.85

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:34:ASN:CG	3:B:1001:NAG:C1	2.46	0.83
2:B:39:ARG:HB2	2:B:39:ARG:NH1	1.93	0.82
1:A:279:TYR:CD2	3:B:1001:NAG:H82	2.23	0.72
2:B:49:GLU:HG3	2:B:79:MET:CG	2.22	0.70
1:A:179:TRP:CD1	1:A:275:GLN:NE2	2.56	0.70
1:A:207:VAL:HG21	1:A:225:VAL:HG11	1.75	0.68
1:A:210:TRP:CD1	1:A:290:ASN:HB3	2.30	0.67
2:B:113:ASN:HA	2:B:114:PRO:C	2.16	0.66
1:A:158:ASP:HB3	1:A:214:ARG:HH12	1.61	0.64
1:A:207:VAL:HG11	1:A:225:VAL:HG21	1.83	0.61
2:B:88:GLY:O	2:B:89:SER:CB	2.50	0.59
2:B:42:PRO:HA	2:B:86:GLU:HG3	1.83	0.59
2:B:78:THR:HG22	2:B:105:HIS:CE1	2.37	0.59
1:A:279:TYR:CE2	3:B:1001:NAG:H82	2.39	0.58
1:A:153:GLU:HB2	1:A:172:TYR:CZ	2.39	0.58
2:B:106:ILE:HG13	2:B:106:ILE:O	2.03	0.58
2:B:103:ASN:C	2:B:103:ASN:HD22	2.05	0.57
2:B:11:GLN:HG2	2:B:35:LEU:CD2	2.34	0.57
1:A:130:ASN:ND2	1:A:135:GLU:H	2.04	0.56
2:B:39:ARG:CB	2:B:39:ARG:NH1	2.62	0.55
1:A:214:ARG:HD3	1:A:284:PHE:HA	1.89	0.55
1:A:172:TYR:HA	1:A:289:THR:HB	1.89	0.54
2:B:25:TYR:CZ	2:B:29:MET:HG3	2.42	0.54
2:B:78:THR:CG2	2:B:105:HIS:CE1	2.89	0.54
1:A:245:VAL:O	1:A:246:PRO:C	2.45	0.54
1:A:189:GLN:HA	1:A:193:TYR:O	2.08	0.54
2:B:79:MET:O	2:B:103:ASN:HA	2.09	0.53
2:B:4:ALA:O	2:B:118:VAL:CG2	2.56	0.53
1:A:296:GLU:HB2	1:A:297:PRO:HD2	1.91	0.52
1:A:132:CYS:HB3	1:A:171:LYS:O	2.10	0.52
1:A:159:ASN:HB2	1:A:214:ARG:NH2	2.24	0.51
1:A:181:ARG:HH11	1:A:181:ARG:HG3	1.74	0.51
2:B:29:MET:HB3	2:B:46:PHE:CZ	2.46	0.51
1:A:157:GLY:HA3	1:A:172:TYR:OH	2.10	0.51
1:A:144:GLU:OE2	1:A:181:ARG:HD2	2.10	0.50
1:A:142:ASP:N	1:A:142:ASP:OD1	2.44	0.50
2:B:11:GLN:HG2	2:B:35:LEU:HD22	1.94	0.50
2:B:4:ALA:O	2:B:118:VAL:HG23	2.12	0.49
1:A:210:TRP:HD1	1:A:290:ASN:HB3	1.77	0.49
2:B:34:ASN:OD1	3:B:1001:NAG:C1	2.59	0.49
2:B:56:ALA:O	2:B:60:GLN:HG3	2.13	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:49:GLU:CG	2:B:79:MET:HG2	2.34	0.49
1:A:229:SER:HA	1:A:264:GLY:O	2.13	0.48
1:A:176:SER:OG	1:A:282:GLY:O	2.32	0.47
1:A:158:ASP:OD2	3:B:1004:MAN:H62	2.14	0.47
2:B:67:ASN:HD21	2:B:69:GLN:NE2	2.13	0.47
2:B:69:GLN:HB3	2:B:70:THR:H	1.48	0.47
2:B:66:LYS:HE3	2:B:121:ASP:O	2.15	0.46
2:B:11:GLN:HG2	2:B:35:LEU:HD21	1.96	0.46
1:A:211:TYR:HA	1:A:287:ARG:O	2.15	0.46
2:B:33:ARG:HH11	2:B:33:ARG:HG3	1.81	0.46
1:A:177:PHE:HB3	1:A:280:TRP:O	2.16	0.45
2:B:103:ASN:ND2	2:B:103:ASN:C	2.68	0.44
2:B:69:GLN:HE21	2:B:69:GLN:HB2	1.63	0.44
1:A:208:LYS:HD2	1:A:253:GLU:OE2	2.17	0.44
2:B:17:THR:O	2:B:48:HIS:HB3	2.18	0.44
1:A:270:PHE:C	1:A:270:PHE:CD2	2.91	0.44
2:B:34:ASN:ND2	3:B:1001:NAG:O5	2.43	0.43
1:A:288:VAL:O	1:A:289:THR:HB	2.19	0.43
1:A:206:VAL:O	1:A:293:VAL:HA	2.18	0.42
1:A:130:ASN:HD21	1:A:135:GLU:H	1.68	0.42
1:A:158:ASP:HB3	1:A:214:ARG:NH1	2.29	0.42
1:A:214:ARG:HH21	1:A:216:ASP:CG	2.23	0.42
2:B:8:PHE:C	2:B:8:PHE:CD2	2.94	0.41
1:A:183:ALA:HA	1:A:270:PHE:O	2.21	0.41
1:A:280:TRP:HA	3:B:1001:NAG:HN2	1.84	0.41
2:B:4:ALA:O	2:B:118:VAL:HG22	2.20	0.41
1:A:140:TRP:CE3	1:A:184:GLN:HG3	2.55	0.41
1:A:134:GLU:HA	1:A:134:GLU:OE1	2.21	0.41
2:B:10:ARG:O	2:B:33:ARG:HG3	2.21	0.41
1:A:231:ASN:O	1:A:232:GLU:HB2	2.20	0.40
1:A:210:TRP:CH2	1:A:253:GLU:HB2	2.56	0.40
2:B:71:ASN:ND2	2:B:110:CYS:O	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	173/197 (88%)	156 (90%)	13 (8%)	4 (2%)	7	19
2	B	122/124 (98%)	106 (87%)	16 (13%)	0	100	100
All	All	295/321 (92%)	262 (89%)	29 (10%)	4 (1%)	13	33

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	149	GLY
1	A	166	ASP
1	A	260	ASP
1	A	278	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	148/167 (89%)	131 (88%)	17 (12%)	6	15
2	B	109/109 (100%)	94 (86%)	15 (14%)	4	10
All	All	257/276 (93%)	225 (88%)	32 (12%)	5	12

All (32) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	137	LEU
1	A	142	ASP
1	A	155	LEU
1	A	161	VAL
1	A	165	GLN
1	A	170	LYS
1	A	176	SER
1	A	194	TRP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	195	GLU
1	A	196	GLU
1	A	200	THR
1	A	231	ASN
1	A	250	SER
1	A	255	SER
1	A	270	PHE
1	A	272	HIS
1	A	281	LYS
2	B	18	SER
2	B	39	ARG
2	B	55	GLN
2	B	59	SER
2	B	63	VAL
2	B	69	GLN
2	B	70	THR
2	B	85	ARG
2	B	87	THR
2	B	89	SER
2	B	98	LYS
2	B	103	ASN
2	B	115	TYR
2	B	116	VAL
2	B	118	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	130	ASN
1	A	159	ASN
1	A	242	GLN
2	B	24	ASN
2	B	55	GLN
2	B	69	GLN
2	B	103	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

5 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	NAG	B	1001	3,2	14,14,15	1.00	1 (7%)	15,19,21	2.39	7 (46%)
3	NAG	B	1002	3	14,14,15	1.01	1 (7%)	15,19,21	2.39	5 (33%)
3	BMA	B	1003	3	11,11,12	1.34	2 (18%)	13,15,17	2.35	5 (38%)
3	MAN	B	1004	3	11,11,12	0.90	1 (9%)	13,15,17	2.06	3 (23%)
3	MAN	B	1005	3	11,11,12	1.70	2 (18%)	13,15,17	2.34	7 (53%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	B	1001	3,2	-	0/6/23/26	0/1/1/1
3	NAG	B	1002	3	-	0/6/23/26	0/1/1/1
3	BMA	B	1003	3	-	0/2/19/22	0/1/1/1
3	MAN	B	1004	3	-	0/2/19/22	0/1/1/1
3	MAN	B	1005	3	-	0/2/19/22	0/1/1/1

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	1003	BMA	O5-C1	-2.94	1.38	1.43
3	B	1002	NAG	O5-C1	-2.31	1.39	1.43
3	B	1003	BMA	O2-C2	-2.21	1.38	1.43
3	B	1001	NAG	O3-C3	-2.07	1.38	1.43

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	1004	MAN	O4-C4	-2.04	1.38	1.43
3	B	1005	MAN	C4-C3	2.67	1.59	1.52
3	B	1005	MAN	C2-C3	4.17	1.58	1.52

All (27) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	1002	NAG	C2-N2-C7	-5.68	114.65	122.94
3	B	1002	NAG	O5-C1-C2	-4.56	105.13	111.47
3	B	1003	BMA	O4-C4-C3	-4.21	101.19	110.36
3	B	1001	NAG	O6-C6-C5	-3.98	97.96	111.34
3	B	1004	MAN	C6-C5-C4	-3.98	103.70	113.00
3	B	1002	NAG	C3-C4-C5	-3.26	104.48	110.22
3	B	1001	NAG	O7-C7-C8	-3.07	116.47	122.06
3	B	1003	BMA	O2-C2-C3	-2.99	104.30	110.17
3	B	1003	BMA	O5-C1-C2	-2.52	106.84	110.79
3	B	1005	MAN	O4-C4-C5	-2.39	103.27	109.28
3	B	1001	NAG	O3-C3-C2	-2.32	104.42	109.39
3	B	1002	NAG	O4-C4-C5	-2.12	103.95	109.28
3	B	1001	NAG	O3-C3-C4	-2.11	105.76	110.36
3	B	1004	MAN	C3-C4-C5	2.07	113.86	110.22
3	B	1005	MAN	C1-C2-C3	2.10	112.31	109.65
3	B	1005	MAN	O4-C4-C3	2.23	115.21	110.36
3	B	1002	NAG	O4-C4-C3	2.82	116.50	110.36
3	B	1003	BMA	C2-C3-C4	2.89	115.92	110.88
3	B	1005	MAN	O5-C1-C2	3.01	115.50	110.79
3	B	1001	NAG	C1-O5-C5	3.08	116.42	112.17
3	B	1001	NAG	C3-C4-C5	3.27	115.97	110.22
3	B	1005	MAN	O2-C2-C3	3.44	116.94	110.17
3	B	1005	MAN	C3-C4-C5	3.83	116.96	110.22
3	B	1005	MAN	C2-C3-C4	3.85	117.58	110.88
3	B	1001	NAG	O5-C1-C2	4.29	117.44	111.47
3	B	1003	BMA	C1-C2-C3	4.62	115.51	109.65
3	B	1004	MAN	C1-O5-C5	4.79	118.77	112.17

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	1001	NAG	6	0
3	B	1004	MAN	1	0

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	175/197 (88%)	-0.17	0 100 100	37, 49, 66, 75	0
2	B	124/124 (100%)	-0.17	0 100 100	41, 57, 78, 82	0
All	All	299/321 (93%)	-0.17	0 100 100	37, 52, 75, 82	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	NAG	B	1001	14/15	0.97	0.14	-0.89	54,57,63,65	0
3	MAN	B	1005	11/12	0.74	0.21	-	88,91,96,97	0
3	NAG	B	1002	14/15	0.94	0.21	-	61,65,77,78	0
3	MAN	B	1004	11/12	0.96	0.23	-	70,72,75,77	0
3	BMA	B	1003	11/12	0.95	0.19	-	59,65,70,73	0

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.