



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 2, 2017 – 09:00 AM EDT

PDB ID : 2E3A
Title : Crystal structure of the NO-bound form of Arthromyces ramosus peroxidase
at 1.3 Angstroms resolution
Authors : Fukuyama, K.; Okada, T.
Deposited on : unknown
Resolution : 1.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20030345
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20030345


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	112137	1185 (1.32-1.28)
Ramachandran outliers	110173	1138 (1.32-1.28)
Sidechain outliers	110143	1138 (1.32-1.28)
RSRZ outliers	101464	1133 (1.32-1.28)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	344	

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 4478 atoms, of which 1515 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Peroxidase.


Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	336	3987	1539	1515	422	496	15	0	3	0

- Molecule 2 is N-ACETYL-D-GLUCOSAMINE (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	15	8	1	6	0	1
2	A	1	14	8	1	5	0	0

- Molecule 3 is ALPHA-D-MANNOSE (three-letter code: MAN) (formula: $C_6H_{12}O_6$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			11	6	5		

- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	2	Total Ca 2 2	0	0

- Molecule 5 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	Fe	N	O	
			44	35	1	4	4	
							0	1

- Molecule 6 is NITRIC OXIDE (three-letter code: NO) (formula: NO).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	N	O		
			2	1	1	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	403	Total	O		
			403	403	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

● Molecule 1: Peroxidase



4 Data and refinement statistics

Property	Value	Source
Space group	P 4 ₂ 2 ₁ 2	Depositor
Cell constants a, b, c, α , β , γ	73.77Å 73.77Å 115.80Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	10.00 – 1.30 9.98 – 1.30	Depositor EDS
% Data completeness (in resolution range)	(Not available) (10.00-1.30) 91.9 (9.98-1.30)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	6.85 (at 1.30Å)	Xtriage
Refinement program	SHELX, SHELXL-97	Depositor
R, R_{free}	0.148 , 0.201 0.152 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	9.7	Xtriage
Anisotropy	0.665	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.43 , 60.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	4478	wwPDB-VP
Average B, all atoms (Å ²)	15.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.67% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, CA, MAN, NAG, NO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.58	0/2543	1.14	11/3467 (0.3%)

There are no bond length outliers.

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	15	GLY	C-N-CA	10.22	147.25	121.70
1	A	242	ARG	NE-CZ-NH2	9.12	124.86	120.30
1	A	52	ARG	NE-CZ-NH2	-6.60	117.00	120.30
1	A	254	ARG	NE-CZ-NH1	-6.17	117.21	120.30
1	A	258	ARG	NE-CZ-NH2	5.96	123.28	120.30
1	A	134	ARG	CD-NE-CZ	5.94	131.91	123.60
1	A	337	LEU	CA-CB-CG	5.91	128.90	115.30
1	A	258	ARG	NE-CZ-NH1	-5.71	117.44	120.30
1	A	268	VAL	CG1-CB-CG2	-5.47	102.15	110.90
1	A	244	ARG	CD-NE-CZ	5.25	130.95	123.60
1	A	90	PHE	CB-CG-CD2	-5.11	117.22	120.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2472	1515	2374	7	0
2	A	29	0	19	0	0
3	A	11	0	10	0	0
4	A	2	0	0	0	0
5	A	44	0	6	0	0
6	A	2	0	0	0	0
7	A	403	0	0	2	0
All	All	2963	1515	2409	7	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (7) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:34:GLN:HE21	1:A:34:GLN:HA	1.49	0.78
1:A:237:PHE:HB3	1:A:337:LEU:HG	1.68	0.76
1:A:193:ASN:HD22	1:A:195:ALA:H	1.56	0.53
1:A:34:GLN:HA	1:A:34:GLN:NE2	2.23	0.51
1:A:194:SER:HA	7:A:1148:HOH:O	2.19	0.42
1:A:244:ARG:HG2	7:A:1048:HOH:O	2.20	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	337/344 (98%)	328 (97%)	9 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	274/273 (100%)	268 (98%)	6 (2%)	57 15

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	17	SER
1	A	34	GLN
1	A	50	ILE
1	A	193	ASN
1	A	334	SER
1	A	337	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	34	GLN
1	A	109	ASN
1	A	128	ASN
1	A	193	ASN
1	A	288	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 9 ligands modelled in this entry, 2 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	HEM	A	401[A]	-	28,50,50	1.54	6 (21%)	17,82,82	1.56	4 (23%)
5	HEM	A	401[B]	-	28,50,50	1.53	5 (17%)	17,82,82	1.56	4 (23%)
2	NAG	A	402[A]	-	14,14,15	1.54	3 (21%)	15,19,21	1.66	3 (20%)
2	NAG	A	402[B]	-	14,14,15	1.55	3 (21%)	15,19,21	1.66	3 (20%)
2	NAG	A	403	2	14,14,15	1.51	3 (21%)	15,19,21	1.49	5 (33%)
3	MAN	A	404	1	11,11,12	1.38	1 (9%)	13,15,17	1.73	4 (30%)
6	NO	A	503	5	0,1,1	0.00	-	0,0,0	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	HEM	A	401[A]	-	-	0/6/54/54	0/0/8/8
5	HEM	A	401[B]	-	-	0/6/54/54	0/0/8/8
2	NAG	A	402[A]	-	-	0/6/23/26	0/1/1/1
2	NAG	A	402[B]	-	-	0/6/23/26	0/1/1/1
2	NAG	A	403	2	-	0/6/23/26	0/1/1/1
3	MAN	A	404	1	-	0/2/19/22	0/1/1/1
6	NO	A	503	5	-	0/0/0/0	0/0/0/0

All (21) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	401[B]	HEM	C3C-CAC	-3.49	1.40	1.47
5	A	401[A]	HEM	C3C-CAC	-3.49	1.40	1.47
2	A	403	NAG	O7-C7	-2.89	1.16	1.23
2	A	402[A]	NAG	O7-C7	-2.75	1.16	1.23
2	A	402[B]	NAG	O7-C7	-2.75	1.16	1.23

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	401[B]	HEM	C3B-CAB	-2.60	1.42	1.47
5	A	401[A]	HEM	C3B-CAB	-2.60	1.42	1.47
2	A	402[A]	NAG	C8-C7	-2.07	1.46	1.50
2	A	402[B]	NAG	C8-C7	-2.07	1.46	1.50
2	A	403	NAG	O5-C1	-2.02	1.40	1.43
5	A	401[A]	HEM	CBB-CAB	2.04	1.43	1.28
5	A	401[B]	HEM	CBC-CAC	2.20	1.44	1.28
5	A	401[A]	HEM	CBC-CAC	2.20	1.44	1.28
2	A	403	NAG	C1-C2	2.26	1.55	1.52
5	A	401[B]	HEM	C1A-NA	2.42	1.41	1.36
5	A	401[A]	HEM	C1A-NA	2.42	1.41	1.36
3	A	404	MAN	C2-C3	3.07	1.56	1.52
5	A	401[B]	HEM	C1C-NC	3.36	1.40	1.36
5	A	401[A]	HEM	C1C-NC	3.36	1.40	1.36
2	A	402[A]	NAG	C1-C2	3.78	1.57	1.52
2	A	402[B]	NAG	C1-C2	3.78	1.57	1.52

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	402[A]	NAG	O5-C1-C2	-4.14	105.71	111.47
2	A	402[B]	NAG	O5-C1-C2	-4.14	105.71	111.47
5	A	401[B]	HEM	C4C-C3C-C2C	-3.23	104.64	106.90
5	A	401[A]	HEM	C4C-C3C-C2C	-3.23	104.64	106.90
2	A	403	NAG	C1-O5-C5	-2.98	108.06	112.17
2	A	402[A]	NAG	O4-C4-C5	-2.64	102.63	109.28
2	A	402[B]	NAG	O4-C4-C5	-2.64	102.63	109.28
3	A	404	MAN	O2-C2-C3	-2.36	105.54	110.17
5	A	401[B]	HEM	CMA-C3A-C4A	-2.32	124.89	128.46
5	A	401[A]	HEM	CMA-C3A-C4A	-2.32	124.89	128.46
2	A	402[A]	NAG	C2-N2-C7	-2.29	119.61	122.94
2	A	402[B]	NAG	C2-N2-C7	-2.29	119.61	122.94
5	A	401[B]	HEM	CMD-C2D-C1D	-2.11	125.22	128.46
5	A	401[A]	HEM	CMD-C2D-C1D	-2.11	125.22	128.46
2	A	403	NAG	O4-C4-C3	-2.11	105.76	110.36
2	A	403	NAG	O5-C1-C2	-2.08	108.58	111.47
2	A	403	NAG	C2-N2-C7	-2.07	119.92	122.94
2	A	403	NAG	C6-C5-C4	2.21	118.17	113.00
3	A	404	MAN	O5-C1-C2	2.23	114.29	110.79
5	A	401[B]	HEM	C3B-C4B-NB	2.47	112.40	109.21
5	A	401[A]	HEM	C3B-C4B-NB	2.47	112.40	109.21
3	A	404	MAN	O3-C3-C2	2.75	115.03	110.02

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	404	MAN	C1-C2-C3	3.02	113.48	109.65

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	336/344 (97%)	-0.44	8 (2%) 59 58	8, 12, 27, 51	1 (0%)

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	15	GLY	3.5
1	A	333	ALA	3.2
1	A	13	PRO	2.9
1	A	9	SER	2.8
1	A	334	SER	2.7
1	A	17	SER	2.6
1	A	11	THR	2.4
1	A	16	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	NAG	A	402[A]	14/15	0.97	0.06	1.00	12,13,22,23	1
2	NAG	A	402[B]	14/15	0.97	0.06	1.00	12,13,22,23	1
5	HEM	A	401[A]	43/43	0.99	0.05	0.48	7,8,10,18	1
5	HEM	A	401[B]	43/43	0.99	0.05	0.48	7,8,10,13	1
3	MAN	A	404	11/12	0.96	0.06	0.05	13,15,17,17	0
4	CA	A	502	1/1	1.00	0.04	-1.18	8,8,8,8	0
4	CA	A	501	1/1	1.00	0.03	-1.58	9,9,9,9	0
2	NAG	A	403	14/15	0.95	0.07	-	12,18,26,31	0
6	NO	A	503	2/2	0.98	0.09	-	12,12,12,15	0

6.5 Other polymers

There are no such residues in this entry.