



wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 5, 2017 – 08:14 AM EST

PDB ID : 4E6H
Title : CRYSTAL STRUCTURE OF THE HAT domain of k. lactis RNA14
Authors : Paulson, A.R.; Tong, L.
Deposited on : unknown
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20030345
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20030345


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	4130 (2.30-2.30)
Clashscore	112137	4751 (2.30-2.30)
Ramachandran outliers	110173	4705 (2.30-2.30)
Sidechain outliers	110143	4704 (2.30-2.30)
RSRZ outliers	101464	4156 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	679	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 4838 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called mRNA 3'-end-processing protein RNA14.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	553	4608	2967	764	852	25	0	0	0

There are 35 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-17	MET	-	EXPRESSION TAG	UNP Q6CII8
A	-16	GLY	-	EXPRESSION TAG	UNP Q6CII8
A	-15	SER	-	EXPRESSION TAG	UNP Q6CII8
A	-14	SER	-	EXPRESSION TAG	UNP Q6CII8
A	-13	HIS	-	EXPRESSION TAG	UNP Q6CII8
A	-12	HIS	-	EXPRESSION TAG	UNP Q6CII8
A	-11	HIS	-	EXPRESSION TAG	UNP Q6CII8
A	-10	HIS	-	EXPRESSION TAG	UNP Q6CII8
A	-9	HIS	-	EXPRESSION TAG	UNP Q6CII8
A	-8	HIS	-	EXPRESSION TAG	UNP Q6CII8
A	-7	SER	-	EXPRESSION TAG	UNP Q6CII8
A	-6	SER	-	EXPRESSION TAG	UNP Q6CII8
A	-5	GLY	-	EXPRESSION TAG	UNP Q6CII8
A	-4	LEU	-	EXPRESSION TAG	UNP Q6CII8
A	-3	VAL	-	EXPRESSION TAG	UNP Q6CII8
A	-2	PRO	-	EXPRESSION TAG	UNP Q6CII8
A	-1	ARG	-	EXPRESSION TAG	UNP Q6CII8
A	0	GLY	-	EXPRESSION TAG	UNP Q6CII8
A	1	SER	-	EXPRESSION TAG	UNP Q6CII8
A	2	HIS	-	EXPRESSION TAG	UNP Q6CII8
A	3	MET	-	EXPRESSION TAG	UNP Q6CII8
A	4	ALA	-	EXPRESSION TAG	UNP Q6CII8
A	5	SER	-	EXPRESSION TAG	UNP Q6CII8
A	6	MET	-	EXPRESSION TAG	UNP Q6CII8
A	7	THR	-	EXPRESSION TAG	UNP Q6CII8
A	8	GLY	-	EXPRESSION TAG	UNP Q6CII8
A	9	GLY	-	EXPRESSION TAG	UNP Q6CII8

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	10	GLN	-	EXPRESSION TAG	UNP Q6CII8
A	11	GLN	-	EXPRESSION TAG	UNP Q6CII8
A	12	MET	-	EXPRESSION TAG	UNP Q6CII8
A	13	GLY	-	EXPRESSION TAG	UNP Q6CII8
A	14	ARG	-	EXPRESSION TAG	UNP Q6CII8
A	15	GLY	-	EXPRESSION TAG	UNP Q6CII8
A	16	SER	-	EXPRESSION TAG	UNP Q6CII8
A	17	MET	-	EXPRESSION TAG	UNP Q6CII8

- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	230	Total O 230 230	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	117.35Å 59.79Å 123.48Å 90.00° 101.55° 90.00°	Depositor
Resolution (Å)	29.32 – 2.30 29.32 – 2.28	Depositor EDS
% Data completeness (in resolution range)	89.2 (29.32-2.30) 87.8 (29.32-2.28)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.18 (at 2.29Å)	Xtriage
Refinement program	CNS 1.2	Depositor
R, R_{free}	0.210 , 0.270 0.204 , 0.265	Depositor DCC
R_{free} test set	1676 reflections (5.00%)	DCC
Wilson B-factor (Å ²)	41.4	Xtriage
Anisotropy	0.501	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 44.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4838	wwPDB-VP
Average B, all atoms (Å ²)	56.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.07% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.69	0/4708	0.79	5/6363 (0.1%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	77	ARG	NE-CZ-NH2	-8.51	116.05	120.30
1	A	202	LEU	CA-CB-CG	6.09	129.30	115.30
1	A	77	ARG	CG-CD-NE	-5.50	100.24	111.80
1	A	77	ARG	NE-CZ-NH1	5.10	122.85	120.30
1	A	443	ARG	NE-CZ-NH1	-5.09	117.76	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4608	0	4608	157	0
2	A	230	0	0	17	0
All	All	4838	0	4608	157	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

The worst 5 of 157 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:258:LYS:HG3	1:A:260:ASN:HD21	1.28	0.98
1:A:202:LEU:HD21	1:A:213:MET:HG2	1.47	0.97
1:A:563:THR:CG2	1:A:572:ASN:HD21	1.80	0.93
1:A:580:ASP:HB2	2:A:736:HOH:O	1.70	0.91
1:A:450:THR:HG23	1:A:451:PRO:HD2	1.58	0.83

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	549/679 (81%)	503 (92%)	33 (6%)	13 (2%)	7 5

5 of 13 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	237	SER
1	A	255	LYS
1	A	518	ASP
1	A	519	SER
1	A	520	HIS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
-----	-------	----------	-----------	----------	-------------

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	514/624 (82%)	482 (94%)	32 (6%)	21 28

5 of 32 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	275	LYS
1	A	372	LEU
1	A	539	LEU
1	A	358	ASN
1	A	385	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 16 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	277	ASN
1	A	282	GLN
1	A	354	GLN
1	A	260	ASN
1	A	358	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	553/679 (81%)	0.12	18 (3%) 47 54	29, 53, 90, 121	0

The worst 5 of 18 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	339	ASN	4.1
1	A	189	GLN	3.9
1	A	185	LYS	3.8
1	A	28	ARG	3.8
1	A	183	VAL	3.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.