



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 14, 2017 – 04:39 am GMT

PDB ID : 4E7H
Title : PFV intasome prior to 3'-processing, Apo form (UI-Apo)
Authors : Hare, S.; Cherepanov, P.
Deposited on : 2012-03-17
Resolution : 2.57 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.57 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	2899 (2.60-2.56)
Clashscore	112137	3268 (2.60-2.56)
Ramachandran outliers	110173	3218 (2.60-2.56)
Sidechain outliers	110143	3218 (2.60-2.56)
RSRZ outliers	101464	2907 (2.60-2.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	395	<div> <div>8%</div> <div> <div></div> <div>79%</div> <div>14%</div> <div>7%</div> </div> </div>
1	B	395	<div> <div>5%</div> <div> <div></div> <div>42%</div> <div>53%</div> </div> </div>
2	C	19	<div> <div>5%</div> <div> <div></div> <div>42%</div> <div>47%</div> <div>11%</div> </div> </div>
3	D	19	<div> <div></div> <div> <div></div> <div>37%</div> <div>47%</div> <div>16%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	SO4	A	405	-	-	-	X
4	SO4	A	408	-	-	-	X
5	GOL	B	402	-	-	-	X
5	GOL	B	403	-	-	-	X

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 5484 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pro-Pol polypeptide.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	368	Total	C	N	O	S	0	0	0
			2920	1872	512	532	4			
1	B	184	Total	C	N	O	S	0	0	0
			1451	942	235	273	1			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	EXPRESSION TAG	UNP P14350
A	-1	PRO	-	EXPRESSION TAG	UNP P14350
A	0	GLY	-	EXPRESSION TAG	UNP P14350
A	217	SER	GLY	VARIANT	UNP P14350
A	218	GLY	SER	VARIANT	UNP P14350
B	-2	GLY	-	EXPRESSION TAG	UNP P14350
B	-1	PRO	-	EXPRESSION TAG	UNP P14350
B	0	GLY	-	EXPRESSION TAG	UNP P14350
B	217	SER	GLY	VARIANT	UNP P14350
B	218	GLY	SER	VARIANT	UNP P14350

- Molecule 2 is a DNA chain called DNA (5'-D(*AP*TP*TP*GP*TP*CP*AP*TP*GP*GP*AP*AP*TP*TP*TP*CP*GP*CP*A)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	19	Total	C	N	O	P	0	0	0
			387	187	68	114	18			

- Molecule 3 is a DNA chain called DNA (5'-D(*TP*GP*CP*GP*AP*AP*AP*TP*TP*CP*CP*AP*TP*GP*AP*CP*AP*AP*T)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	19	Total	C	N	O	P	0	1	0
			406	196	74	117	19			

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		
5	D	1	Total	C	O	0	0
			6	3	3		
5	C	1	Total	C	O	0	0
			6	3	3		
5	B	1	Total	C	O	0	0
			6	3	3		
5	B	1	Total	C	O	0	0
			6	3	3		
5	D	1	Total	C	O	0	0
			6	3	3		

- Molecule 6 is HEXANE-1,6-DIOL (three-letter code: HEZ) (formula: C₆H₁₄O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			8	6	2		

- Molecule 7 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	1	Total	Zn	0	0
			1	1		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	134	Total	O	0	0
			134	134		
8	B	37	Total	O	0	0
			37	37		
8	C	35	Total	O	0	0
			35	35		
8	D	26	Total	O	0	0
			26	26		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Pro-Pol polyprotein



• Molecule 1: Pro-Pol polyprotein




• Molecule 2: DNA (5'-D(*AP*TP*TP*GP*TP*CP*AP*TP*GP*GP*AP*AP*TP*TP*TP*CP*GP*CP*A)-3')





● Molecule 3: DNA (5'-D(*TP*GP*CP*GP*AP*AP*AP*TP*TP*CP*CP*AP*TP*GP*AP*CP*AP*AP*T)-3')



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	159.47Å 159.47Å 123.75Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	36.92 – 2.57 36.92 – 2.57	Depositor EDS
% Data completeness (in resolution range)	99.2 (36.92-2.57) 99.2 (36.92-2.57)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.01 (at 2.58Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.7.3_928)	Depositor
R, R_{free}	0.186 , 0.219 0.169 , 0.204	Depositor DCC
R_{free} test set	2562 reflections (5.04%)	DCC
Wilson B-factor (Å ²)	63.8	Xtriage
Anisotropy	0.075	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 49.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5484	wwPDB-VP
Average B, all atoms (Å ²)	64.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.90% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, ZN, SO4, HEZ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.46	0/2999	0.62	0/4094
1	B	0.43	0/1491	0.56	0/2039
2	C	0.85	0/433	1.69	9/667 (1.3%)
3	D	0.89	0/454	1.83	17/696 (2.4%)
All	All	0.54	0/5377	0.92	26/7496 (0.3%)

There are no bond length outliers.

All (26) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	12	DA	O4'-C1'-N9	-12.70	99.11	108.00
2	C	5	DT	O4'-C1'-N1	-10.26	100.82	108.00
3	D	17	DA	O4'-C1'-N9	8.76	114.13	108.00
3	D	7	DA	O4'-C1'-N9	-8.56	102.01	108.00
3	D	18	DA	O4'-C1'-N9	-8.18	102.28	108.00
3	D	18	DA	O4'-C4'-C3'	-8.12	101.12	106.00
2	C	14	DT	O4'-C1'-N1	-8.04	102.37	108.00
3	D	1	DT	O4'-C4'-C3'	-7.88	101.27	106.00
2	C	12	DA	O4'-C1'-N9	-7.30	102.89	108.00
2	C	3	DT	O4'-C1'-N1	7.26	113.08	108.00
3	D	9	DT	O4'-C1'-N1	-7.26	102.92	108.00
3	D	10	DC	O4'-C1'-N1	-6.87	103.19	108.00
3	D	8	DT	O4'-C1'-N1	-6.71	103.30	108.00
2	C	11	DA	O4'-C1'-N9	-6.67	103.33	108.00
3	D	19[A]	DT	N3-C4-O4	6.33	123.69	119.90
3	D	19[B]	DT	N3-C4-O4	6.33	123.69	119.90
3	D	6	DA	O4'-C1'-N9	-5.86	103.90	108.00
3	D	18	DA	C3'-C2'-C1'	-5.73	95.62	102.50
2	C	18	DC	C1'-O4'-C4'	-5.72	104.38	110.10
3	D	14	DG	O4'-C1'-N9	-5.68	104.02	108.00

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	8	DT	O4'-C1'-N1	-5.07	104.45	108.00
3	D	4	DG	O4'-C1'-N9	-5.03	104.48	108.00
2	C	17	DG	O4'-C1'-N9	-5.02	104.49	108.00
2	C	19	DA	O4'-C1'-N9	5.02	111.51	108.00
3	D	19[A]	DT	C5-C4-O4	-5.02	121.39	124.90
3	D	19[B]	DT	C5-C4-O4	-5.02	121.39	124.90

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2920	0	2945	31	0
1	B	1451	0	1440	10	0
2	C	387	0	218	4	0
3	D	406	0	229	3	0
4	A	20	0	0	1	0
4	B	5	0	0	0	0
5	A	24	0	32	2	0
5	B	12	0	16	0	0
5	C	6	0	8	0	0
5	D	12	0	16	1	0
6	A	8	0	14	1	0
7	A	1	0	0	0	0
8	A	134	0	0	1	0
8	B	37	0	0	0	0
8	C	35	0	0	1	0
8	D	26	0	0	0	0
All	All	5484	0	4918	47	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:406:GOL:HO1	5:A:406:GOL:HO3	1.08	0.89
1:A:162:SER:HB2	5:A:402:GOL:H2	1.70	0.74
1:B:235:LEU:HG	1:B:239:PRO:HA	1.78	0.64
2:C:19:DA:N7	8:C:231:HOH:O	2.31	0.61
1:B:236:VAL:HB	1:B:294:LEU:HD13	1.84	0.60
1:A:358:LEU:N	1:A:359:GLY:HA2	2.17	0.59
1:A:315:TRP:CE2	1:A:371:PRO:HD3	2.39	0.58
1:B:141:TYR:CE2	1:B:161:PRO:HD3	2.41	0.56
1:A:202:ARG:NE	4:A:404:SO4:O2	2.31	0.55
1:A:69:ARG:HD2	1:A:70:GLU:OE2	2.08	0.54
1:B:127:ILE:HA	1:B:144:VAL:O	2.09	0.53
1:A:73:LEU:HD13	1:A:86:ARG:HG3	1.91	0.51
1:A:135:PRO:HG3	1:A:140:LEU:HD21	1.92	0.51
2:C:1:DA:H3'	2:C:2:DT:H71	1.93	0.49
1:A:138:GLY:O	1:A:159:LYS:HG2	2.13	0.49
1:A:341:SER:HB2	1:A:355:LEU:O	2.12	0.49
1:A:67:THR:HB	1:A:71:ALA:HB3	1.93	0.49
3:D:19[A]:DT:H6	3:D:19[A]:DT:H5''	1.77	0.48
1:A:25:PRO:HB2	1:A:28:TYR:HD2	1.78	0.47
1:A:70:GLU:OE1	6:A:409:HEZ:H12	2.14	0.47
1:A:161:PRO:O	1:A:189:ALA:HB2	2.14	0.47
1:A:75:LYS:NZ	8:A:621:HOH:O	2.46	0.47
2:C:1:DA:H2''	2:C:2:DT:H5'	1.95	0.47
3:D:18:DA:H2'	3:D:18:DA:O5'	2.15	0.46
1:A:14:GLN:O	1:A:17:GLN:HG2	2.16	0.46
2:C:18:DC:H2''	2:C:19:DA:O5'	2.17	0.45
1:B:231:LEU:HD23	1:B:231:LEU:HA	1.83	0.45
1:B:284:LEU:HA	1:B:284:LEU:HD12	1.76	0.45
1:A:246:LEU:HB2	1:A:247:PRO:HD3	1.99	0.44
1:A:73:LEU:HD22	1:A:86:ARG:CZ	2.48	0.43
1:A:360:ASN:N	1:A:360:ASN:OD1	2.51	0.43
1:A:284:LEU:HD12	1:A:284:LEU:HA	1.81	0.43
1:A:167:VAL:HG13	1:A:198:TRP:CD1	2.54	0.42
1:A:324:GLN:O	1:A:369:LEU:HA	2.18	0.42
1:A:141:TYR:CE2	1:A:161:PRO:HD3	2.55	0.42
1:A:213:HIS:CE1	1:A:215:GLN:HB2	2.54	0.42
1:A:337:TRP:CD2	1:A:370:LYS:HE2	2.54	0.42
3:D:17:DA:H8	5:D:101:GOL:H32	1.84	0.42
1:A:50:PRO:HG2	1:A:53:ASP:HB2	2.02	0.41
1:A:247:PRO:HB3	1:B:175:SER:OG	2.20	0.41
1:A:17:GLN:HG3	1:A:19:HIS:CD2	2.56	0.41
1:A:295:LEU:HD21	1:A:299:ARG:NH2	2.35	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:325:GLU:HA	1:A:369:LEU:HD23	2.02	0.41
1:A:275:ASN:HB3	1:B:178:ILE:HG22	2.03	0.40
1:A:127:ILE:HG22	1:A:145:VAL:HG13	2.03	0.40
1:B:141:TYR:CZ	1:B:161:PRO:HD3	2.56	0.40
1:B:236:VAL:HG21	1:B:294:LEU:HB2	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	366/395 (93%)	355 (97%)	10 (3%)	1 (0%)	44	68
1	B	182/395 (46%)	172 (94%)	10 (6%)	0	100	100
All	All	548/790 (69%)	527 (96%)	20 (4%)	1 (0%)	51	74

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	201	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	327/354 (92%)	319 (98%)	8 (2%)	54 78
1	B	162/354 (46%)	156 (96%)	6 (4%)	39 65
All	All	489/708 (69%)	475 (97%)	14 (3%)	48 73

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	9	ASP
1	A	13	ASP
1	A	36	LYS
1	A	137	GLN
1	A	149	MET
1	A	357	HIS
1	A	360	ASN
1	A	364	VAL
1	B	149	MET
1	B	164	SER
1	B	215	GLN
1	B	274	SER
1	B	284	LEU
1	B	298	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	19	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 16 ligands modelled in this entry, 1 is monoatomic - leaving 15 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	SO4	A	401	-	4,4,4	0.20	0	6,6,6	0.19	0
5	GOL	A	402	-	5,5,5	0.35	0	5,5,5	0.53	0
5	GOL	A	403	-	5,5,5	0.40	0	5,5,5	0.55	0
4	SO4	A	404	-	4,4,4	0.25	0	6,6,6	0.30	0
4	SO4	A	405	-	4,4,4	0.18	0	6,6,6	0.12	0
5	GOL	A	406	-	5,5,5	0.26	0	5,5,5	0.59	0
5	GOL	A	407	-	5,5,5	0.33	0	5,5,5	0.57	0
4	SO4	A	408	-	4,4,4	0.28	0	6,6,6	0.31	0
6	HEZ	A	409	-	7,7,7	0.49	0	6,6,6	0.60	0
4	SO4	B	401	-	4,4,4	0.22	0	6,6,6	0.23	0
5	GOL	B	402	-	5,5,5	0.37	0	5,5,5	0.25	0
5	GOL	B	403	-	5,5,5	0.35	0	5,5,5	0.85	0
5	GOL	C	101	-	5,5,5	0.36	0	5,5,5	0.62	0
5	GOL	D	101	-	5,5,5	0.33	0	5,5,5	0.75	0
5	GOL	D	102	-	5,5,5	0.26	0	5,5,5	0.44	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	SO4	A	401	-	-	0/0/0/0	0/0/0/0
5	GOL	A	402	-	-	0/4/4/4	0/0/0/0
5	GOL	A	403	-	-	0/4/4/4	0/0/0/0
4	SO4	A	404	-	-	0/0/0/0	0/0/0/0
4	SO4	A	405	-	-	0/0/0/0	0/0/0/0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	A	406	-	-	0/4/4/4	0/0/0/0
5	GOL	A	407	-	-	0/4/4/4	0/0/0/0
4	SO4	A	408	-	-	0/0/0/0	0/0/0/0
6	HEZ	A	409	-	-	0/5/5/5	0/0/0/0
4	SO4	B	401	-	-	0/0/0/0	0/0/0/0
5	GOL	B	402	-	-	0/4/4/4	0/0/0/0
5	GOL	B	403	-	-	0/4/4/4	0/0/0/0
5	GOL	C	101	-	-	0/4/4/4	0/0/0/0
5	GOL	D	101	-	-	0/4/4/4	0/0/0/0
5	GOL	D	102	-	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	402	GOL	1	0
4	A	404	SO4	1	0
5	A	406	GOL	1	0
6	A	409	HEZ	1	0
5	D	101	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	368/395 (93%)	0.44	32 (8%) 11 8	41, 55, 97, 133	0
1	B	184/395 (46%)	0.43	19 (10%) 7 5	43, 62, 138, 161	0
2	C	19/19 (100%)	-0.00	1 (5%) 27 22	46, 57, 87, 106	0
3	D	19/19 (100%)	-0.12	0 100 100	49, 53, 77, 100	0
All	All	590/828 (71%)	0.40	52 (8%) 11 8	41, 57, 110, 161	0

All (52) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	8	LEU	6.2
1	A	12	LEU	5.3
1	B	280	ASN	5.2
1	B	296	GLN	5.1
1	B	294	LEU	5.0
1	B	298	ILE	4.9
1	A	15	LEU	4.4
1	A	358	LEU	4.1
1	B	279	ALA	4.0
1	A	21	ILE	3.8
1	A	17	GLN	3.8
1	B	297	GLU	3.7
1	A	146	VAL	3.7
1	B	295	LEU	3.4
1	B	292	LEU	3.3
1	A	19	HIS	3.3
1	A	145	VAL	3.3
1	A	16	LEU	3.3
1	B	281	GLN	3.2
1	A	9	ASP	3.2
1	B	257	TYR	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	32	LEU	3.1
1	A	357	HIS	3.1
1	A	20	TYR	3.0
1	A	154	TRP	3.0
1	B	288	ARG	2.9
1	A	152	PHE	2.9
1	B	116	ASP	2.8
1	A	254	ASN	2.8
1	A	253	LEU	2.8
1	A	153	THR	2.8
1	A	239	PRO	2.7
1	B	299	ARG	2.7
1	A	265	PRO	2.6
1	A	250	GLN	2.5
1	B	293	SER	2.5
1	A	144	VAL	2.5
1	B	263	TYR	2.4
1	B	260	VAL	2.4
1	A	138	GLY	2.2
1	A	318	VAL	2.2
1	A	257	TYR	2.2
1	A	14	GLN	2.1
1	B	176	ILE	2.1
1	B	286	LEU	2.1
1	A	29	THR	2.1
1	A	176	ILE	2.1
1	A	155	LEU	2.1
1	B	237	GLY	2.0
2	C	1	DA	2.0
1	A	71	ALA	2.0
1	A	264	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	SO4	A	405	5/5	0.82	0.33	4.24	73,82,101,106	5
5	GOL	B	402	6/6	0.84	0.39	2.89	66,76,77,93	0
5	GOL	B	403	6/6	0.79	0.29	2.65	67,81,92,94	0
4	SO4	A	408	5/5	0.94	0.24	2.03	49,52,65,77	5
5	GOL	D	102	6/6	0.90	0.24	1.99	63,75,83,98	0
4	SO4	A	404	5/5	0.91	0.22	1.95	60,63,71,92	5
5	GOL	A	403	6/6	0.88	0.19	1.70	74,86,91,94	0
6	HEZ	A	409	8/8	0.93	0.27	1.34	63,70,84,91	0
5	GOL	C	101	6/6	0.95	0.23	0.95	55,57,62,80	0
7	ZN	A	410	1/1	1.00	0.15	0.29	51,51,51,51	0
4	SO4	B	401	5/5	0.93	0.16	-0.60	71,77,93,106	0
5	GOL	D	101	6/6	0.87	0.25	-	63,81,89,99	0
5	GOL	A	402	6/6	0.84	0.21	-	81,82,92,95	0
5	GOL	A	407	6/6	0.94	0.29	-	74,77,81,84	0
4	SO4	A	401	5/5	0.99	0.12	-	70,72,94,116	0
5	GOL	A	406	6/6	0.83	0.31	-	61,79,92,105	0

6.5 Other polymers [i](#)

There are no such residues in this entry.