



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 09:21 am GMT

PDB ID : 5E9B
Title : Crystal structure of human heparanase in complex with HepMer M09S05a
Authors : Wu, L.; Davies, G.J.
Deposited on : 2015-10-14
Resolution : 1.88 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.88 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	7505 (1.90-1.86)
Clashscore	112137	8369 (1.90-1.86)
Ramachandran outliers	110173	8279 (1.90-1.86)
Sidechain outliers	110143	8280 (1.90-1.86)
RSRZ outliers	101464	7571 (1.90-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	389	<div> <div></div> <div>93%</div> <div>6%</div> </div>
2	B	77	<div> <div></div> <div>91%</div> <div>5%</div> </div>

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 3962 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Heparanase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	385	Total	C	N	O	S	0	0	0
			3057	1974	522	550	11			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	155	ASP	-	expression tag	UNP Q9Y251
A	156	PRO	-	expression tag	UNP Q9Y251
A	157	GLY	-	expression tag	UNP Q9Y251
A	307	ARG	LYS	conflict	UNP Q9Y251

- Molecule 2 is a protein called Heparanase.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	74	Total	C	N	O	0	1	0
			595	388	97	110			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	33	ASP	-	expression tag	UNP Q9Y251
B	34	PRO	-	expression tag	UNP Q9Y251
B	35	GLY	-	expression tag	UNP Q9Y251

- Molecule 3 is N-ACETYL-D-GLUCOSAMINE (three-letter code: NAG) (formula: C₈H₁₅NO₆).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Cl	0	0
			1	1		

- Molecule 5 is BETA-D-GLUCOPYRANURONIC ACID (three-letter code: BDP) (formula: C₆H₁₀O₇).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			13	6	7		
5	A	1	Total	C	O	0	0
			12	6	6		

- Molecule 6 is N,O6-DISULFO-GLUCOSAMINE (three-letter code: SGN) (formula: $C_6H_{13}NO_{11}S_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	A	1	Total	C	N	O	S	0	0
			19	6	1	10	2		

- Molecule 7 is N-SULFO-ALPHA-D-GLUCOSAMINE (three-letter code: GNS) (formula: $C_6H_{13}NO_8S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
7	A	1	Total	C	N	O	S	0	0
			15	6	1	7	1		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	180	Total	O	0	0
			180	180		
8	B	28	Total	O	0	0
			28	28		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Heparanase



• Molecule 2: Heparanase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	45.99Å 71.06Å 78.29Å 90.00° 95.09° 90.00°	Depositor
Resolution (Å)	52.52 – 1.88 52.52 – 1.88	Depositor EDS
% Data completeness (in resolution range)	99.7 (52.52-1.88) 99.7 (52.52-1.88)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.80 (at 1.88Å)	Xtriage
Refinement program	REFMAC 5.8.0131	Depositor
R, R_{free}	0.175 , 0.215 0.184 , 0.218	Depositor DCC
R_{free} test set	2036 reflections (5.24%)	DCC
Wilson B-factor (Å ²)	31.2	Xtriage
Anisotropy	0.386	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 43.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	3962	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.76% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GNS, BDP, SGN, NAG, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.76	0/3131	0.85	3/4235 (0.1%)
2	B	0.73	0/609	0.86	1/826 (0.1%)
All	All	0.76	0/3740	0.85	4/5061 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	481	ARG	NE-CZ-NH1	6.54	123.57	120.30
1	A	432	ARG	NE-CZ-NH2	5.24	122.92	120.30
1	A	171	ASP	CB-CG-OD1	5.21	122.99	118.30
2	B	85	ARG	NE-CZ-NH1	5.06	122.83	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	425	GLY	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3057	0	3090	9	0
2	B	595	0	604	2	0
3	A	42	0	39	0	0
4	A	1	0	0	0	0
5	A	25	0	14	0	0
6	A	19	0	9	0	0
7	A	15	0	11	0	0
8	A	180	0	0	3	1
8	B	28	0	0	0	0
All	All	3962	0	3767	10	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (10) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:61:ILE:HG12	2:B:92:LEU:HD11	1.84	0.59
1:A:512:MET:HG2	8:A:878:HOH:O	2.07	0.52
1:A:178:ASN:ND2	8:A:704:HOH:O	2.44	0.50
1:A:423:VAL:HG12	2:B:41:LEU:HD23	1.94	0.49
1:A:477:LYS:HE2	1:A:479:LEU:HD22	1.98	0.46
1:A:303:ARG:HD2	8:A:842:HOH:O	2.16	0.46
1:A:198:GLN:HE21	1:A:241:GLN:HE22	1.64	0.45
1:A:383:GLN:HG3	1:A:384:VAL:HG13	1.99	0.44
1:A:315:VAL:O	1:A:318:ILE:HG12	2.18	0.43
1:A:448:GLY:O	1:A:537:ALA:HB3	2.19	0.43

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
8:A:870:HOH:O	8:A:873:HOH:O[2_547]	2.06	0.14

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	383/389 (98%)	373 (97%)	9 (2%)	1 (0%)	44	32
2	B	73/77 (95%)	72 (99%)	1 (1%)	0	100	100
All	All	456/466 (98%)	445 (98%)	10 (2%)	1 (0%)	51	39

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	272	ARG

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	335/338 (99%)	329 (98%)	6 (2%)	64	57
2	B	67/68 (98%)	67 (100%)	0	100	100
All	All	402/406 (99%)	396 (98%)	6 (2%)	70	64

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	159	LYS
1	A	240	SER
1	A	391	TYR
1	A	392	HIS
1	A	522	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	529	TYR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	198	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 1 is monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	A	601	1	14,14,15	0.80	1 (7%)	15,19,21	1.37	1 (6%)
3	NAG	A	602	1	14,14,15	0.53	0	15,19,21	1.35	1 (6%)
3	NAG	A	603	1	14,14,15	0.80	1 (7%)	15,19,21	1.82	5 (33%)
5	BDP	A	605	6	10,13,13	0.89	0	15,19,19	1.33	2 (13%)
6	SGN	A	606	5	18,19,20	2.36	2 (11%)	21,29,31	1.39	4 (19%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	BDP	A	607	7,6	9,12,13	0.71	0	13,17,19	1.51	3 (23%)
7	GNS	A	608	5	14,15,16	2.23	3 (21%)	15,22,24	3.27	7 (46%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	601	1	-	0/6/23/26	0/1/1/1
3	NAG	A	602	1	-	0/6/23/26	0/1/1/1
3	NAG	A	603	1	-	0/6/23/26	0/1/1/1
5	BDP	A	605	6	-	0/0/24/24	0/1/1/1
6	SGN	A	606	5	-	0/10/28/31	0/1/1/1
5	BDP	A	607	7,6	-	0/0/21/24	0/1/1/1
7	GNS	A	608	5	-	0/6/24/27	0/1/1/1

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	603	NAG	C1-C2	2.07	1.55	1.52
3	A	601	NAG	C1-C2	2.16	1.55	1.52
7	A	608	GNS	S1-N	2.65	1.62	1.59
6	A	606	SGN	O2S-S1	4.67	1.48	1.42
7	A	608	GNS	O3S-S1	5.00	1.48	1.42
7	A	608	GNS	O2S-S1	5.75	1.49	1.42
6	A	606	SGN	O1S-S1	8.16	1.52	1.42

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	A	608	GNS	O3S-S1-O2S	-4.46	108.33	120.23
7	A	608	GNS	C4-C3-C2	-4.16	104.93	111.02
3	A	603	NAG	C4-C3-C2	-3.30	106.19	111.02
6	A	606	SGN	O6-C6-C5	-3.24	101.48	107.67
7	A	608	GNS	O5-C1-C2	-3.13	107.12	111.47
3	A	603	NAG	C3-C4-C5	-2.73	105.40	110.22
5	A	607	BDP	O3-C3-C4	-2.44	105.05	110.36
6	A	606	SGN	O2S-S1-O1S	-2.35	113.96	120.23
5	A	607	BDP	O2-C2-C1	-2.27	104.57	109.18
6	A	606	SGN	O1S-S1-N	-2.17	105.63	108.49
7	A	608	GNS	C6-C5-C4	-2.04	108.22	113.00

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	601	NAG	C2-N2-C7	2.02	125.89	122.94
5	A	605	BDP	O4-C4-C3	2.13	115.00	110.36
3	A	603	NAG	O3-C3-C2	2.15	113.99	109.39
3	A	603	NAG	O4-C4-C5	2.40	115.34	109.28
5	A	605	BDP	C1-O5-C5	2.51	115.91	112.22
5	A	607	BDP	C1-C2-C3	2.78	113.17	109.65
7	A	608	GNS	O3-C3-C2	2.79	115.36	109.39
6	A	606	SGN	O2S-S1-N	2.79	112.16	108.49
3	A	603	NAG	C1-O5-C5	3.12	116.46	112.17
3	A	602	NAG	C1-O5-C5	4.12	117.84	112.17
7	A	608	GNS	O2S-S1-N	5.71	116.02	108.49
7	A	608	GNS	C1-O5-C5	7.93	123.09	112.17

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	385/389 (98%)	-0.11	5 (1%) 77 79	23, 38, 65, 113	0
2	B	74/77 (96%)	-0.07	1 (1%) 75 78	27, 40, 64, 95	0
All	All	459/466 (98%)	-0.10	6 (1%) 77 79	23, 38, 65, 113	0

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	159	LYS	3.7
1	A	258	PHE	3.0
2	B	109	GLU	3.0
1	A	427	LYS	2.6
1	A	160	PHE	2.4
1	A	428	ARG	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
5	BDP	A	605	13/13	0.95	0.09	-0.06	31,36,42,42	13
6	SGN	A	606	19/20	0.94	0.11	-0.13	34,40,85,87	19
3	NAG	A	603	14/15	0.81	0.14	-	66,72,80,87	0
3	NAG	A	602	14/15	0.89	0.13	-	59,68,74,77	0
5	BDP	A	607	12/13	0.78	0.18	-	53,61,69,79	12
7	GNS	A	608	15/16	0.70	0.18	-	69,78,104,105	15
3	NAG	A	601	14/15	0.75	0.18	-	60,71,76,78	0
4	CL	A	604	1/1	0.92	0.14	-	66,66,66,66	0

6.5 Other polymers [i](#)

There are no such residues in this entry.