



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 05:47 pm GMT

PDB ID : 3ECV  
Title : Crystal structure of the ALS-related pathological mutant I113T of human apo Cu,Zn Superoxide Dismutase (SOD1)  
Authors : Calderone, V.  
Deposited on : 2008-09-02  
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	5047 (1.90-1.90)
Clashscore	112137	5731 (1.90-1.90)
Ramachandran outliers	110173	5669 (1.90-1.90)
Sidechain outliers	110143	5670 (1.90-1.90)
RSRZ outliers	101464	5100 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	153	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 2%, green 76%, yellow 22%, red 2%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>%</span> <span>76%</span> <span>22%</span> <span>..</span> </div> </div>
1	B	153	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, green 73%, yellow 24%, red 3%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>%</span> <span>73%</span> <span>24%</span> <span>..</span> </div> </div>
1	C	153	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 2%, green 61%, yellow 18%, orange 1%, grey 18%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>%</span> <span>61%</span> <span>18%</span> <span>•</span> <span>18%</span> </div> </div>
1	D	153	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 2%, green 64%, yellow 15%, orange 1%, grey 18%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>2%</span> <span>64%</span> <span>15%</span> <span>..</span> <span>18%</span> </div> </div>

## 2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 4291 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Superoxide dismutase [Cu-Zn].

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	153	Total	C	N	O	S	0	0	0
			1109	677	203	225	4			
1	B	153	Total	C	N	O	S	0	0	0
			1109	677	203	225	4			
1	C	126	Total	C	N	O	S	0	0	0
			914	565	165	180	4			
1	D	125	Total	C	N	O	S	0	0	0
			910	563	164	179	4			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	113	THR	ILE	CONFLICT	UNP P00441
B	113	THR	ILE	CONFLICT	UNP P00441
C	113	THR	ILE	CONFLICT	UNP P00441
D	113	THR	ILE	CONFLICT	UNP P00441


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	71	Total	O	0	0
			71	71		
2	B	78	Total	O	0	0
			78	78		
2	C	49	Total	O	0	0
			49	49		
2	D	51	Total	O	0	0
			51	51		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: Superoxide dismutase [Cu-Zn]




#### • Molecule 1: Superoxide dismutase [Cu-Zn]



#### • Molecule 1: Superoxide dismutase [Cu-Zn]



#### • Molecule 1: Superoxide dismutase [Cu-Zn]



GLY	
LYS	
GLY	
GLY	
ASN	
GLU	
GLU	
SER	
THR	
LYS	
THR	
GLY	
ASN	
ALA	
GLY	
S142	
R143	
L144	
A145	
C146	
Q153	

## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	155.90Å 34.40Å 115.00Å 90.00° 112.10° 90.00°	Depositor
Resolution (Å)	38.95 – 1.90 38.94 – 1.90	Depositor EDS
% Data completeness (in resolution range)	100.0 (38.95-1.90) 98.3 (38.94-1.90)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	0.11	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.77 (at 1.89Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.238 , 0.277 0.238 , 0.275	Depositor DCC
$R_{free}$ test set	4053 reflections (9.98%)	DCC
Wilson B-factor (Å <sup>2</sup> )	12.4	Xtriage
Anisotropy	1.382	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 41.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	4291	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	16.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.75% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.11	1/1127 (0.1%)	1.12	6/1519 (0.4%)
1	B	1.19	4/1127 (0.4%)	1.13	5/1519 (0.3%)
1	C	1.14	1/928 (0.1%)	1.11	5/1252 (0.4%)
1	D	1.08	2/924 (0.2%)	1.04	3/1247 (0.2%)
All	All	1.13	8/4106 (0.2%)	1.10	19/5537 (0.3%)

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	57	CYS	CB-SG	-8.02	1.68	1.82
1	D	6	CYS	CB-SG	-6.16	1.71	1.82
1	D	111	CYS	CB-SG	-5.34	1.73	1.81
1	B	87	VAL	CB-CG2	5.27	1.64	1.52
1	B	27	GLY	C-O	5.26	1.32	1.23
1	B	145	ALA	CA-CB	5.15	1.63	1.52
1	B	132	GLU	CG-CD	5.04	1.59	1.51
1	C	146	CYS	CB-SG	-5.04	1.73	1.81

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	38	LEU	CA-CB-CG	7.73	133.09	115.30
1	C	115	ARG	NE-CZ-NH1	7.73	124.17	120.30
1	B	38	LEU	CA-CB-CG	7.50	132.56	115.30
1	A	115	ARG	NE-CZ-NH1	-7.32	116.64	120.30
1	A	38	LEU	CA-CB-CG	6.99	131.39	115.30
1	B	8	LEU	CA-CB-CG	6.87	131.09	115.30
1	C	42	LEU	CA-CB-CG	6.69	130.69	115.30
1	A	115	ARG	NE-CZ-NH2	6.64	123.62	120.30
1	D	38	LEU	CB-CG-CD1	6.22	121.58	111.00
1	C	53	ASN	CB-CA-C	6.20	122.79	110.40
1	D	42	LEU	CA-CB-CG	5.91	128.89	115.30
1	B	115	ARG	NE-CZ-NH2	-5.85	117.37	120.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	57	CYS	CA-CB-SG	-5.63	103.87	114.00
1	B	9	LYS	CD-CE-NZ	5.56	124.48	111.70
1	B	143	ARG	NE-CZ-NH1	5.54	123.07	120.30
1	D	67	LEU	CA-CB-CG	5.48	127.91	115.30
1	A	5	VAL	CB-CA-C	5.05	120.99	111.40
1	C	90	ASP	CB-CG-OD2	5.03	122.82	118.30
1	A	90	ASP	CB-CG-OD2	5.02	122.82	118.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1109	0	1073	18	0
1	B	1109	0	1073	27	0
1	C	914	0	890	20	0
1	D	910	0	887	20	0
2	A	71	0	0	3	0
2	B	78	0	0	8	0
2	C	49	0	0	1	0
2	D	51	0	0	3	0
All	All	4291	0	3923	84	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (84) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:229:HOH:O	1:C:24:GLU:HG2	1.39	1.20
1:D:46:HIS:HD2	1:D:80:HIS:CE1	1.79	1.01
1:D:46:HIS:CD2	1:D:80:HIS:HE1	1.79	0.99
1:B:77:GLU:HB2	2:B:238:HOH:O	1.64	0.96
1:D:64:PHE:O	1:D:80:HIS:HD2	1.65	0.79

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:63:HIS:HB3	1:D:80:HIS:CD2	2.19	0.78
1:C:79:ARG:NH2	1:C:101:ASP:OD1	2.17	0.78
1:C:38:LEU:O	1:C:93:GLY:HA2	1.88	0.74
1:D:27:GLY:HA3	2:D:165:HOH:O	1.88	0.73
1:B:6:CYS:SG	1:B:147:GLY:O	2.49	0.70
1:A:69:ARG:NH1	1:A:77:GLU:O	2.24	0.70
1:D:110:HIS:HB2	2:D:185:HOH:O	1.92	0.69
1:C:40:GLU:OE1	1:C:91:LYS:HB3	1.94	0.66
2:A:168:HOH:O	1:B:9:LYS:HD3	1.96	0.66
1:A:121:GLU:HB2	1:A:142:SER:HB3	1.79	0.65
1:C:49:GLU:O	1:C:115:ARG:HG2	1.97	0.63
1:B:42:LEU:HB3	1:B:123:ALA:HB2	1.82	0.62
1:B:52:ASP:OD1	1:B:54:THR:HG23	1.99	0.61
1:C:100:GLU:HG2	2:C:176:HOH:O	2.00	0.61
1:D:80:HIS:CD2	1:D:82:GLY:H	2.19	0.60
1:D:46:HIS:CD2	1:D:80:HIS:CE1	2.68	0.58
1:B:26:ASN:HA	1:C:26:ASN:HA	1.86	0.57
1:B:9:LYS:HG3	2:B:183:HOH:O	2.07	0.55
1:B:126:LEU:H	1:B:126:LEU:HD23	1.72	0.55
1:A:39:THR:HG23	2:A:160:HOH:O	2.07	0.54
1:D:80:HIS:HB3	2:D:205:HOH:O	2.08	0.54
1:D:90:ASP:C	1:D:90:ASP:OD1	2.47	0.53
1:D:64:PHE:O	1:D:80:HIS:CD2	2.56	0.52
1:B:126:LEU:N	1:B:126:LEU:HD23	2.26	0.51
1:A:125:ASP:OD1	1:A:125:ASP:C	2.49	0.51
1:A:52:ASP:O	1:A:59:SER:HB2	2.12	0.50
1:A:66:PRO:HD2	1:A:81:VAL:CG2	2.41	0.50
1:C:38:LEU:O	1:C:93:GLY:CA	2.59	0.50
1:D:14:VAL:HG13	1:D:38:LEU:HD13	1.95	0.49
1:A:5:VAL:HG12	1:A:152:ALA:HB2	1.94	0.49
1:A:66:PRO:HD2	1:A:81:VAL:HG23	1.95	0.49
1:B:79:ARG:CD	1:B:80:HIS:O	2.61	0.49
1:B:79:ARG:NH2	1:B:101:ASP:OD1	2.40	0.48
1:B:42:LEU:HB3	1:B:123:ALA:CB	2.42	0.48
1:D:63:HIS:HB3	1:D:80:HIS:NE2	2.28	0.48
1:C:66:PRO:HD2	1:C:81:VAL:CG2	2.43	0.48
1:D:14:VAL:CG1	1:D:38:LEU:HD13	2.44	0.48
1:B:52:ASP:OD1	1:B:52:ASP:C	2.51	0.48
2:B:229:HOH:O	1:C:24:GLU:CG	2.22	0.47
1:C:80:HIS:N	1:C:80:HIS:CD2	2.83	0.47
1:C:79:ARG:HH22	1:C:101:ASP:CG	2.13	0.47

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:121:GLU:HB3	1:C:141:GLY:HA3	1.97	0.47
1:D:117:LEU:O	1:D:146:CYS:HA	2.15	0.47
1:B:79:ARG:HH22	1:B:101:ASP:CG	2.18	0.47
1:A:121:GLU:HB2	1:A:142:SER:CB	2.44	0.47
1:A:52:ASP:O	1:A:59:SER:CB	2.62	0.46
1:D:45:PHE:O	1:D:84:LEU:HB2	2.16	0.46
1:B:15:GLN:NE2	1:B:36:LYS:HE2	2.29	0.46
1:C:14:VAL:HG21	1:C:144:LEU:HB3	1.97	0.46
1:B:105:SER:O	1:B:111:CYS:HA	2.16	0.46
1:B:46:HIS:CE1	2:B:193:HOH:O	2.68	0.46
1:B:100:GLU:OE1	2:B:239:HOH:O	2.21	0.45
1:A:38:LEU:O	1:A:93:GLY:HA2	2.17	0.45
1:A:41:GLY:O	1:A:89:ALA:N	2.43	0.45
1:A:117:LEU:O	1:A:146:CYS:HA	2.17	0.44
1:B:79:ARG:HD3	1:B:80:HIS:O	2.17	0.44
1:C:101:ASP:HB3	1:C:104:ILE:HG12	1.98	0.44
1:A:84:LEU:O	2:A:204:HOH:O	2.21	0.44
1:A:43:HIS:HB3	1:A:120:HIS:O	2.17	0.44
1:B:20:PHE:CD1	1:B:20:PHE:N	2.85	0.44
1:D:63:HIS:HB3	1:D:80:HIS:CG	2.52	0.43
1:A:14:VAL:HG21	1:A:144:LEU:HB3	1.99	0.43
1:D:121:GLU:HA	1:D:144:LEU:HD11	2.01	0.43
1:C:46:HIS:CD2	1:C:83:ASP:HA	2.53	0.43
1:A:116:THR:CG2	1:A:146:CYS:HB2	2.48	0.43
1:B:31:VAL:HB	1:B:99:ILE:HB	1.99	0.43
1:B:74:PRO:HG2	2:B:216:HOH:O	2.18	0.43
1:B:118:VAL:HG11	1:B:143:ARG:HG2	2.01	0.43
1:B:40:GLU:HG3	1:B:90:ASP:O	2.18	0.42
1:A:71:HIS:NE2	1:A:124:ASP:OD1	2.43	0.42
1:D:14:VAL:HG21	1:D:144:LEU:HB3	2.02	0.42
1:C:31:VAL:HB	1:C:99:ILE:HB	2.03	0.41
1:C:28:PRO:HB3	1:C:102:SER:OG	2.20	0.41
1:B:125:ASP:OD1	1:B:134:SER:OG	2.38	0.41
1:C:141:GLY:C	1:C:142:SER:O	2.58	0.41
1:D:65:ASN:OD1	1:D:80:HIS:HA	2.20	0.41
1:C:35:ILE:CG2	1:C:38:LEU:HD21	2.50	0.41
1:B:85:GLY:N	2:B:162:HOH:O	2.54	0.40
1:B:79:ARG:HD2	1:B:80:HIS:O	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	151/153 (99%)	147 (97%)	4 (3%)	0	100	100
1	B	151/153 (99%)	143 (95%)	8 (5%)	0	100	100
1	C	120/153 (78%)	112 (93%)	8 (7%)	0	100	100
1	D	119/153 (78%)	117 (98%)	2 (2%)	0	100	100
All	All	541/612 (88%)	519 (96%)	22 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	118/118 (100%)	109 (92%)	9 (8%)	15	6
1	B	118/118 (100%)	109 (92%)	9 (8%)	15	6
1	C	98/118 (83%)	92 (94%)	6 (6%)	22	11
1	D	98/118 (83%)	88 (90%)	10 (10%)	8	3
All	All	432/472 (92%)	398 (92%)	34 (8%)	14	6

All (34) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2	THR
1	A	8	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	24	GLU
1	A	25	SER
1	A	31	VAL
1	A	38	LEU
1	A	98	SER
1	A	128	LYS
1	A	142	SER
1	B	8	LEU
1	B	9	LYS
1	B	34	SER
1	B	38	LEU
1	B	75	LYS
1	B	79	ARG
1	B	91	LYS
1	B	98	SER
1	B	122	LYS
1	C	36	LYS
1	C	53	ASN
1	C	79	ARG
1	C	91	LYS
1	C	122	LYS
1	C	142	SER
1	D	24	GLU
1	D	31	VAL
1	D	38	LEU
1	D	42	LEU
1	D	62	PRO
1	D	67	LEU
1	D	80	HIS
1	D	91	LYS
1	D	96	ASP
1	D	105	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	15	GLN
1	C	46	HIS
1	C	53	ASN
1	C	80	HIS
1	D	46	HIS
1	D	80	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	153/153 (100%)	-0.17	1 (0%) 87 89	4, 15, 27, 39	0
1	B	153/153 (100%)	-0.15	0 100 100	3, 14, 26, 34	0
1	C	126/153 (82%)	0.00	2 (1%) 72 75	3, 16, 32, 41	2 (1%)
1	D	125/153 (81%)	-0.03	3 (2%) 59 63	6, 17, 31, 43	0
All	All	557/612 (91%)	-0.09	6 (1%) 80 83	3, 16, 30, 43	2 (0%)

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	67	LEU	4.8
1	C	141	GLY	4.1
1	A	1	ALA	3.3
1	D	1	ALA	2.9
1	C	79	ARG	2.8
1	D	79	ARG	2.3

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.