



# Full wwPDB X-ray Structure Validation Report ⓘ

Sep 3, 2017 – 09:23 AM EDT

PDB ID : 5ED2  
Title : Human Adenosine Deaminase Acting on dsRNA (ADAR2) mutant E488Q  
bound to dsRNA sequence derived from human GLI1 gene  
Authors : Matthews, M.M.; Fisher, A.J.  
Deposited on : unknown  
Resolution : 2.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20029824  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20029824


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R <sub>free</sub>	100719	2395 (3.00-2.92)
Clashscore	112137	2773 (3.00-2.92)
Ramachandran outliers	110173	2680 (3.00-2.92)
Sidechain outliers	110143	2683 (3.00-2.92)
RSRZ outliers	101464	2421 (3.00-2.92)
RNA backbone	2435	1008 (3.30-2.62)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	403	<div><div>84%</div><div>9%</div><div>• •</div></div>
1	D	403	<div><div>82%</div><div>12%</div><div>• •</div></div>
2	B	23	<div><div>83%</div><div>17%</div></div>
2	E	23	<div><div>83%</div><div>17%</div></div>

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	C	23	 96% .
3	F	23	 96% .

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 8063 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Double-stranded RNA-specific editase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	385	Total	C	N	O	S	0	0	0
			3019	1903	550	555	11			
1	D	385	Total	C	N	O	S	0	1	0
			3020	1903	550	555	12			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	488	GLN	GLU	engineered mutation	UNP P78563
D	488	GLN	GLU	engineered mutation	UNP P78563


- Molecule 2 is a RNA chain called RNA (5'-R(P\*GP\*CP\*UP\*CP\*GP\*CP\*GP\*AP\*UP\*GP\*CP\*UP\*(8AZ)P\*GP\*AP\*GP\*GP\*GP\*CP\*UP\*CP\*UP\*G)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	23	Total	C	N	O	P	0	0	0
			490	218	88	162	22			
2	E	23	Total	C	N	O	P	0	0	0
			490	218	88	162	22			

- Molecule 3 is a RNA chain called RNA (5'-R(P\*CP\*AP\*GP\*AP\*GP\*CP\*CP\*CP\*CP\*CP\*CP\*AP\*GP\*CP\*AP\*UP\*CP\*GP\*CP\*GP\*AP\*GP\*C)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	23	Total	C	N	O	P	0	0	0
			485	218	90	155	22			
3	F	23	Total	C	N	O	P	0	0	0
			485	218	90	155	22			

- Molecule 4 is INOSITOL HEXAKISPHOSPHATE (three-letter code: IHP) (formula: C<sub>6</sub>H<sub>18</sub>O<sub>24</sub>P<sub>6</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	O	P	0	0
			36	6	24	6		
4	D	1	Total	C	O	P	0	0
			36	6	24	6		

- Molecule 5 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Zn	0	0
			1	1		
5	D	1	Total	Zn	0	0
			1	1		



- Molecule 1: Double-stranded RNA-specific editase 1





● Molecule 3: RNA (5'-R(P\*CP\*AP\*GP\*AP\*GP\*CP\*CP\*CP\*CP\*CP\*CP\*AP\*GP\*CP\*AP\*U  
P\*CP\*GP\*CP\*GP\*AP\*GP\*C)-3')

Chain C: 96%



● Molecule 3: RNA (5'-R(P\*CP\*AP\*GP\*AP\*GP\*CP\*CP\*CP\*CP\*CP\*CP\*AP\*GP\*CP\*AP\*U  
P\*CP\*GP\*CP\*GP\*AP\*GP\*C)-3')

Chain F: 96%



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	79.13Å 81.61Å 256.62Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.56 – 2.95 39.56 – 2.95	Depositor EDS
% Data completeness (in resolution range)	98.2 (39.56-2.95) 98.3 (39.56-2.95)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.49 (at 2.95Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, $R_{free}$	0.188 , 0.207 0.198 , 0.220	Depositor DCC
$R_{free}$ test set	1743 reflections (5.20%)	DCC
Wilson B-factor (Å <sup>2</sup> )	74.6	Xtriage
Anisotropy	0.123	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.29 , 44.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.021 for k,h,-l	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	8063	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	94.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.36% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, IHP, 8AZ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.29	0/3081	0.46	0/4165
1	D	0.28	0/3087	0.46	0/4173
2	B	0.28	0/521	0.78	0/809
2	E	0.26	0/521	0.75	0/809
3	C	0.28	0/541	0.73	0/841
3	F	0.28	0/541	0.73	0/841
All	All	0.28	0/8292	0.55	0/11638

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3019	0	3043	29	0
1	D	3020	0	3044	30	0
2	B	490	0	239	2	0
2	E	490	0	239	2	0
3	C	485	0	254	1	0
3	F	485	0	254	1	0
4	A	36	0	6	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	D	36	0	6	0	0
5	A	1	0	0	0	0
5	D	1	0	0	0	0
All	All	8063	0	7085	59	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:448:THR:OG1	1:D:455:ARG:NH2	2.28	0.66
1:D:463:ILE:HD13	1:D:465:GLU:HG3	1.79	0.65
1:A:502:TRP:HZ3	1:A:694:GLN:HG2	1.61	0.65
1:A:485:GLU:HB3	1:A:510:ARG:HG3	1.81	0.62
1:D:400:ARG:HD3	1:D:523:TRP:CE2	2.35	0.62
1:D:485:GLU:HB3	1:D:510:ARG:HG3	1.82	0.61
1:D:621:LEU:HD12	1:D:623:ARG:HH21	1.67	0.60
1:A:653:THR:HG23	1:A:654:LYS:H	1.67	0.59
1:A:355:VAL:HG11	1:A:403:LEU:HD22	1.85	0.58
1:D:376:LYS:NZ	2:E:15:A:OP2	2.26	0.57
1:A:455:ARG:NH1	2:B:13:8AZ:OP1	2.37	0.57
1:A:463:ILE:HD11	1:A:466:GLU:HB3	1.87	0.56
1:A:348:ARG:NH1	3:C:3:G:OP1	2.38	0.56
1:D:355:VAL:HG11	1:D:403:LEU:HD22	1.88	0.56
1:D:502:TRP:HZ3	1:D:694:GLN:HG2	1.71	0.55
1:A:400:ARG:HD3	1:A:523:TRP:CE2	2.42	0.55
1:D:652:ILE:O	1:D:654:LYS:N	2.39	0.55
1:A:566:ASN:H	1:A:566:ASN:HD22	1.56	0.54
1:D:477:ARG:NH1	3:F:15:A:OP1	2.40	0.54
1:A:629:LYS:HE3	1:A:694:GLN:NE2	2.24	0.52
1:A:463:ILE:HD12	1:A:466:GLU:H	1.75	0.52
1:D:420:ASP:OD1	1:D:423:ARG:NH1	2.40	0.51
1:A:558:ARG:HA	1:A:562:GLN:HB3	1.93	0.51
1:A:485:GLU:O	1:A:510:ARG:NH1	2.44	0.49
1:D:504:GLY:O	1:D:509:GLU:HB2	2.11	0.49
1:D:392:ASP:OD2	1:D:483:LYS:NZ	2.46	0.49
1:A:621:LEU:HD12	1:A:623:ARG:HH21	1.77	0.48
1:D:629:LYS:HE3	1:D:694:GLN:NE2	2.28	0.48
1:A:522:ARG:O	1:A:526:VAL:HG22	2.12	0.48
1:D:359:THR:HG22	1:D:440:VAL:HG22	1.95	0.48

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:394:HIS:CE1	1:A:483:LYS:HE2	2.50	0.47
1:A:652:ILE:O	1:A:654:LYS:N	2.48	0.47
1:A:462:PRO:HG3	1:A:551:TYR:O	2.15	0.47
1:D:522:ARG:O	1:D:526:VAL:HG22	2.15	0.46
1:D:394:HIS:CE1	1:D:483:LYS:HE2	2.51	0.45
1:A:323:LEU:O	1:A:327:VAL:HG13	2.18	0.44
1:A:514:MET:HG3	1:A:687:TRP:CE2	2.53	0.44
1:D:600:VAL:HG13	1:D:609:ILE:HB	2.00	0.43
1:A:390:LEU:HD12	1:A:612:ILE:HG21	2.00	0.43
1:D:451:CYS:HB3	2:E:13:8AZ:N7	2.33	0.43
1:A:471:HIS:N	1:A:472:PRO:HD3	2.34	0.43
1:D:554:ASP:OD1	1:D:554:ASP:N	2.50	0.43
1:A:475:LYS:O	1:A:479:GLN:NE2	2.51	0.43
1:A:448:THR:OG1	1:A:550:LEU:HD12	2.17	0.43
1:A:376:LYS:NZ	2:B:15:A:OP2	2.30	0.43
1:D:396:GLU:N	1:D:396:GLU:OE2	2.40	0.43
1:A:458:SER:OG	1:A:458:SER:O	2.37	0.42
1:D:420:ASP:HA	1:D:423:ARG:NH1	2.34	0.42
1:D:350:LYS:NZ	1:D:591:GLN:O	2.47	0.42
1:D:532:LEU:HB3	1:D:636:TRP:CD1	2.55	0.42
1:A:590:ARG:HD2	1:A:590:ARG:HA	1.82	0.42
1:D:418:LYS:HA	1:D:421:GLN:HG3	2.01	0.41
1:D:514:MET:HG3	1:D:687:TRP:CE2	2.55	0.41
1:D:494:ARG:HB2	1:D:497:ALA:HB2	2.03	0.41
1:A:566:ASN:H	1:A:566:ASN:ND2	2.18	0.41
1:D:451:CYS:HA	1:D:455:ARG:HG2	2.02	0.40
1:D:470:ARG:HG2	1:D:471:HIS:CD2	2.56	0.40
1:A:479:GLN:HE21	1:A:479:GLN:HB2	1.69	0.40
1:D:635:ARG:O	1:D:639:VAL:HG23	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	383/403 (95%)	364 (95%)	18 (5%)	1 (0%)	44	79
1	D	384/403 (95%)	366 (95%)	17 (4%)	1 (0%)	44	79
All	All	767/806 (95%)	730 (95%)	35 (5%)	2 (0%)	44	79

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	653	THR
1	D	653	THR

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	328/347 (94%)	319 (97%)	9 (3%)	50	81
1	D	329/347 (95%)	323 (98%)	6 (2%)	64	87
All	All	657/694 (95%)	642 (98%)	15 (2%)	56	84

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	327	VAL
1	A	413	LEU
1	A	455	ARG
1	A	463	ILE
1	A	510	ARG
1	A	566	ASN
1	A	669	GLN
1	A	684	LEU
1	A	694	GLN
1	D	413	LEU
1	D	455	ARG
1	D	491	ILE
1	D	510	ARG
1	D	684	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	D	694	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	318	HIS
1	A	321	GLN
1	A	479	GLN
1	A	566	ASN
1	A	659	HIS
1	A	694	GLN
1	D	479	GLN
1	D	694	GLN

### 5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	B	21/23 (91%)	2 (9%)	0
2	E	21/23 (91%)	2 (9%)	0
3	C	22/23 (95%)	0	0
3	F	22/23 (95%)	0	0
All	All	86/92 (93%)	4 (4%)	0

All (4) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
2	B	5	G
2	B	7	G
2	E	5	G
2	E	7	G

There are no RNA pucker outliers to report.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link

column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	8AZ	B	13	2,5	16,24,25	0.79	0	15,35,38	0.60	0
2	8AZ	E	13	2,5	16,24,25	0.77	0	15,35,38	0.65	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	8AZ	B	13	2,5	-	0/3/35/36	0/3/3/3
2	8AZ	E	13	2,5	-	0/3/35/36	0/3/3/3

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	13	8AZ	1	0
2	E	13	8AZ	1	0

## 5.5 Carbohydrates

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length

(or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	IHP	A	801	-	36,36,36	0.70	0	54,60,60	0.99	0
4	IHP	D	801	-	36,36,36	0.70	0	54,60,60	1.02	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	IHP	A	801	-	-	0/30/54/54	0/1/1/1
4	IHP	D	801	-	-	0/30/54/54	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	385/403 (95%)	0.06	6 (1%) 72 53	51, 90, 138, 174	0
1	D	385/403 (95%)	-0.01	4 (1%) 82 66	50, 85, 124, 153	0
2	B	22/23 (95%)	-0.17	0 100 100	78, 116, 128, 133	0
2	E	22/23 (95%)	-0.15	0 100 100	77, 109, 137, 137	0
3	C	23/23 (100%)	-0.14	0 100 100	78, 109, 125, 127	0
3	F	23/23 (100%)	-0.21	0 100 100	76, 108, 135, 146	0
All	All	860/898 (95%)	0.01	10 (1%) 79 61	50, 89, 135, 174	0

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	496	ASN	4.4
1	D	506	LEU	3.1
1	A	652	ILE	3.1
1	A	496	ASN	2.2
1	A	495	SER	2.2
1	A	653	THR	2.2
1	D	470	ARG	2.2
1	A	466	GLU	2.1
1	D	590	ARG	2.1
1	A	506	LEU	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy



less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	8AZ	E	13	22/23	0.98	0.21	-	56,62,66,68	0
2	8AZ	B	13	22/23	0.98	0.23	-	55,64,70,71	0

### 6.3 Carbohydrates

There are no carbohydrates in this entry.

### 6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
4	IHP	D	801	36/36	0.98	0.24	0.77	39,63,79,88	0
4	IHP	A	801	36/36	0.98	0.26	0.20	47,64,96,100	0
5	ZN	D	802	1/1	0.99	0.21	-0.79	63,63,63,63	0
5	ZN	A	802	1/1	0.98	0.25	-1.43	65,65,65,65	0

### 6.5 Other polymers

There are no such residues in this entry.