



# wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 4, 2017 – 11:31 PM EDT

PDB ID : 4EE7  
Title : Crystal Structure of the Novel Phenazine Prenyltransferase EpzP in complex with S-thiolodiphosphate (methylated)  
Authors : Zocher, G.; Stehle, T.  
Deposited on : unknown  
Resolution : 1.67 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20030345  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20030345


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.67 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	5252 (1.70-1.66)
Clashscore	112137	5803 (1.70-1.66)
Ramachandran outliers	110173	5704 (1.70-1.66)
Sidechain outliers	110143	5703 (1.70-1.66)
RSRZ outliers	101464	5298 (1.70-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	304	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 1%, green 91%, grey 5%);"></div> <div style="display: flex; justify-content: space-between; padding: 0 5px;"> <span>1%</span> <span>91%</span> <span>5%</span> </div> </div>
1	B	304	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 2%, orange 2%, yellow 7%, green 88%, grey 5%);"></div> <div style="display: flex; justify-content: space-between; padding: 0 5px;"> <span>2%</span> <span>88%</span> <span>7%</span> <span>5%</span> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	PIS	A	401	-	-	-	X
3	PG4	A	402	-	-	-	X
4	SO4	B	403	-	-	X	-
5	CL	A	406	-	-	-	X

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 5237 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Prenyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	290	Total	C	N	O	S	0	6	0
			2273	1465	365	435	8			
1	B	288	Total	C	N	O	S	0	5	0
			2247	1450	362	427	8			

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	EXPRESSION TAG	UNP E5KWG9
A	0	SER	-	EXPRESSION TAG	UNP E5KWG9
B	-1	GLY	-	EXPRESSION TAG	UNP E5KWG9
B	0	SER	-	EXPRESSION TAG	UNP E5KWG9

- Molecule 2 is TRIHYDROGEN THIODIPHOSPHATE (three-letter code: PIS) (formula:  $\text{H}_3\text{O}_6\text{P}_2\text{S}$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	O	P	S	0	0
			9	6	2	1		

- Molecule 3 is TETRAETHYLENE GLYCOL (three-letter code: PG4) (formula:  $C_8H_{18}O_5$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			7	4	3		
3	A	1	Total	C	O	0	0
			13	8	5		
3	B	1	Total	C	O	0	0
			10	6	4		
3	B	1	Total	C	O	0	0
			13	8	5		

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula:  $O_4S$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	B	1	Total	O	S	0	0
			5	4	1		
4	B	1	Total	O	S	0	0
			5	4	1		
4	B	1	Total	O	S	0	0
			5	4	1		
4	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 5 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Cl	0	0
			1	1		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	320	Total	O	0	0
			320	320		
6	B	314	Total	O	0	0
			314	314		



- Molecule 1: Prenyltransferase



- |     |      |
|-----|------|
| ARG | GLY  |
| SER | A5   |
| ASP | D6   |
| ASP | L7   |
| GLY | K31  |
|     | L34  |
|     | R49  |
|     | C61  |
|     | R62  |
|     | K68  |
|     | D66  |
|     | L93  |
|     | D108 |
|     | V111 |
|     | W119 |
|     | K152 |
|     | R169 |
|     | G173 |
|     | G186 |
|     | G192 |
|     | E199 |
|     | K258 |
|     | T264 |
|     | E265 |
|     | G266 |
|     | R267 |
|     | M268 |
|     | M290 |
|     | Q291 |
|     | K392 |
|     | ARG  |
|     | VAL  |
|     | GLU  |
|     | LYS  |
|     | LEU  |
|     | ILE  |

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	42.11Å 97.01Å 135.86Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.39 – 1.67 29.39 – 1.67	Depositor EDS
% Data completeness (in resolution range)	100.0 (29.39-1.67) 99.9 (29.39-1.67)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.29 (at 1.67Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, $R_{free}$	0.188 , 0.224 0.195 , 0.229	Depositor DCC
$R_{free}$ test set	2619 reflections (4.17%)	DCC
Wilson B-factor (Å <sup>2</sup> )	11.6	Xtriage
Anisotropy	0.233	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 43.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	5237	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	15.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 50.01 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 6.8807e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PIS, PG4, MLY, SO4, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.55	0/2294	0.65	0/3129
1	B	0.54	0/2266	0.67	1/3091 (0.0%)
All	All	0.54	0/4560	0.66	1/6220 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	B	169	ARG	NE-CZ-NH1	-5.25	117.68	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	264	THR	Peptide

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2273	0	2236	10	0
1	B	2247	0	2216	12	0
2	A	9	0	0	2	0
3	A	20	0	26	1	0
3	B	23	0	31	2	0
4	A	10	0	0	0	0
4	B	20	0	0	2	0
5	A	1	0	0	0	0
6	A	320	0	0	1	0
6	B	314	0	0	0	1
All	All	5237	0	4509	22	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 22 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:61[B]:CYS:SG	1:A:111:VAL:CG1	2.80	0.69
1:A:39:ASP:OD2	6:A:788:HOH:O	2.16	0.62
1:A:61[B]:CYS:SG	1:A:111:VAL:HG12	2.44	0.58
1:A:49:ARG:HD2	2:A:401:PIS:S1	2.48	0.54
1:B:61[B]:CYS:SG	1:B:111:VAL:CG1	2.97	0.53

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:B:515:HOH:O	6:B:755:HOH:O[1_655]	1.88	0.32

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	289/304 (95%)	283 (98%)	5 (2%)	1 (0%)	44	24
1	B	286/304 (94%)	281 (98%)	4 (1%)	1 (0%)	44	24
All	All	575/608 (95%)	564 (98%)	9 (2%)	2 (0%)	38	24

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	265	GLU
1	B	265	GLU

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	242/251 (96%)	241 (100%)	1 (0%)	93	89
1	B	239/251 (95%)	234 (98%)	5 (2%)	59	37
All	All	481/502 (96%)	475 (99%)	6 (1%)	73	62

5 of 6 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	62	ARG
1	B	267	ARG
1	B	86	ASP
1	B	49	ARG
1	B	258	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	19	GLN
1	A	79	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

10 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
1	MLY	A	126	1	10,10,11	0.51	0	8,11,13	0.84	0
1	MLY	A	16	1	10,10,11	0.63	0	8,11,13	0.88	0
1	MLY	A	283	1	10,10,11	0.59	0	8,11,13	1.19	1 (12%)
1	MLY	A	68	1	10,10,11	0.48	0	8,11,13	0.99	0
1	MLY	A	84	1	10,10,11	0.69	0	8,11,13	1.74	3 (37%)
1	MLY	B	126	1	10,10,11	0.53	0	8,11,13	0.88	0
1	MLY	B	16	1	10,10,11	0.49	0	8,11,13	0.71	0
1	MLY	B	283	1	9,9,11	0.45	0	6,9,13	1.78	2 (33%)
1	MLY	B	68	1	10,10,11	0.48	0	8,11,13	1.35	0
1	MLY	B	84	1	10,10,11	0.84	0	8,11,13	1.48	2 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	MLY	A	126	1	-	0/7/9/11	0/0/0/0
1	MLY	A	16	1	-	0/7/9/11	0/0/0/0
1	MLY	A	283	1	-	0/7/9/11	0/0/0/0
1	MLY	A	68	1	-	0/7/9/11	0/0/0/0
1	MLY	A	84	1	-	0/7/9/11	0/0/0/0
1	MLY	B	126	1	-	0/7/9/11	0/0/0/0
1	MLY	B	16	1	-	0/7/9/11	0/0/0/0
1	MLY	B	283	1	-	0/6/8/11	0/0/0/0
1	MLY	B	68	1	-	0/7/9/11	0/0/0/0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	MLY	B	84	1	-	0/7/9/11	0/0/0/0

There are no bond length outliers.

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	283	MLY	CB-CA-C	-3.51	105.86	111.65
1	A	283	MLY	CB-CA-C	-2.32	107.83	111.65
1	B	283	MLY	O-C-CA	-2.13	119.14	125.02
1	A	84	MLY	CB-CA-C	-2.10	108.19	111.65
1	B	84	MLY	CD-CE-NZ	2.05	119.35	113.77

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	B	68	MLY	2	0

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 1 is monoatomic - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	PIS	A	401	-	5,8,8	2.90	3 (60%)	6,13,13	2.46	1 (16%)
3	PG4	A	402	-	6,6,12	0.46	0	5,5,11	0.25	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	PG4	A	403	-	12,12,12	0.69	0	11,11,11	0.51	0
4	SO4	A	404	-	4,4,4	0.23	0	6,6,6	0.31	0
4	SO4	A	405	-	4,4,4	0.27	0	6,6,6	0.26	0
3	PG4	B	401	-	9,9,12	0.48	0	8,8,11	0.51	0
3	PG4	B	402	-	12,12,12	0.46	0	11,11,11	0.42	0
4	SO4	B	403	-	4,4,4	0.21	0	6,6,6	0.40	0
4	SO4	B	404	-	4,4,4	0.34	0	6,6,6	0.32	0
4	SO4	B	405	-	4,4,4	0.25	0	6,6,6	0.43	0
4	SO4	B	406	-	4,4,4	0.28	0	6,6,6	0.30	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PIS	A	401	-	-	0/4/6/6	0/0/0/0
3	PG4	A	402	-	-	0/4/4/10	0/0/0/0
3	PG4	A	403	-	-	0/10/10/10	0/0/0/0
4	SO4	A	404	-	-	0/0/0/0	0/0/0/0
4	SO4	A	405	-	-	0/0/0/0	0/0/0/0
3	PG4	B	401	-	-	0/7/7/10	0/0/0/0
3	PG4	B	402	-	-	0/10/10/10	0/0/0/0
4	SO4	B	403	-	-	0/0/0/0	0/0/0/0
4	SO4	B	404	-	-	0/0/0/0	0/0/0/0
4	SO4	B	405	-	-	0/0/0/0	0/0/0/0
4	SO4	B	406	-	-	0/0/0/0	0/0/0/0

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	PIS	P2-O4	-3.36	1.48	1.56
2	A	401	PIS	P1-O5	2.72	1.60	1.50
2	A	401	PIS	P1-O1	4.49	1.67	1.60

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	PIS	O4-P2-O6	5.50	121.53	109.87

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	401	PIS	2	0
3	A	403	PG4	1	0
3	B	401	PG4	2	0
4	B	403	SO4	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	285/304 (93%)	-0.26	3 (1%) 80 84	6, 12, 24, 42	0
1	B	283/304 (93%)	-0.26	6 (2%) 64 68	6, 12, 27, 45	0
All	All	568/608 (93%)	-0.26	9 (1%) 72 76	6, 12, 26, 45	0

The worst 5 of 9 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	292	LYS	3.7
1	A	265	GLU	3.6
1	A	290	TRP	3.1
1	B	5	ALA	2.9
1	B	178	GLY	2.7

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
1	MLY	B	68	11/12	0.92	0.12	-	13,15,21,22	0
1	MLY	B	16	11/12	0.95	0.11	-	20,25,30,32	0
1	MLY	A	68	11/12	0.96	0.07	-	9,11,13,14	0
1	MLY	A	16	11/12	0.95	0.10	-	15,18,30,31	0
1	MLY	A	126	11/12	0.93	0.10	-	10,14,25,26	0
1	MLY	A	84	11/12	0.94	0.10	-	9,11,16,19	0
1	MLY	B	84	11/12	0.93	0.10	-	10,14,19,20	0
1	MLY	A	283	11/12	0.95	0.14	-	7,13,27,29	0

*Continued on next page...*



*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
1	MLY	B	283	10/12	0.96	0.10	-	7,10,20,22	0
1	MLY	B	126	11/12	0.94	0.10	-	12,13,21,23	0

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	PG4	A	402	7/13	0.87	0.14	4.01	23,23,26,28	0
2	PIS	A	401	9/9	0.90	0.14	3.39	12,21,30,31	9
5	CL	A	406	1/1	0.99	0.10	3.18	20,20,20,20	0
4	SO4	B	405	5/5	0.94	0.14	1.58	27,28,29,33	0
4	SO4	A	404	5/5	0.98	0.13	1.53	24,25,29,33	0
3	PG4	B	402	13/13	0.86	0.15	1.33	20,26,36,38	0
3	PG4	A	403	13/13	0.80	0.14	1.19	21,27,31,36	0
3	PG4	B	401	10/13	0.79	0.15	0.60	21,26,31,31	0
4	SO4	B	403	5/5	0.95	0.22	-	22,28,32,37	0
4	SO4	A	405	5/5	0.93	0.22	-	18,28,32,35	0
4	SO4	B	404	5/5	0.93	0.27	-	31,32,34,35	0
4	SO4	B	406	5/5	0.94	0.41	-	31,33,34,37	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.