



wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 15, 2018 – 05:34 AM EST

PDB ID : 3EH5
Title : Structure of the reduced form of cytochrome ba3 oxidase from *Thermus thermophilus*
Authors : Liu, B.; Chen, Y.; Doukov, T.; Soltis, S.M.; Stout, D.; Fee, J.A.
Deposited on : 2008-09-11
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20030736
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20030736


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	2583 (2.80-2.80)
Clashscore	112137	3033 (2.80-2.80)
Ramachandran outliers	110173	2983 (2.80-2.80)
Sidechain outliers	110143	2985 (2.80-2.80)
RSRZ outliers	101464	2610 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	618	<div> <div>7%</div> <div>55%</div> <div>29%</div> <div>5%</div> <div>10%</div> </div>
2	B	166	<div> <div>7%</div> <div>68%</div> <div>27%</div> <div>5%</div> </div>
3	C	33	<div> <div>6%</div> <div>73%</div> <div>27%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
7	BNG	A	807	-	-	-	X
7	BNG	C	804	-	-	-	X

2 Entry composition

There are 9 unique types of molecules in this entry. The entry contains 6244 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cytochrome c oxidase subunit 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	557	4409	2985	709	699	16	0	0	0

There are 58 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-55	SER	-	EXPRESSION TAG	UNP Q5SJ79
A	-54	GLU	-	EXPRESSION TAG	UNP Q5SJ79
A	-53	ILE	-	EXPRESSION TAG	UNP Q5SJ79
A	-52	SER	-	EXPRESSION TAG	UNP Q5SJ79
A	-51	ARG	-	EXPRESSION TAG	UNP Q5SJ79
A	-50	VAL	-	EXPRESSION TAG	UNP Q5SJ79
A	-49	TYR	-	EXPRESSION TAG	UNP Q5SJ79
A	-48	GLU	-	EXPRESSION TAG	UNP Q5SJ79
A	-47	ALA	-	EXPRESSION TAG	UNP Q5SJ79
A	-46	TYR	-	EXPRESSION TAG	UNP Q5SJ79
A	-45	PRO	-	EXPRESSION TAG	UNP Q5SJ79
A	-44	GLU	-	EXPRESSION TAG	UNP Q5SJ79
A	-43	LYS	-	EXPRESSION TAG	UNP Q5SJ79
A	-42	LYS	-	EXPRESSION TAG	UNP Q5SJ79
A	-41	ALA	-	EXPRESSION TAG	UNP Q5SJ79
A	-40	THR	-	EXPRESSION TAG	UNP Q5SJ79
A	-39	LEU	-	EXPRESSION TAG	UNP Q5SJ79
A	-38	TYR	-	EXPRESSION TAG	UNP Q5SJ79
A	-37	PHE	-	EXPRESSION TAG	UNP Q5SJ79
A	-36	LEU	-	EXPRESSION TAG	UNP Q5SJ79
A	-35	VAL	-	EXPRESSION TAG	UNP Q5SJ79
A	-34	LEU	-	EXPRESSION TAG	UNP Q5SJ79
A	-33	GLY	-	EXPRESSION TAG	UNP Q5SJ79
A	-32	PHE	-	EXPRESSION TAG	UNP Q5SJ79
A	-31	LEU	-	EXPRESSION TAG	UNP Q5SJ79
A	-30	ALA	-	EXPRESSION TAG	UNP Q5SJ79
A	-29	LEU	-	EXPRESSION TAG	UNP Q5SJ79

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	-28	ILE	-	EXPRESSION TAG	UNP Q5SJ79
A	-27	VAL	-	EXPRESSION TAG	UNP Q5SJ79
A	-26	GLY	-	EXPRESSION TAG	UNP Q5SJ79
A	-25	SER	-	EXPRESSION TAG	UNP Q5SJ79
A	-24	LEU	-	EXPRESSION TAG	UNP Q5SJ79
A	-23	PHE	-	EXPRESSION TAG	UNP Q5SJ79
A	-22	GLY	-	EXPRESSION TAG	UNP Q5SJ79
A	-21	PRO	-	EXPRESSION TAG	UNP Q5SJ79
A	-20	PHE	-	EXPRESSION TAG	UNP Q5SJ79
A	-19	GLN	-	EXPRESSION TAG	UNP Q5SJ79
A	-18	ALA	-	EXPRESSION TAG	UNP Q5SJ79
A	-17	LEU	-	EXPRESSION TAG	UNP Q5SJ79
A	-16	ASN	-	EXPRESSION TAG	UNP Q5SJ79
A	-15	TYR	-	EXPRESSION TAG	UNP Q5SJ79
A	-14	GLY	-	EXPRESSION TAG	UNP Q5SJ79
A	-13	ASN	-	EXPRESSION TAG	UNP Q5SJ79
A	-12	VAL	-	EXPRESSION TAG	UNP Q5SJ79
A	-11	ASP	-	EXPRESSION TAG	UNP Q5SJ79
A	-10	ALA	-	EXPRESSION TAG	UNP Q5SJ79
A	-9	TYR	-	EXPRESSION TAG	UNP Q5SJ79
A	-8	PRO	-	EXPRESSION TAG	UNP Q5SJ79
A	-7	LEU	-	EXPRESSION TAG	UNP Q5SJ79
A	-6	LEU	-	EXPRESSION TAG	UNP Q5SJ79
A	-5	MET	-	EXPRESSION TAG	UNP Q5SJ79
A	-4	HIS	-	EXPRESSION TAG	UNP Q5SJ79
A	-3	HIS	-	EXPRESSION TAG	UNP Q5SJ79
A	-2	HIS	-	EXPRESSION TAG	UNP Q5SJ79
A	-1	HIS	-	EXPRESSION TAG	UNP Q5SJ79
A	0	HIS	-	EXPRESSION TAG	UNP Q5SJ79
A	1	HIS	-	EXPRESSION TAG	UNP Q5SJ79
A	258	ARG	LYS	ENGINEERED	UNP Q5SJ79

- Molecule 2 is a protein called Cytochrome c oxidase subunit 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	166	Total	C	N	O	S	0	0	0
			1298	844	217	233	4			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	4	GLN	GLU	ENGINEERED	UNP Q5SJ80


- Molecule 3 is a protein called Cytochrome c oxidase polypeptide 2A.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	33	Total	C	N	O	0	0	0
			259	179	39	41			

- Molecule 4 is COPPER (I) ION (three-letter code: CU1) (formula: Cu).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Cu	0	0
			1	1		

- Molecule 5 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 6 is HEME-AS (three-letter code: HAS) (formula: $C_{54}H_{64}FeN_4O_6$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	A	1	Total	C	Fe	N	O	0	0
			65	54	1	4	6		

- Molecule 7 is B-NONYLGLUCOSIDE (three-letter code: BNG) (formula: C₁₅H₃₀O₆).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
7	A	1	21	15	6	0	0
7	A	1	21	15	6	0	0
7	A	1	21	15	6	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	C	1	Total	C	O	0	0
			21	15	6		


- Molecule 8 is DINUCLEAR COPPER ION (three-letter code: CUA) (formula: Cu₂).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	B	1	Total	Cu	0	0
			2	2		

- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	63	Total	O	0	0
			63	63		
9	B	19	Total	O	0	0
			19	19		
9	C	1	Total	O	0	0
			1	1		



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	115.63 Å 115.63 Å 149.17 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.80 19.99 – 2.55	Depositor EDS
% Data completeness (in resolution range)	99.8 (20.00-2.80) 99.5 (19.99-2.55)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.40 (at 2.56 Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.197 , 0.257 0.198 , 0.263	Depositor DCC
R_{free} test set	1274 reflections (5.27%)	DCC
Wilson B-factor (Å ²)	79.6	Xtriage
Anisotropy	0.189	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 103.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6244	wwPDB-VP
Average B, all atoms (Å ²)	89.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.08% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, HAS, BNG, CUA, CU1

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.87	1/4566 (0.0%)	0.94	7/6266 (0.1%)
2	B	0.91	0/1335	0.92	1/1822 (0.1%)
3	C	1.04	0/265	0.87	0/359
All	All	0.89	1/6166 (0.0%)	0.93	8/8447 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
2	B	0	1
All	All	0	3

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	337	ARG	CG-CD	5.78	1.66	1.51

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	435	MET	CG-SD-CE	8.98	114.58	100.20
1	A	326	LEU	CA-CB-CG	6.54	130.34	115.30
2	B	146	ARG	CB-CG-CD	-5.74	96.67	111.60
1	A	432	MET	CG-SD-CE	5.31	108.70	100.20
1	A	450	ARG	NE-CZ-NH1	-5.11	117.74	120.30

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	503	GLU	Peptide
1	A	8	ILE	Peptide
2	B	87	ALA	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4409	0	4516	204	1
2	B	1298	0	1282	43	0
3	C	259	0	279	5	0
4	A	1	0	0	0	0
5	A	43	0	30	6	0
6	A	65	0	61	8	0
7	A	63	0	90	2	0
7	C	21	0	30	0	0
8	B	2	0	0	0	0
9	A	63	0	0	47	1
9	B	19	0	0	7	0
9	C	1	0	0	0	0
All	All	6244	0	6288	243	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 20.

The worst 5 of 243 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:233:HIS:NE2	1:A:237:TYR:CE2	1.73	1.30
1:A:233:HIS:NE2	1:A:237:TYR:HE2	0.80	1.29
1:A:185:MET:CE	9:A:624:HOH:O	1.84	1.26
1:A:281:PHE:HD2	9:A:572:HOH:O	1.18	1.23
1:A:185:MET:HE3	9:A:624:HOH:O	1.38	1.23

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:330:ARG:CD	9:A:590:HOH:O[4_444]	2.19	0.01

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	555/618 (90%)	464 (84%)	71 (13%)	20 (4%)	4	13
2	B	164/166 (99%)	146 (89%)	14 (8%)	4 (2%)	7	23
3	C	31/33 (94%)	29 (94%)	2 (6%)	0	100	100
All	All	750/817 (92%)	639 (85%)	87 (12%)	24 (3%)	5	16

5 of 24 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	7	GLU
1	A	57	ARG
1	A	58	LEU
1	A	330	ARG
1	A	501	LEU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	453/503 (90%)	407 (90%)	46 (10%)	8	25
2	B	136/136 (100%)	127 (93%)	9 (7%)	19	49

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	C	26/26 (100%)	24 (92%)	2 (8%)	15	39
All	All	615/665 (92%)	558 (91%)	57 (9%)	10	30

5 of 57 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	339	LEU
1	A	401	LEU
2	B	107	ILE
1	A	354	LEU
1	A	370	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 14 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	254	GLN
1	A	284	GLN
1	A	554	ASN
1	A	127	ASN
1	A	446	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 1 is monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	HEM	A	800	1	28,50,50	2.94	14 (50%)	17,82,82	2.39	5 (29%)
6	HAS	A	801	1	55,72,72	3.98	20 (36%)	48,109,109	2.87	22 (45%)
7	BNG	A	805	-	21,21,21	0.82	1 (4%)	26,26,26	1.05	1 (3%)
7	BNG	A	806	-	21,21,21	1.03	1 (4%)	26,26,26	1.28	4 (15%)
7	BNG	A	807	-	21,21,21	0.67	1 (4%)	26,26,26	1.26	4 (15%)
8	CUA	B	802	2	0,1,1	0.00	-	0,0,0	0.00	-
7	BNG	C	804	-	21,21,21	0.82	1 (4%)	26,26,26	1.46	7 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	HEM	A	800	1	-	0/6/54/54	0/0/8/8
6	HAS	A	801	1	-	0/35/122/122	0/0/8/8
7	BNG	A	805	-	-	0/12/32/32	0/1/1/1
7	BNG	A	806	-	-	0/12/32/32	0/1/1/1
7	BNG	A	807	-	-	0/12/32/32	0/1/1/1
8	CUA	B	802	2	-	0/0/0/0	0/0/0/0
7	BNG	C	804	-	-	0/12/32/32	0/1/1/1

The worst 5 of 38 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	801	HAS	C1B-NB	-10.78	1.35	1.49
6	A	801	HAS	C1D-ND	-10.53	1.35	1.49
6	A	801	HAS	C4B-NB	-9.22	1.37	1.49
6	A	801	HAS	C4D-ND	-7.78	1.39	1.49
6	A	801	HAS	C1D-C2D	-7.43	1.39	1.51

The worst 5 of 43 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	800	HEM	CMA-C3A-C4A	-6.74	118.10	128.46

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	801	HAS	CAA-CBA-CGA	-6.17	102.12	112.66
6	A	801	HAS	CHD-C4A-C3A	-5.80	119.84	129.53
6	A	801	HAS	CAD-CBD-CGD	-5.56	103.16	112.66
6	A	801	HAS	CHD-C4C-C3C	-5.24	122.74	129.61

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 16 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	800	HEM	6	0
6	A	801	HAS	8	0
7	A	805	BNG	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	557/618 (90%)	0.13	46 (8%)	12 6	63, 85, 124, 190	0
2	B	166/166 (100%)	0.02	11 (6%)	19 11	66, 85, 118, 160	0
3	C	33/33 (100%)	-0.45	2 (6%)	22 14	67, 77, 125, 144	0
All	All	756/817 (92%)	0.08	59 (7%)	14 7	63, 85, 126, 190	0

The worst 5 of 59 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	8	ILE	10.1
1	A	495	ARG	8.8
1	A	494	SER	8.0
1	A	6	SER	7.9
2	B	69	GLN	7.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
7	BNG	A	807	21/21	0.49	0.37	3.10	120,150,158,159	0
7	BNG	C	804	21/21	0.65	0.39	2.35	89,134,146,146	0
6	HAS	A	801	65/65	0.97	0.12	-0.84	58,66,77,83	0
5	HEM	A	800	43/43	0.98	0.10	-1.20	56,62,74,79	0
8	CUA	B	802	2/2	0.98	0.05	-2.35	80,80,80,80	0
7	BNG	A	805	21/21	0.64	0.35	-	146,158,162,163	0
7	BNG	A	806	21/21	0.61	0.40	-	134,141,143,144	0
4	CU1	A	803	1/1	0.99	0.08	-	80,80,80,80	0

6.5 Other polymers ⓘ

There are no such residues in this entry.