



# wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 5, 2017 – 06:16 AM EST

PDB ID : 4EJH  
Title : Human Cytochrome P450 2A13 in complex with 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK)  
Authors : DeVore, N.M.; Scott, E.E.  
Deposited on : unknown  
Resolution : 2.35 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20030345  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20030345


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	1522 (2.38-2.34)
Clashscore	112137	1626 (2.38-2.34)
Ramachandran outliers	110173	1605 (2.38-2.34)
Sidechain outliers	110143	1606 (2.38-2.34)
RSRZ outliers	101464	1528 (2.38-2.34)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	476	
1	B	476	
1	C	476	
1	D	476	
1	E	476	

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	F	476	
1	G	476	
1	H	476	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	0QA	A	502	-	-	-	X
3	0QA	B	502	-	-	-	X
3	0QA	C	502	-	-	-	X
3	0QA	D	502	-	-	-	X
3	0QA	E	502	-	-	-	X
3	0QA	F	502	-	-	X	X
4	GOL	H	502	-	-	-	X

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 30804 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cytochrome P450 2A13.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	464	Total	C	N	O	S	8	2	0
			3777	2430	653	676	18			
1	B	464	Total	C	N	O	S	0	1	0
			3771	2426	653	674	18			
1	C	464	Total	C	N	O	S	0	2	0
			3776	2429	653	676	18			
1	D	464	Total	C	N	O	S	0	1	0
			3771	2426	653	674	18			
1	E	464	Total	C	N	O	S	0	1	0
			3771	2426	653	674	18			
1	F	464	Total	C	N	O	S	0	1	0
			3771	2426	653	674	18			
1	G	463	Total	C	N	O	S	0	0	0
			3737	2404	642	673	18			
1	H	459	Total	C	N	O	S	0	1	0
			3723	2396	640	669	18			

There are 96 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	23	MET	-	INITIATING METHIONINE	UNP Q16696
A	24	ALA	-	EXPRESSION TAG	UNP Q16696
A	25	LYS	-	EXPRESSION TAG	UNP Q16696
A	26	LYS	-	EXPRESSION TAG	UNP Q16696
A	27	THR	-	EXPRESSION TAG	UNP Q16696
A	28	SER	-	EXPRESSION TAG	UNP Q16696
A	29	SER	-	EXPRESSION TAG	UNP Q16696
A	30	LYS	-	EXPRESSION TAG	UNP Q16696
A	495	HIS	-	EXPRESSION TAG	UNP Q16696
A	496	HIS	-	EXPRESSION TAG	UNP Q16696
A	497	HIS	-	EXPRESSION TAG	UNP Q16696
A	498	HIS	-	EXPRESSION TAG	UNP Q16696
B	23	MET	-	INITIATING METHIONINE	UNP Q16696

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	24	ALA	-	EXPRESSION TAG	UNP Q16696
B	25	LYS	-	EXPRESSION TAG	UNP Q16696
B	26	LYS	-	EXPRESSION TAG	UNP Q16696
B	27	THR	-	EXPRESSION TAG	UNP Q16696
B	28	SER	-	EXPRESSION TAG	UNP Q16696
B	29	SER	-	EXPRESSION TAG	UNP Q16696
B	30	LYS	-	EXPRESSION TAG	UNP Q16696
B	495	HIS	-	EXPRESSION TAG	UNP Q16696
B	496	HIS	-	EXPRESSION TAG	UNP Q16696
B	497	HIS	-	EXPRESSION TAG	UNP Q16696
B	498	HIS	-	EXPRESSION TAG	UNP Q16696
C	23	MET	-	INITIATING METHIONINE	UNP Q16696
C	24	ALA	-	EXPRESSION TAG	UNP Q16696
C	25	LYS	-	EXPRESSION TAG	UNP Q16696
C	26	LYS	-	EXPRESSION TAG	UNP Q16696
C	27	THR	-	EXPRESSION TAG	UNP Q16696
C	28	SER	-	EXPRESSION TAG	UNP Q16696
C	29	SER	-	EXPRESSION TAG	UNP Q16696
C	30	LYS	-	EXPRESSION TAG	UNP Q16696
C	495	HIS	-	EXPRESSION TAG	UNP Q16696
C	496	HIS	-	EXPRESSION TAG	UNP Q16696
C	497	HIS	-	EXPRESSION TAG	UNP Q16696
C	498	HIS	-	EXPRESSION TAG	UNP Q16696
D	23	MET	-	INITIATING METHIONINE	UNP Q16696
D	24	ALA	-	EXPRESSION TAG	UNP Q16696
D	25	LYS	-	EXPRESSION TAG	UNP Q16696
D	26	LYS	-	EXPRESSION TAG	UNP Q16696
D	27	THR	-	EXPRESSION TAG	UNP Q16696
D	28	SER	-	EXPRESSION TAG	UNP Q16696
D	29	SER	-	EXPRESSION TAG	UNP Q16696
D	30	LYS	-	EXPRESSION TAG	UNP Q16696
D	495	HIS	-	EXPRESSION TAG	UNP Q16696
D	496	HIS	-	EXPRESSION TAG	UNP Q16696
D	497	HIS	-	EXPRESSION TAG	UNP Q16696
D	498	HIS	-	EXPRESSION TAG	UNP Q16696
E	23	MET	-	INITIATING METHIONINE	UNP Q16696
E	24	ALA	-	EXPRESSION TAG	UNP Q16696
E	25	LYS	-	EXPRESSION TAG	UNP Q16696
E	26	LYS	-	EXPRESSION TAG	UNP Q16696
E	27	THR	-	EXPRESSION TAG	UNP Q16696
E	28	SER	-	EXPRESSION TAG	UNP Q16696
E	29	SER	-	EXPRESSION TAG	UNP Q16696

*Continued on next page...*

*Continued from previous page...*


Chain	Residue	Modelled	Actual	Comment	Reference
E	30	LYS	-	EXPRESSION TAG	UNP Q16696
E	495	HIS	-	EXPRESSION TAG	UNP Q16696
E	496	HIS	-	EXPRESSION TAG	UNP Q16696
E	497	HIS	-	EXPRESSION TAG	UNP Q16696
E	498	HIS	-	EXPRESSION TAG	UNP Q16696
F	23	MET	-	INITIATING METHIONINE	UNP Q16696
F	24	ALA	-	EXPRESSION TAG	UNP Q16696
F	25	LYS	-	EXPRESSION TAG	UNP Q16696
F	26	LYS	-	EXPRESSION TAG	UNP Q16696
F	27	THR	-	EXPRESSION TAG	UNP Q16696
F	28	SER	-	EXPRESSION TAG	UNP Q16696
F	29	SER	-	EXPRESSION TAG	UNP Q16696
F	30	LYS	-	EXPRESSION TAG	UNP Q16696
F	495	HIS	-	EXPRESSION TAG	UNP Q16696
F	496	HIS	-	EXPRESSION TAG	UNP Q16696
F	497	HIS	-	EXPRESSION TAG	UNP Q16696
F	498	HIS	-	EXPRESSION TAG	UNP Q16696
G	23	MET	-	INITIATING METHIONINE	UNP Q16696
G	24	ALA	-	EXPRESSION TAG	UNP Q16696
G	25	LYS	-	EXPRESSION TAG	UNP Q16696
G	26	LYS	-	EXPRESSION TAG	UNP Q16696
G	27	THR	-	EXPRESSION TAG	UNP Q16696
G	28	SER	-	EXPRESSION TAG	UNP Q16696
G	29	SER	-	EXPRESSION TAG	UNP Q16696
G	30	LYS	-	EXPRESSION TAG	UNP Q16696
G	495	HIS	-	EXPRESSION TAG	UNP Q16696
G	496	HIS	-	EXPRESSION TAG	UNP Q16696
G	497	HIS	-	EXPRESSION TAG	UNP Q16696
G	498	HIS	-	EXPRESSION TAG	UNP Q16696
H	23	MET	-	INITIATING METHIONINE	UNP Q16696
H	24	ALA	-	EXPRESSION TAG	UNP Q16696
H	25	LYS	-	EXPRESSION TAG	UNP Q16696
H	26	LYS	-	EXPRESSION TAG	UNP Q16696
H	27	THR	-	EXPRESSION TAG	UNP Q16696
H	28	SER	-	EXPRESSION TAG	UNP Q16696
H	29	SER	-	EXPRESSION TAG	UNP Q16696
H	30	LYS	-	EXPRESSION TAG	UNP Q16696
H	495	HIS	-	EXPRESSION TAG	UNP Q16696
H	496	HIS	-	EXPRESSION TAG	UNP Q16696
H	497	HIS	-	EXPRESSION TAG	UNP Q16696
H	498	HIS	-	EXPRESSION TAG	UNP Q16696

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (for-

# HEM

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
2	B	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
2	C	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
2	D	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
2	E	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
2	F	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
2	G	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
2	H	1	Total 43	C 34	Fe 1	N 4	O 4	0	0


- Molecule 3 is 4-[methyl(nitroso)amino]-1-(pyridin-3-yl)butan-1-one (three-letter code: 0QA) (formula: C<sub>10</sub>H<sub>13</sub>N<sub>3</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			15	10	3	2		
3	B	1	Total	C	N	O	0	0
			15	10	3	2		
3	C	1	Total	C	N	O	0	0
			15	10	3	2		
3	D	1	Total	C	N	O	0	0
			15	10	3	2		
3	E	1	Total	C	N	O	0	0
			15	10	3	2		
3	F	1	Total	C	N	O	0	0
			15	10	3	2		

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	G	1	Total	C	O	0	0
			6	3	3		
4	H	1	Total	C	O	0	0
			6	3	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	56	Total	O	0	0
			56	56		
5	B	39	Total	O	0	0
			39	39		
5	C	71	Total	O	0	0
			71	71		
5	D	49	Total	O	0	0
			49	49		
5	E	13	Total	O	0	0
			13	13		
5	F	11	Total	O	0	0
			11	11		
5	G	12	Total	O	0	0
			12	12		
5	H	10	Total	O	0	0
			10	10		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: Cytochrome P450 2A13



#### • Molecule 1: Cytochrome P450 2A13




#### • Molecule 1: Cytochrome P450 2A13




Chain D:  81% 15% . .



• Molecule 1: Cytochrome P450 2A13



• Molecule 1: Cytochrome P450 2A13



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	71.68Å 119.27Å 153.66Å 100.59° 101.86° 93.56°	Depositor
Resolution (Å)	69.77 – 2.35 69.77 – 2.35	Depositor EDS
% Data completeness (in resolution range)	97.5 (69.77-2.35) 91.0 (69.77-2.35)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	0.10	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.17 (at 2.34Å)	Xtriage
Refinement program	REFMAC 6.1.13	Depositor
R, $R_{free}$	0.214 , 0.273 0.214 , 0.271	Depositor DCC
$R_{free}$ test set	9958 reflections (5.30%)	DCC
Wilson B-factor (Å <sup>2</sup> )	42.3	Xtriage
Anisotropy	0.027	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 46.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	30804	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	47.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.85% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, 0QA, HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	1.01	2/3880 (0.1%)	0.90	1/5224 (0.0%)
1	B	0.99	3/3871 (0.1%)	0.88	4/5212 (0.1%)
1	C	1.04	4/3879 (0.1%)	0.89	2/5223 (0.0%)
1	D	0.99	1/3871 (0.0%)	0.87	1/5212 (0.0%)
1	E	0.85	0/3871	0.79	0/5212
1	F	0.80	0/3871	0.76	1/5212 (0.0%)
1	G	0.82	0/3834	0.81	0/5170
1	H	0.77	0/3822	0.75	1/5149 (0.0%)
All	All	0.92	10/30899 (0.0%)	0.83	10/41614 (0.0%)

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	473	VAL	CB-CG2	5.94	1.65	1.52
1	C	383	PHE	CE2-CZ	5.37	1.47	1.37
1	B	63	GLU	CG-CD	5.28	1.59	1.51
1	B	322	GLU	CG-CD	5.24	1.59	1.51
1	A	133	ALA	CA-CB	5.20	1.63	1.52

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	123	ARG	NE-CZ-NH1	-5.97	117.31	120.30
1	C	143	ARG	NE-CZ-NH2	-5.90	117.35	120.30
1	H	288	LEU	CA-CB-CG	5.86	128.78	115.30
1	C	143	ARG	NE-CZ-NH1	5.85	123.22	120.30
1	F	316	LEU	CA-CB-CG	5.79	128.63	115.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3777	0	3747	50	0
1	B	3771	0	3741	47	0
1	C	3776	0	3745	53	0
1	D	3771	0	3741	56	0
1	E	3771	0	3741	55	0
1	F	3771	0	3741	78	0
1	G	3737	0	3670	85	0
1	H	3723	0	3668	81	0
2	A	43	0	30	2	0
2	B	43	0	30	1	0
2	C	43	0	30	3	0
2	D	43	0	30	2	0
2	E	43	0	30	3	0
2	F	43	0	30	9	0
2	G	43	0	30	6	0
2	H	43	0	30	6	0
3	A	15	0	13	4	0
3	B	15	0	13	5	0
3	C	15	0	13	4	0
3	D	15	0	13	5	0
3	E	15	0	13	3	0
3	F	15	0	13	9	0
4	G	6	0	8	2	0
4	H	6	0	8	0	0
5	A	56	0	0	0	0
5	B	39	0	0	3	0
5	C	71	0	0	4	0
5	D	49	0	0	3	0
5	E	13	0	0	0	0
5	F	11	0	0	0	0
5	G	12	0	0	1	0
5	H	10	0	0	0	0
All	All	30804	0	30128	524	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 524 close contacts within the same asymmetric unit are listed below, sorted by

their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:32:LYS:HD2	1:C:33:LEU:H	1.09	1.07
1:D:381:ARG:O	1:D:382:ASP:HB2	1.55	1.03
1:D:143:ARG:HH12	1:D:147:GLU:HB2	1.22	1.03
3:B:502:0QA:O1	3:B:502:0QA:H5	1.58	0.99
1:A:77:ARG:HG2	1:A:77:ARG:HH11	1.26	0.98

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	464/476 (98%)	439 (95%)	24 (5%)	1 (0%)	51	61
1	B	463/476 (97%)	445 (96%)	18 (4%)	0	100	100
1	C	464/476 (98%)	443 (96%)	20 (4%)	1 (0%)	51	61
1	D	463/476 (97%)	444 (96%)	18 (4%)	1 (0%)	51	61
1	E	463/476 (97%)	434 (94%)	28 (6%)	1 (0%)	51	61
1	F	463/476 (97%)	413 (89%)	44 (10%)	6 (1%)	14	12
1	G	461/476 (97%)	417 (90%)	36 (8%)	8 (2%)	11	8
1	H	456/476 (96%)	396 (87%)	53 (12%)	7 (2%)	12	10
All	All	3697/3808 (97%)	3431 (93%)	241 (6%)	25 (1%)	25	27

5 of 25 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	346	ARG
1	G	364	ASP
1	G	468	PRO
1	A	369	GLY

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type
1	E	162	GLY

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	410/419 (98%)	391 (95%)	19 (5%)	31	39
1	B	409/419 (98%)	392 (96%)	17 (4%)	34	43
1	C	410/419 (98%)	394 (96%)	16 (4%)	37	47
1	D	409/419 (98%)	388 (95%)	21 (5%)	28	33
1	E	409/419 (98%)	380 (93%)	29 (7%)	17	18
1	F	409/419 (98%)	383 (94%)	26 (6%)	20	22
1	G	403/419 (96%)	375 (93%)	28 (7%)	18	19
1	H	403/419 (96%)	354 (88%)	49 (12%)	6	5
All	All	3262/3352 (97%)	3057 (94%)	205 (6%)	21	23

5 of 205 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	382	ASP
1	F	283	ASN
1	H	376	LYS
1	E	424	LYS
1	F	135	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 8 such sidechains are listed below:

Mol	Chain	Res	Type
1	F	283	ASN
1	G	255	ASN
1	G	104	GLN
1	E	210	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	F	422	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

16 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	HEM	A	501	1,3	28,50,50	2.32	8 (28%)	17,82,82	1.80	5 (29%)
3	0QA	A	502	2	13,15,15	0.44	0	14,18,18	2.18	5 (35%)
2	HEM	B	501	1	28,50,50	2.52	11 (39%)	17,82,82	2.28	5 (29%)
3	0QA	B	502	-	13,15,15	0.62	0	14,18,18	1.97	5 (35%)
2	HEM	C	501	1,3	28,50,50	2.44	8 (28%)	17,82,82	1.89	7 (41%)
3	0QA	C	502	2	13,15,15	0.62	0	14,18,18	1.51	4 (28%)
2	HEM	D	501	1	28,50,50	2.33	8 (28%)	17,82,82	1.13	1 (5%)
3	0QA	D	502	-	13,15,15	0.63	0	14,18,18	1.51	4 (28%)
2	HEM	E	501	1,3	28,50,50	2.24	8 (28%)	17,82,82	1.42	2 (11%)
3	0QA	E	502	2	13,15,15	0.70	0	14,18,18	1.52	3 (21%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	HEM	F	501	1,3	28,50,50	2.21	8 (28%)	17,82,82	1.80	4 (23%)
3	0QA	F	502	2	13,15,15	0.53	0	14,18,18	1.41	2 (14%)
2	HEM	G	501	1	28,50,50	2.42	9 (32%)	17,82,82	1.55	2 (11%)
4	GOL	G	502	-	5,5,5	0.62	0	5,5,5	0.47	0
2	HEM	H	501	1	28,50,50	2.17	6 (21%)	17,82,82	2.36	5 (29%)
4	GOL	H	502	-	5,5,5	1.34	1 (20%)	5,5,5	2.38	2 (40%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	501	1,3	-	0/6/54/54	0/0/8/8
3	0QA	A	502	2	-	0/12/12/12	0/1/1/1
2	HEM	B	501	1	-	0/6/54/54	0/0/8/8
3	0QA	B	502	-	-	0/12/12/12	0/1/1/1
2	HEM	C	501	1,3	-	0/6/54/54	0/0/8/8
3	0QA	C	502	2	-	0/12/12/12	0/1/1/1
2	HEM	D	501	1	-	0/6/54/54	0/0/8/8
3	0QA	D	502	-	-	0/12/12/12	0/1/1/1
2	HEM	E	501	1,3	-	0/6/54/54	0/0/8/8
3	0QA	E	502	2	-	1/12/12/12	0/1/1/1
2	HEM	F	501	1,3	-	0/6/54/54	0/0/8/8
3	0QA	F	502	2	-	0/12/12/12	0/1/1/1
2	HEM	G	501	1	-	0/6/54/54	0/0/8/8
4	GOL	G	502	-	-	0/4/4/4	0/0/0/0
2	HEM	H	501	1	-	0/6/54/54	0/0/8/8
4	GOL	H	502	-	-	0/4/4/4	0/0/0/0

The worst 5 of 67 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	501	HEM	C3C-C2C	-6.30	1.32	1.40
2	D	501	HEM	C3C-C2C	-5.93	1.32	1.40
2	B	501	HEM	C3B-C2B	-5.84	1.32	1.40
2	A	501	HEM	C3C-C2C	-5.82	1.32	1.40
2	C	501	HEM	C3B-C2B	-5.81	1.32	1.40

The worst 5 of 56 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	501	HEM	C1D-C2D-C3D	-5.39	103.25	107.00
2	H	501	HEM	CAA-CBA-CGA	-5.36	103.50	112.66
2	B	501	HEM	CBD-CAD-C3D	-4.98	102.97	112.47
3	A	502	0QA	C9-C8-C7	-4.76	100.96	112.06
2	H	501	HEM	C1D-C2D-C3D	-4.65	103.76	107.00

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	E	502	0QA	O2-N3-N2-C10

There are no ring outliers.

15 monomers are involved in 57 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	501	HEM	2	0
3	A	502	0QA	4	0
2	B	501	HEM	1	0
3	B	502	0QA	5	0
2	C	501	HEM	3	0
3	C	502	0QA	4	0
2	D	501	HEM	2	0
3	D	502	0QA	5	0
2	E	501	HEM	3	0
3	E	502	0QA	3	0
2	F	501	HEM	9	0
3	F	502	0QA	9	0
2	G	501	HEM	6	0
4	G	502	GOL	2	0
2	H	501	HEM	6	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	464/476 (97%)	0.22	0 100 100	20, 35, 53, 63	0
1	B	464/476 (97%)	0.22	0 100 100	19, 35, 54, 62	0
1	C	464/476 (97%)	0.17	1 (0%) 94 97	19, 33, 48, 58	0
1	D	464/476 (97%)	0.20	1 (0%) 94 97	17, 35, 55, 63	0
1	E	464/476 (97%)	0.26	12 (2%) 56 66	27, 48, 70, 78	0
1	F	464/476 (97%)	0.63	30 (6%) 20 27	27, 60, 85, 91	0
1	G	463/476 (97%)	0.66	34 (7%) 16 23	29, 59, 78, 88	0
1	H	459/476 (96%)	1.01	89 (19%) 1 2	28, 71, 92, 96	0
All	All	3706/3808 (97%)	0.42	167 (4%) 34 46	17, 44, 81, 96	0

The worst 5 of 167 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	141	GLY	6.4
1	H	491	PHE	6.0
1	F	491	PHE	5.2
1	H	453	PHE	4.9
1	H	468	PRO	4.9

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
4	GOL	H	502	6/6	0.94	0.27	6.75	33,36,42,50	0
3	0QA	F	502	15/15	0.90	0.36	5.32	84,85,86,87	0
3	0QA	C	502	15/15	0.91	0.22	4.88	48,55,58,67	0
3	0QA	E	502	15/15	0.90	0.29	4.57	71,74,76,78	0
3	0QA	A	502	15/15	0.94	0.24	4.08	57,62,71,72	0
3	0QA	B	502	15/15	0.91	0.21	2.20	42,51,59,59	0
3	0QA	D	502	15/15	0.94	0.21	2.17	48,55,58,67	0
2	HEM	A	501	43/43	0.98	0.17	0.74	21,27,31,36	0
2	HEM	B	501	43/43	0.98	0.17	0.44	18,25,29,32	0
2	HEM	D	501	43/43	0.98	0.16	0.24	18,26,29,30	0
2	HEM	G	501	43/43	0.98	0.15	-0.09	33,38,43,47	0
2	HEM	E	501	43/43	0.98	0.15	-0.10	30,35,37,39	0
2	HEM	C	501	43/43	0.98	0.15	-0.18	12,22,27,28	0
2	HEM	F	501	43/43	0.97	0.16	-0.20	37,49,54,59	0
4	GOL	G	502	6/6	0.92	0.16	-0.31	54,56,58,58	0
2	HEM	H	501	43/43	0.96	0.16	-0.43	47,52,60,63	0

## 6.5 Other polymers

There are no such residues in this entry.