



# wwPDB NMR Structure Validation Summary Report ⓘ

Feb 12, 2017 – 05:58 pm GMT

PDB ID : 1EKZ  
Title : NMR STRUCTURE OF THE COMPLEX BETWEEN THE THIRD DSRBD FROM DROSOPHILA STAUFEN AND A RNA HAIRPIN  
Authors : Ramos, A.; Grunert, S.; Bycroft, M.; St Johnston, D.; Varani, G.  
Deposited on : 2000-03-11

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)  
NmrClust : Kelley et al. (1996)  
MolProbity : 4.02b-467  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
RCI : v\_1n\_11\_5\_13\_A (Berjanski et al., 2005)  
PANAV : Wang et al. (2010)  
ShiftChecker : trunk28760  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*SOLUTION NMR*

The overall completeness of chemical shifts assignment is 73%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	125131	11601
Ramachandran outliers	121729	10391
Sidechain outliers	121581	10367
RNA backbone	3398	623

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	B	30	 13% 50% 30% 7%
2	A	76	 13% 53% 30%

## 2 Ensemble composition and analysis i

This entry contains 36 models. Model 27 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:9-A:30, A:40-A:55, A:62-A:76 (53)	0.60	27

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters and 13 single-model clusters were found.

Cluster number	Models
1	2, 3, 4, 7, 8, 10, 11, 13, 17, 20, 23, 24, 25, 27, 28, 30, 32, 33
2	1, 5, 22
3	6, 21
Single-model clusters	9; 12; 14; 15; 16; 18; 19; 26; 29; 31; 34; 35; 36

### 3 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2173 atoms, of which 949 are hydrogens and 0 are deuteriums.

- Molecule 1 is a RNA chain called STAUFEN DOUBLE-STRANDED RNA BINDING DOMAIN.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	P	
1	B	30	961	284	326	112	210	29	0

- Molecule 2 is a protein called MATERNAL EFFECT PROTEIN (STAUFEN).

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
2	A	76	1212	368	623	107	109	5	0

## 4 Residue-property plots

### 4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: STAUFEN DOUBLE-STRANDED RNA BINDING DOMAIN

Chain B: 



- Molecule 2: MATERNAL EFFECT PROTEIN (STAUFEN)

Chain A: 



### 4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 27. Colouring as in section 4.1 above.

- Molecule 1: STAUFEN DOUBLE-STRANDED RNA BINDING DOMAIN

Chain B: 



- Molecule 2: MATERNAL EFFECT PROTEIN (STAUFEN)

Chain A: 



M69	L76
L70	
V71	
E72	
L73	
	L76

## 5 Refinement protocol and experimental data overview

Of the 50 calculated structures, 36 were deposited, based on the following criterion: *structures with the least restraint violations, structures with the lowest energy.*

The authors did not provide any information on software used for structure solution, optimization or refinement.

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	BMRB entry 4894
Number of chemical shift lists	2
Total number of shifts	1223
Number of shifts mapped to atoms	1121
Number of unparsed shifts	0
Number of shifts with mapping errors	102
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	73%

No validations of the models with respect to experimental NMR restraints is performed at this time.

## 6 Model quality i

### 6.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	B	1.10±0.01	0±0/708 (0.0±0.0%)	1.89±0.01	38±0/1102 (3.5±0.0%)
2	A	1.13±0.01	0±0/420 (0.0±0.0%)	0.84±0.01	0±0/558 (0.0±0.0%)
All	All	1.11	0/40608 (0.0%)	1.62	1369/59760 (2.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	B	0.0±0.0	0.1±0.3
2	A	0.0±0.0	3.0±0.0
All	All	0	110

There are no bond-length outliers.

5 of 42 unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	B	17	G	N7-C8-N9	9.47	117.83	113.10	5	36
1	B	1	G	N7-C8-N9	9.45	117.82	113.10	36	36
1	B	9	G	N7-C8-N9	9.37	117.79	113.10	32	36
1	B	18	G	N7-C8-N9	9.31	117.75	113.10	7	36
1	B	20	G	N7-C8-N9	9.29	117.75	113.10	32	36

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
2	A	64	ARG	Sidechain	36

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Group	Models (Total)
2	A	20	ARG	Sidechain	36
2	A	30	ARG	Sidechain	36
1	B	9	G	Sidechain	1
1	B	22	C	Sidechain	1

## 6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	B	635	326	326	10±6
2	A	416	453	453	24±11
All	All	37836	28044	28044	1193

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

5 of 650 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
2:A:41:ILE:HG23	2:A:65:ALA:HB2	1.01	1.26	31	1
2:A:13:VAL:HG22	2:A:70:LEU:HD21	0.96	1.36	25	20
2:A:50:VAL:HG11	2:A:69:MET:CE	0.95	1.92	36	1
2:A:50:VAL:HG11	2:A:69:MET:HE1	0.94	1.35	36	1
2:A:44:CYS:SG	2:A:65:ALA:HB1	0.93	2.02	2	5

## 6.3 Torsion angles [i](#)

### 6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	A	52/76 (68%)	42±3 (81±5%)	7±2 (14±4%)	3±2 (6±3%)	4	23

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	1872/2736 (68%)	1507 (81%)	261 (14%)	104 (6%)	4 23

5 of 25 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
2	A	21	ASN	14
2	A	9	PRO	11
2	A	47	GLY	11
2	A	54	GLU	7
2	A	30	ARG	6

### 6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
2	A	46/65 (71%)	35±3 (77±6%)	11±3 (23±6%)	3 29
All	All	1656/2340 (71%)	1270 (77%)	386 (23%)	3 29

5 of 41 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
2	A	23	THR	27
2	A	11	SER	26
2	A	75	LYS	20
2	A	19	LYS	20
2	A	54	GLU	18

### 6.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers	Suiteness
1	B	29/30 (97%)	13±3 (46±10%)	0±0 (0±0%)	0.13±0.03
All	All	1044/1080 (97%)	479 (46%)	0 (0%)	0.12

The overall RNA backbone suiteness is 0.13.

5 of 29 unique RNA backbone outliers are listed below:

Mol	Chain	Res	Type	Models (Total)
1	B	18	G	30
1	B	17	G	28
1	B	27	G	26
1	B	30	C	23
1	B	9	G	21

There are no RNA pucker outliers to report.

#### 6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

#### 6.6 Ligand geometry [i](#)

There are no ligands in this entry.

#### 6.7 Other polymers [i](#)

There are no such molecules in this entry.

#### 6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 7 Chemical shift validation i

The completeness of assignment taking into account all chemical shift lists is 73% for the well-defined parts and 67% for the entire structure.

### 7.1 Chemical shift list 1

File name: BMRB entry 4894

Chemical shift list name: *assigned\_chem\_shift\_list\_1*

#### 7.1.1 Bookkeeping i

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	736
Number of shifts mapped to atoms	634
Number of unparsed shifts	0
Number of shifts with mapping errors	102
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	4

The following assigned chemical shifts were not mapped to the molecules present in the coordinate file.

- Residue not found in structure. First 5 (of 102) occurrences are reported below.

Chain	Res	Type	Atom	Shift Data		
				Value	Uncertainty	Ambiguity
A	84	THR	H	8.31	-1.0	1
A	87	PHE	CA	56.0	-1.0	1
A	83	PRO	CD	50.0	-1.0	1
A	81	LEU	HB3	1.6	-1.0	1
A	81	LEU	N	122.3	-1.0	1

#### 7.1.2 Chemical shift referencing i

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction $\pm$ precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	70	$1.25 \pm 0.17$	Should be applied
$^{13}\text{C}_\beta$	34	$1.91 \pm 0.28$	Should be applied
$^{13}\text{C}'$	0	—	None (insufficient data)

*Continued on next page...*

Continued from previous page...

Nucleus	# values	Correction $\pm$ precision, ppm	Suggested action
<sup>15</sup> N	70	0.97 $\pm$ 0.39	Should be applied

### 7.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 35%, i.e. 439 atoms were assigned a chemical shift out of a possible 1247. 3 out of 10 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	<sup>1</sup> H	<sup>13</sup> C	<sup>15</sup> N
Backbone	198/263 (75%)	102/105 (97%)	47/106 (44%)	49/52 (94%)
Sidechain	225/384 (59%)	172/224 (77%)	46/142 (32%)	7/18 (39%)
Aromatic	16/32 (50%)	12/18 (67%)	4/12 (33%)	0/2 (0%)
Overall	439/1247 (35%)	286/675 (42%)	97/460 (21%)	56/112 (50%)

### 7.1.4 Statistically unusual chemical shifts [i](#)


The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

Mol	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	12	GLN	CA	4.10	67.31 – 45.91	-24.5
1	A	22	MET	CG	16.00	38.33 – 25.73	-12.7
1	A	38	LYS	HD2	3.00	2.76 – 0.46	6.0
1	A	90	ASP	CB	31.00	49.06 – 32.66	-6.0

### 7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:



## 7.2 Chemical shift list 2

File name: BMRB entry 4894

Chemical shift list name: *assigned\_chem\_shift\_list\_2*

### 7.2.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	487
Number of shifts mapped to atoms	487
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	5

### 7.2.2 Chemical shift referencing [i](#)

No chemical shift referencing corrections were calculated (not enough data).

### 7.2.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 38%, i.e. 479 atoms were assigned a chemical shift out of a possible 1249. 0 out of 10 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	<sup>1</sup> H	<sup>13</sup> C	<sup>15</sup> N
Backbone	0/263 (0%)	0/105 (0%)	0/106 (0%)	0/52 (0%)
Sidechain	0/384 (0%)	0/224 (0%)	0/142 (0%)	0/18 (0%)
Aromatic	0/32 (0%)	0/18 (0%)	0/12 (0%)	0/2 (0%)

*Continued on next page...*

*Continued from previous page...*

	<b>Total</b>	<sup>1</sup> <b>H</b>	<sup>13</sup> <b>C</b>	<sup>15</sup> <b>N</b>
Overall	479/1249 (38%)	268/675 (40%)	194/460 (42%)	17/114 (15%)

#### 7.2.4 Statistically unusual chemical shifts [i](#)

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

Mol	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
2	B	8	U	N1	163.60	152.56 – 139.26	13.3
2	B	14	U	H6	5.81	8.67 – 6.87	-10.9
2	B	14	U	H5	7.83	6.81 – 4.11	8.8
2	B	22	C	H5	7.09	6.75 – 4.15	6.3
2	B	16	C	H5''	2.78	5.47 – 2.87	-5.3

#### 7.2.5 Random Coil Index (RCI) plots [i](#)

No *random coil index* (RCI) plot could be generated from the current chemical shift list (assigned\_chem\_shift\_list\_2). RCI is only applicable to proteins.