



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 02:33 am GMT

PDB ID : 3EKZ
Title : Structural Characterization of tetrameric Mycobacterium tuberculosis fructose 1,6-bisphosphate aldolase - substrate binding and catalysis mechanism of a class IIa bacterial aldolase
Authors : Pegan, S.; Rukseere, K.; Franzblau, S.G.; Mesecar, A.D.
Deposited on : 2008-09-19
Resolution : 2.07 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.07 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	2028 (2.08-2.04)
Clashscore	112137	2143 (2.08-2.04)
Ramachandran outliers	110173	2126 (2.08-2.04)
Sidechain outliers	110143	2126 (2.08-2.04)
RSRZ outliers	101464	2035 (2.08-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	349	<div> <div style="width: 4%; background-color: red;"></div> <div style="width: 80%; background-color: green;"></div> <div style="width: 13%; background-color: yellow;"></div> <div style="width: 3%; background-color: orange;"></div> <div style="width: 2%; background-color: grey;"></div> </div> <div>4% 80% 13% ...</div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	13P	A	350[A]	-	-	X	-
2	13P	A	351[B]	-	-	X	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	G3P	A	352	X	-	X	-

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 2972 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Fructose-bisphosphate aldolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	334	2578	1637	439	492	10	0	10	0

There are 5 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	345	HIS	-	EXPRESSION TAG	UNP P67475
A	346	HIS	-	EXPRESSION TAG	UNP P67475
A	347	HIS	-	EXPRESSION TAG	UNP P67475
A	348	HIS	-	EXPRESSION TAG	UNP P67475
A	349	HIS	-	EXPRESSION TAG	UNP P67475

- Molecule 2 is 1,3-DIHYDROXYACETONEPHOSPHATE (three-letter code: 13P) (formula: $C_3H_7O_6P$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	O	P	0	1
			10	3	6	1		
2	A	1	Total	C	O	P	0	1
			10	3	6	1		

- Molecule 3 is SN-GLYCEROL-3-PHOSPHATE (three-letter code: G3P) (formula: $C_3H_9O_6P$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	O	P	0	0
			10	3	6	1		

- Molecule 4 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Na	0	0
			1	1		

- Molecule 5 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	2	Total	Zn	0	0
			2	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	361	Total 361	O 361	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Fructose-bisphosphate aldolase



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	61.22Å 120.49Å 164.33Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	25.90 – 2.07 25.90 – 2.07	Depositor EDS
% Data completeness (in resolution range)	97.4 (25.90-2.07) 85.6 (25.90-2.07)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	8.09 (at 2.06Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.158 , 0.180 0.181 , 0.201	Depositor DCC
R_{free} test set	1589 reflections (4.95%)	DCC
Wilson B-factor (Å ²)	21.7	Xtriage
Anisotropy	0.567	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 55.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	2972	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.67% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: G3P, ZN, 13P, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.72	0/2633	0.80	6/3565 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	66	LYS	N-CA-CB	-5.75	100.25	110.60
1	A	180	LEU	C-N-CA	5.69	135.93	121.70
1	A	92[A]	LEU	CA-CB-CG	5.56	128.09	115.30
1	A	92[B]	LEU	CA-CB-CG	5.56	128.09	115.30
1	A	305	VAL	CG1-CB-CG2	5.55	119.78	110.90
1	A	117	VAL	N-CA-C	5.04	124.61	111.00

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	117	VAL	Peptide
1	A	180	LEU	Peptide
1	A	65	VAL	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2578	0	2555	43	0
2	A	20	0	9	8	0
3	A	10	0	7	6	0
4	A	1	0	0	0	0
5	A	2	0	0	0	0
6	A	361	0	0	9	0
All	All	2972	0	2571	49	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:231:GLN:HE22	1:A:247:PHE:H	1.23	0.87
1:A:274:ASN:HB3	2:A:350[A]:13P:H31	1.61	0.82
2:A:351[B]:13P:H32	3:A:352:G3P:C2	2.11	0.80
1:A:92[A]:LEU:H	1:A:126:GLN:NE2	1.84	0.76
1:A:92[B]:LEU:H	1:A:126:GLN:NE2	1.84	0.76
1:A:290:HIS:HD2	1:A:319:LYS:NZ	1.85	0.74
1:A:192:GLU:OE2	1:A:237:LYS:HE3	1.91	0.70
1:A:92[B]:LEU:H	1:A:126:GLN:HE21	1.37	0.69
1:A:92[A]:LEU:H	1:A:126:GLN:HE21	1.39	0.68
1:A:115[A]:GLN:HG3	6:A:636:HOH:O	1.93	0.67
2:A:351[B]:13P:H32	3:A:352:G3P:H2	1.78	0.65
1:A:115[A]:GLN:CG	6:A:636:HOH:O	2.45	0.64
1:A:107:ARG:HD3	6:A:478:HOH:O	1.97	0.64
1:A:272:LYS:HZ3	1:A:274:ASN:HD21	1.43	0.64
1:A:231:GLN:HA	1:A:231:GLN:HE21	1.63	0.63
1:A:220[A]:VAL:HG23	6:A:495:HOH:O	1.99	0.62
2:A:351[B]:13P:H32	3:A:352:G3P:O2	2.01	0.60
1:A:180:LEU:N	1:A:181:TYR:HB2	2.16	0.60
1:A:188[A]:GLU:HG3	1:A:233:VAL:CG1	2.33	0.59
1:A:272:LYS:NZ	1:A:274:ASN:HD21	2.01	0.57
1:A:290:HIS:HE1	1:A:310:VAL:O	1.88	0.57

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:290:HIS:HD2	1:A:319:LYS:HZ1	1.54	0.56
1:A:188[A]:GLU:HG3	1:A:233:VAL:HG13	1.86	0.56
1:A:114:ALA:O	1:A:118:SER:HB3	2.07	0.54
1:A:93:HIS:HD2	1:A:127:SER:OG	1.89	0.54
1:A:192:GLU:OE2	1:A:237:LYS:CE	2.56	0.53
1:A:180:LEU:H	1:A:181:TYR:HB2	1.71	0.53
1:A:290:HIS:HD2	1:A:319:LYS:HZ2	1.57	0.53
1:A:259:LYS:HE2	6:A:463:HOH:O	2.09	0.53
3:A:352:G3P:H32	6:A:684:HOH:O	2.09	0.51
1:A:180:LEU:HB2	1:A:181:TYR:CD1	2.46	0.51
1:A:274:ASN:HB3	2:A:350[A]:13P:C3	2.36	0.49
1:A:38:ILE:HG13	1:A:50:ILE:HD11	1.95	0.48
1:A:214:VAL:HG13	1:A:277:THR:CG2	2.43	0.48
1:A:313:PRO:HA	1:A:316:TYR:CE2	2.49	0.48
2:A:351[B]:13P:H32	3:A:352:G3P:C1	2.46	0.46
1:A:58:GLU:HG3	1:A:66:LYS:HB3	1.98	0.46
1:A:80:HIS:HE1	6:A:580:HOH:O	2.00	0.45
1:A:79:THR:HG21	1:A:92[B]:LEU:HD21	1.99	0.45
1:A:162:ILE:HG12	1:A:204:LEU:HD11	1.99	0.44
1:A:214:VAL:HG13	1:A:277:THR:HG21	2.00	0.44
3:A:352:G3P:O4P	6:A:685:HOH:O	2.21	0.44
1:A:253:GLY:HA2	2:A:350[A]:13P:O1	2.17	0.44
1:A:345:HIS:HB2	6:A:490:HOH:O	2.17	0.44
1:A:313:PRO:HA	1:A:316:TYR:CZ	2.54	0.42
1:A:58:GLU:CG	1:A:66:LYS:HB3	2.49	0.42
1:A:93:HIS:CD2	1:A:127:SER:OG	2.72	0.41
1:A:274:ASN:O	2:A:350[A]:13P:H11	2.20	0.41
1:A:290:HIS:CD2	1:A:319:LYS:HZ2	2.38	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	340/349 (97%)	330 (97%)	5 (2%)	5 (2%)	12 3

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	66	LYS
1	A	118	SER
1	A	181	TYR
1	A	96	HIS
1	A	217	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	268/271 (99%)	247 (92%)	21 (8%)	15 6

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	90	VAL
1	A	92[A]	LEU
1	A	92[B]	LEU
1	A	115[A]	GLN
1	A	115[B]	GLN
1	A	141	LEU
1	A	148	LEU
1	A	185[A]	GLU
1	A	185[B]	GLU
1	A	194	LEU
1	A	198	GLU
1	A	201[A]	LYS
1	A	201[B]	LYS
1	A	210	ASN
1	A	229[A]	GLN
1	A	229[B]	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	231	GLN
1	A	251	PHE
1	A	266	LEU
1	A	275	VAL
1	A	305	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	15	GLN
1	A	19	ASN
1	A	80	HIS
1	A	93	HIS
1	A	126	GLN
1	A	219	ASN
1	A	231	GLN
1	A	274	ASN
1	A	280	GLN
1	A	290	HIS
1	A	336	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 3 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	13P	A	350[A]	4	9,9,9	0.58	0	10,12,12	0.98	0
2	13P	A	351[B]	5,4	9,9,9	3.62	3 (33%)	10,12,12	1.27	1 (10%)
3	G3P	A	352	-	9,9,9	0.54	0	11,12,12	0.69	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	13P	A	350[A]	4	-	0/7/8/8	0/0/0/0
2	13P	A	351[B]	5,4	-	0/7/8/8	0/0/0/0
3	G3P	A	352	-	1/1/2/2	0/8/8/8	0/0/0/0

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	351[B]	13P	C3-C2	-6.12	1.34	1.50
2	A	351[B]	13P	O1-C1	4.86	1.46	1.43
2	A	351[B]	13P	O2-C2	7.02	1.34	1.21

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	351[B]	13P	O3-C3-C2	2.77	121.89	112.42

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
3	A	352	G3P	C2

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	350[A]	13P	4	0
2	A	351[B]	13P	4	0
3	A	352	G3P	6	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	334/349 (95%)	-0.16	15 (4%) 34 35	13, 18, 24, 30	0

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	180	LEU	8.6
1	A	181	TYR	5.2
1	A	217	PRO	5.1
1	A	218	GLY	4.3
1	A	242	ALA	4.1
1	A	179	LYS	4.1
1	A	219	ASN	3.1
1	A	240	LEU	2.9
1	A	241	PRO	2.6
1	A	119	LYS	2.5
1	A	220[A]	VAL	2.5
1	A	216	LYS	2.3
1	A	167	GLY	2.2
1	A	164	VAL	2.2
1	A	221	LYS	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	G3P	A	352	10/10	0.96	0.12	0.62	30,36,47,48	0
2	13P	A	350[A]	10/10	0.98	0.11	-0.28	12,15,17,19	10
2	13P	A	351[B]	10/10	0.98	0.09	-0.84	16,18,21,21	10
4	NA	A	353	1/1	0.99	0.04	-1.86	17,17,17,17	0
5	ZN	A	354	1/1	0.98	0.07	-3.08	35,35,35,35	0
5	ZN	A	355	1/1	0.99	0.05	-	24,24,24,24	0

6.5 Other polymers [i](#)

There are no such residues in this entry.