



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 12:41 am GMT

PDB ID : 1EN7
Title : ENDONUCLEASE VII (ENDOVII) FROM PHAGE T4
Authors : Raaijmakers, H.; Vix, O.; Toro, I.; Suck, D.
Deposited on : 1999-02-07
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	112137	3674 (2.40-2.40)
Ramachandran outliers	110173	3616 (2.40-2.40)
Sidechain outliers	110143	3617 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	157	
1	B	157	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 2622 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RECOMBINATION ENDONUCLEASE VII.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	157	Total	C	N	O	S	0	0	0
			1272	796	227	240	9			
1	B	157	Total	C	N	O	S	0	0	0
			1272	796	227	240	9			

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Zn	0	0
			1	1		
2	A	1	Total	Zn	0	0
			1	1		

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	2	Total	Ca	0	0
			2	2		
3	A	1	Total	Ca	0	0
			1	1		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	30	Total	O	0	0
			30	30		
4	B	43	Total	O	0	0
			43	43		

3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: RECOMBINATION ENDONUCLEASE VII



• Molecule 1: RECOMBINATION ENDONUCLEASE VII



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	144.99Å 39.45Å 75.75Å 90.00° 106.20° 90.00°	Depositor
Resolution (Å)	10.00 – 2.40	Depositor
% Data completeness (in resolution range)	99.5 (10.00-2.40)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	0.06	Depositor
Refinement program	REFMAC	Depositor
R, R_{free}	0.239 , 0.308	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	2622	wwPDB-VP
Average B, all atoms (Å ²)	46.0	wwPDB-VP

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.22	4/1293 (0.3%)	1.63	12/1725 (0.7%)
1	B	1.35	3/1293 (0.2%)	1.73	18/1725 (1.0%)
All	All	1.28	7/2586 (0.3%)	1.68	30/3450 (0.9%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	6
1	B	0	11
All	All	0	17

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	147	SER	CB-OG	-7.07	1.33	1.42
1	B	153	ARG	CD-NE	-7.01	1.34	1.46
1	A	131	ARG	CD-NE	-6.81	1.34	1.46
1	B	14	LYS	CB-CG	-5.86	1.36	1.52
1	A	131	ARG	NE-CZ	-5.45	1.25	1.33
1	A	86	GLU	CD-OE2	5.23	1.31	1.25
1	A	10	GLU	CD-OE2	-5.08	1.20	1.25

All (30) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	131	ARG	CD-NE-CZ	25.66	159.53	123.60
1	B	153	ARG	CD-NE-CZ	24.43	157.81	123.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	14	LYS	CA-CB-CG	15.61	147.74	113.40
1	B	54	ARG	NE-CZ-NH2	12.20	126.40	120.30
1	A	131	ARG	NE-CZ-NH2	10.33	125.47	120.30
1	A	33	ASP	CB-CG-OD2	-10.20	109.12	118.30
1	B	54	ARG	NE-CZ-NH1	-8.57	116.02	120.30
1	A	1	MET	CA-CB-CG	8.41	127.60	113.30
1	A	10	GLU	CA-CB-CG	8.37	131.82	113.40
1	B	1	MET	CA-CB-CG	8.25	127.33	113.30
1	A	44	GLU	OE1-CD-OE2	8.05	132.96	123.30
1	B	14	LYS	CB-CG-CD	7.94	132.24	111.60
1	B	28	ARG	NE-CZ-NH1	7.82	124.21	120.30
1	B	28	ARG	CD-NE-CZ	7.37	133.91	123.60
1	A	54	ARG	NE-CZ-NH2	6.69	123.65	120.30
1	B	153	ARG	NE-CZ-NH2	6.68	123.64	120.30
1	A	157	LYS	CA-CB-CG	6.56	127.84	113.40
1	B	141	LYS	CA-CB-CG	6.13	126.88	113.40
1	B	100	THR	CA-CB-CG2	6.11	120.95	112.40
1	B	42	ASP	CB-CG-OD2	-5.85	113.03	118.30
1	A	96	LYS	N-CA-CB	-5.73	100.29	110.60
1	B	153	ARG	CA-CB-CG	5.49	125.47	113.40
1	B	42	ASP	CB-CA-C	5.44	121.27	110.40
1	A	14	LYS	CA-CB-CG	5.43	125.34	113.40
1	B	14	LYS	N-CA-CB	5.38	120.28	110.60
1	A	131	ARG	NE-CZ-NH1	-5.26	117.67	120.30
1	B	94	TYR	CB-CG-CD2	5.10	124.06	121.00
1	B	67	GLN	CB-CG-CD	5.08	124.80	111.60
1	A	17	ASP	CB-CG-OD2	-5.06	113.75	118.30
1	B	114	LYS	CA-CB-CG	5.02	124.44	113.40

There are no chirality outliers.

All (17) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	1	MET	Mainchain
1	A	53	VAL	Mainchain
1	A	62	ASN	Mainchain
1	A	63	ALA	Mainchain
1	A	64	ALA	Mainchain
1	A	75	SER	Mainchain
1	B	102	ASN	Mainchain
1	B	107	ASN	Mainchain
1	B	108	PHE	Mainchain

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group
1	B	113	SER	Mainchain
1	B	135	TYR	Mainchain
1	B	153	ARG	Mainchain
1	B	19	GLN	Mainchain
1	B	22	LYS	Mainchain
1	B	6	LYS	Mainchain
1	B	80	GLN	Mainchain
1	B	83	ASP	Mainchain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1272	0	1265	33	0
1	B	1272	0	1265	19	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	1	0	0	0	0
3	B	2	0	0	0	0
4	A	30	0	0	1	0
4	B	43	0	0	0	0
All	All	2622	0	2530	43	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (43) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:89:GLU:HG2	1:B:92:LEU:HD21	1.54	0.89
1:B:87:TRP:CE3	1:B:88:LEU:HD23	2.37	0.60
1:B:68:MET:HE2	1:B:88:LEU:HD21	1.86	0.58
1:B:105:HIS:CG	1:B:106:PRO:HD2	2.40	0.57
1:A:60:LEU:HG	4:A:471:HOH:O	2.03	0.57
1:A:124:MET:HE1	1:A:141:LYS:HA	1.88	0.55
1:A:77:LEU:HD22	1:A:82:VAL:HG21	1.89	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:86:GLU:HA	1:B:89:GLU:HG2	1.89	0.54
1:A:119:LEU:HB3	1:A:123:GLU:HB2	1.90	0.54
1:A:125:MET:HE2	1:A:137:GLU:HA	1.90	0.53
1:A:49:LYS:HB3	1:B:99:TYR:CE2	2.45	0.52
1:A:116:PHE:CZ	1:A:124:MET:HB3	2.45	0.52
1:A:124:MET:CE	1:A:144:LEU:HB2	2.40	0.52
1:A:124:MET:CE	1:A:141:LYS:HA	2.41	0.50
1:A:31:ASN:HD22	1:A:32:PRO:N	2.10	0.49
1:A:19:GLN:HG2	1:A:24:LEU:HD13	1.95	0.48
1:A:31:ASN:HD22	1:A:32:PRO:HD2	1.79	0.48
1:A:92:LEU:O	1:A:96:LYS:HB2	2.13	0.48
1:A:118:ARG:HH11	1:A:118:ARG:HG3	1.78	0.48
1:A:136:ASN:HB2	1:A:139:ASP:OD2	2.12	0.48
1:A:42:ASP:HB2	1:A:54:ARG:HD3	1.95	0.48
1:B:122:GLU:HA	1:B:125:MET:HE2	1.96	0.48
1:B:22:LYS:HE2	1:B:27:GLN:O	2.13	0.48
1:A:125:MET:HG2	1:A:135:TYR:CD2	2.49	0.47
1:A:60:LEU:HD23	1:B:75:SER:HB3	1.94	0.47
1:A:136:ASN:HB2	1:A:139:ASP:CG	2.34	0.47
1:A:93:THR:OG1	1:A:94:TYR:N	2.47	0.47
1:B:15:PHE:O	1:B:19:GLN:HG3	2.15	0.47
1:B:30:LEU:HB3	1:B:37:ASN:OD1	2.15	0.46
1:A:31:ASN:HD22	1:A:32:PRO:CD	2.28	0.46
1:B:120:GLY:O	1:B:124:MET:HG3	2.15	0.45
1:A:125:MET:CE	1:A:137:GLU:HA	2.46	0.44
1:A:124:MET:HE3	1:A:144:LEU:HB2	1.99	0.43
1:A:75:SER:HB2	1:B:60:LEU:HD22	2.00	0.43
1:A:105:HIS:CD2	1:B:41:HIS:CD2	3.08	0.42
1:B:153:ARG:O	1:B:154:LYS:C	2.58	0.42
1:A:95:LEU:HD13	1:B:84:TYR:HE2	1.85	0.41
1:A:128:MET:HE3	1:A:133:PHE:CB	2.51	0.41
1:A:124:MET:O	1:A:128:MET:HG3	2.20	0.41
1:A:60:LEU:CD2	1:B:75:SER:CB	2.98	0.41
1:A:60:LEU:HD23	1:B:75:SER:CB	2.50	0.41
1:A:111:ASP:HB3	1:B:45:LEU:HD21	2.03	0.41
1:A:3:LEU:HD23	1:A:3:LEU:HA	1.91	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	155/157 (99%)	149 (96%)	4 (3%)	2 (1%)	14	19
1	B	155/157 (99%)	144 (93%)	11 (7%)	0	100	100
All	All	310/314 (99%)	293 (94%)	15 (5%)	2 (1%)	28	41

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	5	GLY
1	A	44	GLU

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	138/138 (100%)	135 (98%)	3 (2%)	57	76
1	B	138/138 (100%)	135 (98%)	3 (2%)	57	76
All	All	276/276 (100%)	270 (98%)	6 (2%)	57	76

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2	LEU
1	A	31	ASN
1	A	96	LYS
1	B	43	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	100	THR
1	B	130	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	31	ASN
1	A	38	HIS
1	A	59	ASN
1	B	38	HIS
1	B	41	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 5 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.