



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 15, 2017 – 04:43 am GMT

PDB ID : 4ENP
Title : Structure of E530A variant E. coli KatE
Authors : Loewen, P.C.; Jha, V.
Deposited on : 2012-04-13
Resolution : 1.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	2279 (1.50-1.50)
Clashscore	112137	2503 (1.50-1.50)
Ramachandran outliers	110173	2445 (1.50-1.50)
Sidechain outliers	110143	2443 (1.50-1.50)
RSRZ outliers	101464	2305 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	753	 3% 83% 12% . .
1	B	753	 7% 82% 13% . .
1	C	753	 5% 81% 13% . .
1	D	753	 2% 82% 12% . .

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 26715 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.



- Molecule 1 is a protein called Catalase HP11.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	726	5825	3699	1018	1096	12	0	17	0
1	B	726	5789	3673	1012	1092	12	0	11	0
1	C	726	5796	3677	1016	1091	12	0	12	0
1	D	726	5789	3674	1015	1088	12	0	11	0



There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	530	ALA	GLU	ENGINEERED MUTATION	UNP P21179
B	530	ALA	GLU	ENGINEERED MUTATION	UNP P21179
C	530	ALA	GLU	ENGINEERED MUTATION	UNP P21179
D	530	ALA	GLU	ENGINEERED MUTATION	UNP P21179

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



• Molecule 1: Catalase HP11



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	93.33Å 133.11Å 122.74Å 90.00° 109.41° 90.00°	Depositor
Resolution (Å)	35.24 – 1.50 35.24 – 1.50	Depositor EDS
% Data completeness (in resolution range)	99.9 (35.24-1.50) 99.9 (35.24-1.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.17 (at 1.50Å)	Xtrriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.138 , 0.162 0.136 , 0.161	Depositor DCC
R_{free} test set	22572 reflections (5.28%)	DCC
Wilson B-factor (Å ²)	13.1	Xtrriage
Anisotropy	0.056	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 45.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.014 for h,-k,-h-l	Xtrriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	26715	wwPDB-VP
Average B, all atoms (Å ²)	17.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.99% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.34	25/6023 (0.4%)	1.37	52/8188 (0.6%)
1	B	1.27	14/5972 (0.2%)	1.25	34/8119 (0.4%)
1	C	1.30	13/5984 (0.2%)	1.36	62/8134 (0.8%)
1	D	1.35	29/5976 (0.5%)	1.35	55/8123 (0.7%)
All	All	1.31	81/23955 (0.3%)	1.33	203/32564 (0.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

The worst 5 of 81 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	377	ARG	CZ-NH1	9.10	1.44	1.33
1	C	59	ASP	CB-CG	8.98	1.70	1.51
1	C	265	SER	CB-OG	-8.40	1.31	1.42
1	A	440	TYR	CE1-CZ	8.24	1.49	1.38
1	A	344	GLU	CD-OE2	7.78	1.34	1.25

The worst 5 of 203 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	377	ARG	NE-CZ-NH2	-18.45	111.08	120.30
1	C	37	ARG	NE-CZ-NH2	-16.93	111.83	120.30
1	D	278	ARG	NE-CZ-NH1	-13.06	113.77	120.30
1	A	740	ARG	NE-CZ-NH2	-11.90	114.35	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	59	ASP	CB-CG-OD1	11.59	128.73	118.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	121	ARG	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5825	0	5671	46	0
1	B	5789	0	5618	62	0
1	C	5796	0	5631	59	0
1	D	5789	0	5632	42	0
2	A	43	0	30	0	0
2	B	43	0	30	2	0
2	C	43	0	30	2	0
2	D	43	0	30	3	0
3	A	874	0	0	14	0
3	B	776	0	0	14	0
3	C	808	0	0	24	0
3	D	886	0	0	18	0
All	All	26715	0	22672	187	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 187 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:313:ARG:HD2	3:C:1684:HOH:O	1.14	1.30
1:B:369:ARG:HD3	3:B:1673:HOH:O	1.44	1.18
1:C:267:ARG:HG3	3:C:1553:HOH:O	1.53	1.08

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:28:SER:HA	3:B:1624:HOH:O	1.53	1.06
1:C:751:ILE:HB	3:C:1456:HOH:O	1.54	1.05

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	741/753 (98%)	723 (98%)	16 (2%)	2 (0%)	44 19
1	B	735/753 (98%)	712 (97%)	21 (3%)	2 (0%)	44 19
1	C	736/753 (98%)	718 (98%)	17 (2%)	1 (0%)	55 26
1	D	735/753 (98%)	717 (98%)	17 (2%)	1 (0%)	55 26
All	All	2947/3012 (98%)	2870 (97%)	71 (2%)	6 (0%)	51 24

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	725	ASP
1	A	75	SER
1	A	725	ASP
1	C	75	SER
1	D	75	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	627/635 (99%)	616 (98%)	11 (2%)	64	33
1	B	621/635 (98%)	596 (96%)	25 (4%)	36	7
1	C	622/635 (98%)	591 (95%)	31 (5%)	28	4
1	D	621/635 (98%)	598 (96%)	23 (4%)	39	9
All	All	2491/2540 (98%)	2401 (96%)	90 (4%)	41	10

5 of 90 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	205	ILE
1	C	488[B]	ARG
1	D	616	LEU
1	C	213	LYS
1	C	377	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 23 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	252	ASN
1	C	507	HIS
1	D	629	HIS
1	C	459	ASN
1	C	572	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	HEM	A	801	1	28,50,50	2.01	4 (14%)	17,82,82	2.13	5 (29%)
2	HEM	B	801	1	28,50,50	2.11	7 (25%)	17,82,82	2.23	6 (35%)
2	HEM	C	801	1	28,50,50	1.95	6 (21%)	17,82,82	2.87	7 (41%)
2	HEM	D	801	1	28,50,50	1.76	5 (17%)	17,82,82	2.41	8 (47%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	801	1	-	0/6/54/54	0/0/8/8
2	HEM	B	801	1	-	0/6/54/54	0/0/8/8
2	HEM	C	801	1	-	0/6/54/54	0/0/8/8
2	HEM	D	801	1	-	0/6/54/54	0/0/8/8

The worst 5 of 22 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	801	HEM	CAA-C2A	-2.72	1.47	1.52
2	D	801	HEM	CAA-C2A	-2.37	1.48	1.52
2	D	801	HEM	CMD-C2D	-2.31	1.47	1.51
2	C	801	HEM	CMA-C3A	-2.20	1.47	1.51
2	C	801	HEM	CAA-C2A	-2.12	1.48	1.52

The worst 5 of 26 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	801	HEM	C1D-C2D-C3D	-6.48	102.49	107.00
2	C	801	HEM	CBD-CAD-C3D	-5.98	101.06	112.47

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	801	HEM	CBD-CAD-C3D	-5.61	101.77	112.47
2	B	801	HEM	CBD-CAD-C3D	-5.27	102.41	112.47
2	A	801	HEM	CBD-CAD-C3D	-4.57	103.76	112.47

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	801	HEM	2	0
2	C	801	HEM	2	0
2	D	801	HEM	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	726/753 (96%)	-0.33	20 (2%) 53 59	7, 13, 29, 57	1 (0%)
1	B	726/753 (96%)	-0.10	49 (6%) 19 21	7, 14, 43, 72	0
1	C	726/753 (96%)	-0.16	41 (5%) 25 28	7, 14, 41, 74	1 (0%)
1	D	726/753 (96%)	-0.29	17 (2%) 61 66	7, 13, 31, 69	0
All	All	2904/3012 (96%)	-0.22	127 (4%) 35 40	7, 13, 38, 74	2 (0%)

The worst 5 of 127 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	726	GLY	6.1
1	B	32	GLU	5.0
1	B	673	ALA	4.6
1	C	714	GLY	4.5
1	C	675	ILE	4.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	HEM	A	801	43/43	0.99	0.10	0.55	6,8,9,15	0
2	HEM	C	801	43/43	0.99	0.08	0.23	7,8,11,16	0
2	HEM	B	801	43/43	0.99	0.06	-0.46	7,9,11,14	0
2	HEM	D	801	43/43	0.99	0.06	-0.60	7,8,10,15	0

6.5 Other polymers [i](#)

There are no such residues in this entry.