



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 03:35 am GMT

PDB ID : 5EN0  
Title : Crystal Structure of T94I rhodopsin mutant  
Authors : Singhal, A.; Guo, Y.; Matkovic, M.; Schertler, G.; Deupi, X.; Yan, E.; Standfuss, J.  
Deposited on : 2015-11-08  
Resolution : 2.81 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.81 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	2917 (2.84-2.80)
Clashscore	112137	3382 (2.84-2.80)
Ramachandran outliers	110173	3324 (2.84-2.80)
Sidechain outliers	110143	3326 (2.84-2.80)
RSRZ outliers	101464	2948 (2.84-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	349	<div> <div>5%</div> <div> <div></div> <div>83%</div> <div>9%</div> <div>6%</div> </div> </div>
2	B	11	<div> <div>9%</div> <div> <div></div> <div>73%</div> <div>18%</div> <div>9%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	RET	A	401	-	-	-	X
4	ACT	A	402	-	-	-	X
5	BOG	A	404	-	-	-	X
5	BOG	A	411	-	-	-	X
6	PLM	A	405	-	-	-	X

## 2 Entry composition

There are 11 unique types of molecules in this entry. The entry contains 2863 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Rhodopsin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	327	Total	C	N	O	S	0	0	0
			2592	1727	399	438	28			


There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	ACE	-	acetylation	UNP P02699
A	2	CYS	ASN	engineered mutation	UNP P02699
A	94	ILE	THR	engineered mutation	UNP P02699
A	282	CYS	ASP	engineered mutation	UNP P02699

- Molecule 2 is a protein called Guanine nucleotide-binding protein G(t) subunit alpha-3.


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	11	Total	C	N	O	0	0	0
			89	59	13	17			

- Molecule 3 is RETINAL (three-letter code: RET) (formula: C<sub>20</sub>H<sub>28</sub>O).




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	C	0	0
			20	20		

- Molecule 4 is ACETATE ION (three-letter code: ACT) (formula:  $C_2H_3O_2$ ).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is B-OCTYLGLUCOSIDE (three-letter code: BOG) (formula:  $C_{14}H_{28}O_6$ ).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			20	14	6		
5	A	1	Total	C	O	0	0
			20	14	6		

- Molecule 6 is PALMITIC ACID (three-letter code: PLM) (formula: C<sub>16</sub>H<sub>32</sub>O<sub>2</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			17	16	1		
6	A	1	Total	C	O	0	0
			17	16	1		

- Molecule 7 is N-ACETYL-D-GLUCOSAMINE (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	A	1	Total	C	N	O	0	0
			14	8	1	5		
7	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 8 is BETA-D-MANNOSE (three-letter code: BMA) (formula:  $C_6H_{12}O_6$ ).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	A	1	Total	C	O	0	0
			11	6	5		

- Molecule 9 is ALPHA-D-MANNOSE (three-letter code: MAN) (formula:  $C_6H_{12}O_6$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	A	1	Total	C	O	0	0
			11	6	5		

- Molecule 10 is SULFATE ION (three-letter code: SO4) (formula:  $O_4S$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
10	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 11 is water.




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
11	A	25	Total 25	O 25	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Rhodopsin



#### • Molecule 2: Guanine nucleotide-binding protein G(t) subunit alpha-3



## 4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	242.27Å 242.27Å 111.15Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	30.00 – 2.81 29.31 – 2.81	Depositor EDS
% Data completeness (in resolution range)	99.3 (30.00-2.81) 99.4 (29.31-2.81)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.94 (at 2.80Å)	Xtriage
Refinement program	REFMAC 5.8.0131	Depositor
R, $R_{free}$	0.207 , 0.253 0.220 , 0.264	Depositor DCC
$R_{free}$ test set	1565 reflections (5.44%)	DCC
Wilson B-factor (Å <sup>2</sup> )	60.4	Xtriage
Anisotropy	0.158	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 53.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	2863	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	59.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.75% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, NAG, ACE, RET, SO4, ACT, PLM, BOG, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.81	1/2672 (0.0%)	0.94	5/3640 (0.1%)
2	B	0.79	0/89	1.10	0/117
All	All	0.81	1/2761 (0.0%)	0.95	5/3757 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1	MET	N-CA	6.41	1.59	1.46

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	282	CYS	CA-CB-SG	-10.07	95.88	114.00
1	A	322	CYS	CA-CB-SG	-9.69	96.56	114.00
1	A	2	CYS	CA-CB-SG	-9.51	96.89	114.00
1	A	257	MET	CG-SD-CE	-6.68	89.52	100.20
1	A	252	ARG	NE-CZ-NH2	-5.21	117.69	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2592	0	2569	12	0
2	B	89	0	93	1	0
3	A	20	0	27	1	0
4	A	8	0	6	0	0
5	A	40	0	56	0	0
6	A	34	0	62	0	0
7	A	28	0	24	0	0
8	A	11	0	9	0	0
9	A	11	0	10	0	0
10	A	5	0	0	0	0
11	A	25	0	0	1	0
All	All	2863	0	2856	14	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (14) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:176:SER:O	11:A:501:HOH:O	2.04	0.74
3:A:401:RET:H181	3:A:401:RET:C8	2.33	0.57
1:A:6:GLY:HA3	1:A:9:PHE:CZ	2.43	0.54
1:A:277:THR:O	1:A:278:HIS:ND1	2.44	0.50
1:A:193:THR:HG22	1:A:195:HIS:HB2	1.93	0.50
2:B:343:ASN:O	2:B:347:VAL:HB	2.13	0.48
1:A:94:ILE:HG12	1:A:186:SER:HB3	1.97	0.47
1:A:135:ARG:HH12	1:A:257:MET:CE	2.28	0.46
1:A:96:TYR:O	1:A:100:HIS:HD2	1.99	0.46
1:A:93:THR:HG21	1:A:113:GLU:HG3	1.99	0.44
1:A:135:ARG:HH12	1:A:257:MET:HE1	1.83	0.43
1:A:61:VAL:HG13	1:A:68:LEU:CD1	2.50	0.42
1:A:41:ALA:HB1	1:A:95:LEU:HD12	2.02	0.41
1:A:216:LEU:HD12	1:A:216:LEU:HA	1.90	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	325/349 (93%)	301 (93%)	24 (7%)	0	100	100
2	B	9/11 (82%)	9 (100%)	0	0	100	100
All	All	334/360 (93%)	310 (93%)	24 (7%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	278/296 (94%)	265 (95%)	13 (5%)	30	63
2	B	10/10 (100%)	8 (80%)	2 (20%)	1	4
All	All	288/306 (94%)	273 (95%)	15 (5%)	27	59

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2	CYS
1	A	7	PRO
1	A	61	VAL
1	A	66	LYS
1	A	108	THR
1	A	154	ILE
1	A	167	CYS
1	A	215	PRO

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	226	LEU
1	A	249	GLU
1	A	293	PHE
1	A	323	CYS
1	A	326	ASN
2	B	340	ILE
2	B	347	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	145	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
1	ACE	A	0	1	2,2,2	0.94	0	1,1,1	1.20	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	ACE	A	0	1	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	RET	A	401	1	19,20,21	0.76	0	27,27,28	1.65	9 (33%)
4	ACT	A	402	-	1,3,3	4.90	1 (100%)	0,3,3	0.00	-
4	ACT	A	403	-	1,3,3	2.61	1 (100%)	0,3,3	0.00	-
5	BOG	A	404	-	20,20,20	0.75	1 (5%)	25,25,25	1.69	7 (28%)
6	PLM	A	405	1	16,16,17	0.64	0	15,15,17	0.49	0
6	PLM	A	406	1	16,16,17	0.52	0	15,15,17	0.98	1 (6%)
7	NAG	A	407	1,7	14,14,15	0.61	0	15,19,21	1.42	2 (13%)
7	NAG	A	408	8,7	14,14,15	1.21	1 (7%)	15,19,21	1.77	3 (20%)
8	BMA	A	409	9,7	11,11,12	1.23	1 (9%)	13,15,17	2.78	7 (53%)
9	MAN	A	410	8	11,11,12	0.67	0	13,15,17	2.22	2 (15%)
5	BOG	A	411	-	20,20,20	0.86	1 (5%)	25,25,25	1.91	7 (28%)
10	SO4	A	412	-	4,4,4	0.48	0	6,6,6	0.39	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical



component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	RET	A	401	1	-	0/13/30/31	0/1/1/1
4	ACT	A	402	-	-	0/0/0/0	0/0/0/0
4	ACT	A	403	-	-	0/0/0/0	0/0/0/0
5	BOG	A	404	-	-	0/11/31/31	0/1/1/1
6	PLM	A	405	1	-	0/13/14/15	0/0/0/0
6	PLM	A	406	1	-	0/13/14/15	0/0/0/0
7	NAG	A	407	1,7	-	0/6/23/26	0/1/1/1
7	NAG	A	408	8,7	-	0/6/23/26	0/1/1/1
8	BMA	A	409	9,7	-	0/2/19/22	0/1/1/1
9	MAN	A	410	8	-	0/2/19/22	0/1/1/1
5	BOG	A	411	-	-	0/11/31/31	0/1/1/1
10	SO4	A	412	-	-	0/0/0/0	0/0/0/0

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
8	A	409	BMA	O5-C1	2.00	1.47	1.43
5	A	404	BOG	O1-C1	2.22	1.44	1.40
5	A	411	BOG	O1-C1	2.35	1.44	1.40
4	A	403	ACT	CH3-C	2.61	1.52	1.48
7	A	408	NAG	O5-C1	2.83	1.48	1.43
4	A	402	ACT	CH3-C	4.90	1.55	1.48

All (38) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
9	A	410	MAN	O5-C1-C2	-4.24	104.14	110.79
5	A	411	BOG	O5-C5-C4	-4.11	102.08	109.66
8	A	409	BMA	O5-C1-C2	-3.90	104.68	110.79
5	A	411	BOG	O3-C3-C4	-3.78	102.14	110.36
3	A	401	RET	C8-C9-C10	-3.31	113.87	118.94
5	A	404	BOG	C4-C3-C2	-3.25	105.10	110.84
3	A	401	RET	C18-C5-C6	-3.20	120.93	124.51
5	A	404	BOG	C1-C2-C3	-3.02	104.37	109.98
3	A	401	RET	C11-C10-C9	-2.72	123.43	127.31
7	A	408	NAG	O3-C3-C2	-2.68	103.64	109.39
8	A	409	BMA	O3-C3-C2	-2.50	105.48	110.02
5	A	404	BOG	C3-C4-C5	-2.49	105.83	110.22
3	A	401	RET	C11-C12-C13	-2.45	119.55	126.42
6	A	406	PLM	C4-C3-C2	-2.41	104.48	113.74

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	401	RET	C10-C11-C12	-2.41	115.85	123.23
5	A	411	BOG	O4-C4-C3	-2.40	105.13	110.36
3	A	401	RET	C8-C7-C6	-2.39	120.56	127.25
3	A	401	RET	C7-C8-C9	-2.28	122.79	126.21
8	A	409	BMA	O2-C2-C1	-2.26	104.57	109.18
8	A	409	BMA	O2-C2-C3	-2.13	106.00	110.17
5	A	404	BOG	O2-C2-C3	2.11	114.95	110.36
8	A	409	BMA	C2-C3-C4	2.12	114.57	110.88
7	A	407	NAG	C2-N2-C7	2.13	126.06	122.94
3	A	401	RET	C18-C5-C4	2.17	117.56	113.45
5	A	411	BOG	O4-C4-C5	2.21	114.86	109.28
5	A	404	BOG	O3-C3-C4	2.24	115.24	110.36
5	A	411	BOG	O5-C1-C2	2.28	114.70	110.30
5	A	411	BOG	C1-C2-C3	2.32	114.30	109.98
3	A	401	RET	C19-C9-C8	2.64	122.30	118.10
5	A	404	BOG	C1-O5-C5	2.73	118.85	113.72
7	A	407	NAG	O7-C7-N2	2.73	127.17	121.92
5	A	404	BOG	O5-C5-C6	2.79	113.09	106.41
7	A	408	NAG	O3-C3-C4	2.81	116.47	110.36
8	A	409	BMA	C1-C2-C3	4.43	115.27	109.65
7	A	408	NAG	O6-C6-C5	4.51	126.52	111.34
5	A	411	BOG	C4-C3-C2	4.70	119.14	110.84
8	A	409	BMA	C3-C4-C5	5.80	120.43	110.22
9	A	410	MAN	C1-O5-C5	6.41	121.00	112.17

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	401	RET	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	326/349 (93%)	-0.11	16 (4%) 30 20	36, 55, 81, 120	0
2	B	11/11 (100%)	0.55	1 (9%) 10 5	55, 66, 87, 92	0
All	All	337/360 (93%)	-0.09	17 (5%) 30 20	36, 55, 84, 120	0

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	145	ASN	3.8
1	A	151	ASN	3.5
1	A	147	ARG	3.4
1	A	278	HIS	3.3
1	A	281	SER	3.2
1	A	199	ASN	2.8
1	A	274	TYR	2.8
1	A	146	PHE	2.8
1	A	277	THR	2.7
1	A	276	PHE	2.5
1	A	279	GLN	2.4
1	A	144	SER	2.3
1	A	196	GLU	2.2
1	A	186	SER	2.1
1	A	245	LYS	2.1
1	A	142	PRO	2.0
2	B	341	LEU	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
1	ACE	A	0	3/3	0.46	0.38	-	83,83,106,109	0

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
5	BOG	A	411	20/20	0.88	0.40	6.00	63,82,98,101	0
6	PLM	A	405	17/18	0.90	0.37	5.59	71,93,113,123	0
3	RET	A	401	20/21	0.82	0.44	4.30	85,95,117,118	0
5	BOG	A	404	20/20	0.81	0.40	3.98	66,107,139,149	0
4	ACT	A	402	4/4	0.87	0.60	3.33	76,80,92,94	0
6	PLM	A	406	17/18	0.86	0.20	1.43	55,65,99,104	0
10	SO4	A	412	5/5	0.92	0.18	-0.03	101,102,123,126	0
7	NAG	A	407	14/15	0.96	0.17	-0.05	39,41,50,56	0
9	MAN	A	410	11/12	0.94	0.42	-	60,74,85,90	0
4	ACT	A	403	4/4	0.69	0.95	-	80,86,89,93	0
7	NAG	A	408	14/15	0.96	0.28	-	30,32,43,58	0
8	BMA	A	409	11/12	0.90	0.36	-	61,74,81,82	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.