



Full wwPDB NMR Structure Validation Report ⓘ

Feb 12, 2017 – 05:58 pm GMT

PDB ID : 1EOQ
Title : ROUS SARCOMA VIRUS CAPSID PROTEIN: C-TERMINAL DOMAIN
Authors : Kingston, R.L.; Fitzon-Ostendorp, T.; Eisenmesser, E.Z.; Schatz, G.W.; Vogt, V.M.; Post, C.B.; Rossman, M.G.
Deposited on : 2000-03-23

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)
NmrClust : Kelley et al. (1996)
MolProbity : 4.02b-467
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : trunk28760
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 82%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	125131	11601
Ramachandran outliers	121729	10391
Sidechain outliers	121581	10367

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	96	 35% 41% • 20%

2 Ensemble composition and analysis

This entry contains 1 models. Identification of well-defined residues and clustering analysis are not possible.

3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 1209 atoms, of which 613 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called GAG POLYPROTEIN CAPSID PROTEIN P27.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	77	1209	376	613	103	114	3	0

There is a discrepancy between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	154	MET	-	INITIATING MET	UNP P03322

4 Residue-property plots [i](#)

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: GAG POLYPROTEIN CAPSID PROTEIN P27

Chain A: 



5 Refinement protocol and experimental data overview

The models were refined using the following method: *torsion angle dynamics/simulated annealing*.

Of the ? calculated structures, 1 were deposited, based on the following criterion: ?.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CNS	refinement	1.0

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	BMRB entry 4593
Number of chemical shift lists	1
Total number of shifts	886
Number of shifts mapped to atoms	886
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	82%

No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality i

6.1 Standard geometry i

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	596	613	610	23
All	All	596	613	610	23

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 19.

All clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)
1:A:204:LEU:O	1:A:208:ALA:HB2	0.99	1.57
1:A:218:ILE:O	1:A:222:VAL:HG23	0.83	1.72
1:A:164:PHE:CE2	1:A:222:VAL:HG21	0.76	2.15
1:A:186:ALA:HB3	1:A:187:PRO:HD3	0.72	1.60
1:A:204:LEU:O	1:A:208:ALA:CB	0.67	2.41
1:A:168:ALA:O	1:A:172:ILE:HG12	0.66	1.90
1:A:184:ALA:O	1:A:188:VAL:HG23	0.66	1.90
1:A:171:LEU:HD23	1:A:189:ILE:HG23	0.62	1.72
1:A:186:ALA:HB3	1:A:187:PRO:CD	0.58	2.28
1:A:171:LEU:CD2	1:A:189:ILE:HG23	0.58	2.29
1:A:180:LEU:HD13	1:A:188:VAL:HG21	0.57	1.74
1:A:189:ILE:CG2	1:A:193:PHE:CE2	0.51	2.94
1:A:171:LEU:O	1:A:175:VAL:HG23	0.47	2.09
1:A:212:LEU:HD21	1:A:221:TYR:CD2	0.45	2.46
1:A:219:ILE:N	1:A:219:ILE:HD12	0.45	2.27

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)
1:A:156:ILE:HG22	1:A:167:PHE:CZ	0.44	2.48
1:A:191:ASP:O	1:A:195:GLN:HG2	0.43	2.14
1:A:219:ILE:HD12	1:A:219:ILE:H	0.41	1.74
1:A:171:LEU:HD11	1:A:192:CYS:SG	0.41	2.56
1:A:192:CYS:O	1:A:196:LYS:N	0.41	2.54
1:A:156:ILE:HG22	1:A:167:PHE:CE1	0.40	2.51
1:A:193:PHE:O	1:A:197:SER:HB2	0.40	2.16
1:A:194:ARG:HG3	1:A:205:ILE:CG2	0.40	2.45

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	75/96 (78%)	66 (88%)	7 (9%)	2 (3%)	10	45
All	All	75/96 (78%)	66 (88%)	7 (9%)	2 (3%)	10	45

All 2 Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type
1	A	210	SER
1	A	209	PRO

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	68/82 (83%)	51 (75%)	17 (25%)	3	25
All	All	68/82 (83%)	51 (75%)	17 (25%)	3	25

All 17 residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type
1	A	161	SER
1	A	207	THR
1	A	179	ASP
1	A	178	SER
1	A	210	SER
1	A	197	SER
1	A	163	SER
1	A	198	GLN
1	A	169	ASN
1	A	191	ASP
1	A	192	CYS
1	A	154	MET
1	A	206	ARG
1	A	220	LYS
1	A	203	GLN
1	A	225	ARG
1	A	211	THR

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues

There are no chain breaks in this entry.

7 Chemical shift validation [i](#)

The completeness of assignment taking into account all chemical shift lists is 82% for the well-defined parts and 82% for the entire structure.

7.1 Chemical shift list 1

File name: BMRB entry 4593

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	886
Number of shifts mapped to atoms	886
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	1

7.1.2 Chemical shift referencing [i](#)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction \pm precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	75	-0.57 ± 0.15	Should be applied
$^{13}\text{C}_\beta$	73	0.25 ± 0.12	None needed (< 0.5 ppm)
$^{13}\text{C}'$	0	—	None (insufficient data)
^{15}N	68	0.79 ± 0.34	Should be applied

7.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 82%, i.e. 783 atoms were assigned a chemical shift out of a possible 952. 1 out of 9 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	287/369 (78%)	144/146 (99%)	75/154 (49%)	68/69 (99%)
Sidechain	473/548 (86%)	293/324 (90%)	168/198 (85%)	12/26 (46%)

Continued on next page...

Continued from previous page...

	Total	¹ H	¹³ C	¹⁵ N
Aromatic	23/35 (66%)	19/19 (100%)	4/16 (25%)	0/0 (—%)
Overall	783/952 (82%)	456/489 (93%)	247/368 (67%)	80/95 (84%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 82%, i.e. 783 atoms were assigned a chemical shift out of a possible 952. 1 out of 9 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	287/369 (78%)	144/146 (99%)	75/154 (49%)	68/69 (99%)
Sidechain	473/548 (86%)	293/324 (90%)	168/198 (85%)	12/26 (46%)
Aromatic	23/35 (66%)	19/19 (100%)	4/16 (25%)	0/0 (—%)
Overall	783/952 (82%)	456/489 (93%)	247/368 (67%)	80/95 (84%)

7.1.4 Statistically unusual chemical shifts [i](#)

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

Mol	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	158	GLN	HB2	0.09	3.30 – 0.80	-7.8

7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:

