



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 14, 2017 – 09:49 am GMT

PDB ID : 5EO7  
Title : Crystal structure of AOL  
Authors : Kato, R.; Kiso, M.; Ishida, H.; Ando, H.; Suzuki, T.; Shimabukuro, S.; Makyio, H.  
Deposited on : 2015-11-10  
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	4130 (2.30-2.30)
Clashscore	112137	4751 (2.30-2.30)
Ramachandran outliers	110173	4705 (2.30-2.30)
Sidechain outliers	110143	4704 (2.30-2.30)
RSRZ outliers	101464	4156 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	310	<div> <div>9%</div> <div>97%</div> </div>
1	B	310	<div> <div>4%</div> <div>96%</div> </div>
1	C	310	<div> <div>3%</div> <div>95%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	SFU	A	404	-	-	-	X
2	SFU	A	405	-	-	-	X
2	SFU	B	401	-	-	-	X
2	SFU	B	404	-	-	-	X
2	SFU	B	405	-	-	-	X
2	SFU	C	401	-	-	-	X
2	SFU	C	403	-	-	-	X
2	SFU	C	404	-	-	-	X
2	SFU	C	405	-	-	-	X
3	SO4	A	408	-	-	-	X

## 2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 15101 atoms, of which 7255 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Predicted protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	310	Total	C	H	N	O	S	0	0	0
			4785	1551	2349	423	457	5			
1	B	310	Total	C	H	N	O	S	0	0	0
			4784	1551	2348	423	457	5			
1	C	310	Total	C	H	N	O	S	0	0	0
			4784	1551	2348	423	457	5			

- Molecule 2 is METHYL 6-DEOXY-1-SELENO-ALPHA-L-GALACTOPYRANOSIDE (three-letter code: SFU) (formula: C<sub>7</sub>H<sub>14</sub>O<sub>4</sub>Se).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	H	O	Se	0	0
			26	7	14	4	1		
2	A	1	Total	C	H	O	Se	0	0
			26	7	14	4	1		
2	A	1	Total	C	H	O	Se	0	0
			26	7	14	4	1		

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	H	O	Se	0	0
			26	7	14	4	1		
2	A	1	Total	C	H	O	Se	0	0
			26	7	14	4	1		
2	B	1	Total	C	H	O	Se	0	0
			26	7	14	4	1		
2	B	1	Total	C	H	O	Se	0	0
			26	7	14	4	1		
2	B	1	Total	C	H	O	Se	0	0
			26	7	14	4	1		
2	B	1	Total	C	H	O	Se	0	0
			26	7	14	4	1		
2	B	1	Total	C	H	O	Se	0	0
			26	7	14	4	1		
2	C	1	Total	C	H	O	Se	0	0
			26	7	14	4	1		
2	C	1	Total	C	H	O	Se	0	0
			26	7	14	4	1		
2	C	1	Total	C	H	O	Se	0	0
			26	7	14	4	1		
2	C	1	Total	C	H	O	Se	0	0
			26	7	14	4	1		
2	C	1	Total	C	H	O	Se	0	0
			26	7	14	4	1		

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	0
3	C	1	Total O S 5 4 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	137	Total O 137 137	0	0
4	B	101	Total O 101 101	0	0
4	C	100	Total O 100 100	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: Predicted protein



#### • Molecule 1: Predicted protein



#### • Molecule 1: Predicted protein



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	125.16Å 196.85Å 110.61Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.68 – 2.30 38.68 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.8 (38.68-2.30) 92.8 (38.68-2.30)	Depositor EDS
$R_{merge}$	0.14	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.74 (at 2.29Å)	Xtriage
Refinement program	PHENIX	Depositor
R, $R_{free}$	0.219 , 0.241 0.204 , 0.228	Depositor DCC
$R_{free}$ test set	2860 reflections (5.07%)	DCC
Wilson B-factor (Å <sup>2</sup> )	24.2	Xtriage
Anisotropy	1.892	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.42 , 39.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	15101	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.48% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SFU, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.25	0/2504	0.46	0/3413
1	B	0.23	0/2504	0.45	0/3413
1	C	0.24	0/2504	0.46	0/3413
All	All	0.24	0/7512	0.46	0/10239

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2436	2349	2348	9	0
1	B	2436	2348	2348	9	0
1	C	2436	2348	2348	10	0
2	A	60	70	50	0	0
2	B	60	70	50	0	0
2	C	60	70	50	2	0
3	A	15	0	0	0	0
3	C	5	0	0	0	0
4	A	137	0	0	5	0
4	B	101	0	0	7	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	C	100	0	0	6	0
All	All	7846	7255	7194	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:8:GLU:OE2	4:A:501:HOH:O	2.07	0.72
1:C:144:ASP:OD1	4:C:501:HOH:O	2.08	0.70
1:B:198:TRP:O	4:B:501:HOH:O	2.12	0.67
2:C:402:SFU:SE	4:C:593:HOH:O	2.61	0.67
1:A:183:ASP:O	4:A:502:HOH:O	2.12	0.67
1:B:225:SER:N	4:B:504:HOH:O	2.25	0.65
1:A:97:GLY:O	4:A:503:HOH:O	2.15	0.64
1:C:308:PRO:O	4:C:502:HOH:O	2.16	0.62
1:B:299:GLY:O	4:B:502:HOH:O	2.16	0.61
1:C:119:GLY:HA3	1:C:167:ARG:HD2	1.85	0.59
1:A:239:THR:HG22	1:A:259:PRO:HA	1.86	0.58
1:A:239:THR:HG21	4:A:553:HOH:O	2.07	0.53
1:B:7:GLN:NE2	4:B:512:HOH:O	2.41	0.53
1:C:2:SER:N	4:C:515:HOH:O	2.44	0.51
1:A:110:SER:O	4:A:505:HOH:O	2.20	0.49
1:C:253:ASP:OD1	4:C:503:HOH:O	2.20	0.47
1:C:85:GLU:OE2	2:C:402:SFU:O3	2.29	0.47
1:B:41:GLU:HA	4:B:538:HOH:O	2.14	0.47
1:B:202:LEU:HD11	4:B:562:HOH:O	2.14	0.46
1:A:169:THR:HB	1:A:224:LYS:HA	1.99	0.44
1:B:273:GLY:N	1:B:274:PRO:HA	2.32	0.44
1:C:109:TYR:OH	4:C:504:HOH:O	2.21	0.43
1:B:288:ALA:O	4:B:503:HOH:O	2.21	0.42
1:B:77:LEU:HD13	1:B:112:ILE:HD11	2.00	0.42
1:C:273:GLY:N	1:C:274:PRO:HA	2.34	0.42
1:A:239:THR:HG22	1:A:259:PRO:CA	2.49	0.42
1:A:2:SER:OG	1:A:3:THR:N	2.54	0.41
1:C:77:LEU:HD13	1:C:112:ILE:HD11	2.03	0.41
1:C:75:TYR:CZ	1:C:127:ILE:HD11	2.56	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	308/310 (99%)	298 (97%)	10 (3%)	0	100	100
1	B	308/310 (99%)	300 (97%)	8 (3%)	0	100	100
1	C	308/310 (99%)	301 (98%)	7 (2%)	0	100	100
All	All	924/930 (99%)	899 (97%)	25 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	251/251 (100%)	251 (100%)	0	100	100
1	B	251/251 (100%)	251 (100%)	0	100	100
1	C	251/251 (100%)	250 (100%)	1 (0%)	93	97
All	All	753/753 (100%)	752 (100%)	1 (0%)	94	98

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	77	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	7	GLN
1	C	7	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

19 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	SFU	A	401	-	10,12,12	0.70	0	14,17,17	2.30	7 (50%)
2	SFU	A	402	-	10,12,12	0.63	0	14,17,17	1.82	6 (42%)
2	SFU	A	403	-	10,12,12	0.65	0	14,17,17	1.67	6 (42%)
2	SFU	A	404	-	10,12,12	0.72	0	14,17,17	1.58	4 (28%)
2	SFU	A	405	-	10,12,12	0.70	0	14,17,17	1.72	6 (42%)
3	SO4	A	406	-	4,4,4	0.17	0	6,6,6	0.19	0
3	SO4	A	407	-	4,4,4	0.19	0	6,6,6	0.08	0
3	SO4	A	408	-	4,4,4	0.20	0	6,6,6	0.10	0
2	SFU	B	401	-	10,12,12	0.69	0	14,17,17	1.68	6 (42%)
2	SFU	B	402	-	10,12,12	0.67	0	14,17,17	1.78	6 (42%)
2	SFU	B	403	-	10,12,12	0.54	0	14,17,17	1.70	4 (28%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	SFU	B	404	-	10,12,12	0.66	0	14,17,17	1.72	5 (35%)
2	SFU	B	405	-	10,12,12	0.60	0	14,17,17	1.80	6 (42%)
2	SFU	C	401	-	10,12,12	0.67	0	14,17,17	1.78	6 (42%)
2	SFU	C	402	-	10,12,12	0.99	0	14,17,17	1.79	5 (35%)
2	SFU	C	403	-	10,12,12	0.67	0	14,17,17	1.73	4 (28%)
2	SFU	C	404	-	10,12,12	0.73	0	14,17,17	1.76	6 (42%)
2	SFU	C	405	-	10,12,12	0.73	0	14,17,17	1.71	6 (42%)
3	SO4	C	406	-	4,4,4	0.17	0	6,6,6	0.11	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SFU	A	401	-	-	0/0/22/22	0/1/1/1
2	SFU	A	402	-	-	0/0/22/22	0/1/1/1
2	SFU	A	403	-	-	0/0/22/22	0/1/1/1
2	SFU	A	404	-	-	0/0/22/22	0/1/1/1
2	SFU	A	405	-	-	0/0/22/22	0/1/1/1
3	SO4	A	406	-	-	0/0/0/0	0/0/0/0
3	SO4	A	407	-	-	0/0/0/0	0/0/0/0
3	SO4	A	408	-	-	0/0/0/0	0/0/0/0
2	SFU	B	401	-	-	0/0/22/22	0/1/1/1
2	SFU	B	402	-	-	0/0/22/22	0/1/1/1
2	SFU	B	403	-	-	0/0/22/22	0/1/1/1
2	SFU	B	404	-	-	0/0/22/22	0/1/1/1
2	SFU	B	405	-	-	0/0/22/22	0/1/1/1
2	SFU	C	401	-	-	0/0/22/22	0/1/1/1
2	SFU	C	402	-	-	0/0/22/22	0/1/1/1
2	SFU	C	403	-	-	0/0/22/22	0/1/1/1
2	SFU	C	404	-	-	0/0/22/22	0/1/1/1
2	SFU	C	405	-	-	0/0/22/22	0/1/1/1
3	SO4	C	406	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

All (83) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	403	SFU	C4-C3-C2	-3.17	105.25	110.84

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	405	SFU	C4-C3-C2	-2.96	105.61	110.84
2	A	402	SFU	O2-C2-C3	-2.83	104.20	110.36
2	B	405	SFU	O2-C2-C3	-2.78	104.30	110.36
2	C	401	SFU	O2-C2-C3	-2.77	104.33	110.36
2	C	401	SFU	C4-C3-C2	-2.77	105.96	110.84
2	B	402	SFU	O2-C2-C3	-2.74	104.39	110.36
2	C	404	SFU	O2-C2-C3	-2.72	104.44	110.36
2	A	401	SFU	O2-C2-C3	-2.72	104.45	110.36
2	C	402	SFU	O2-C2-C3	-2.68	104.52	110.36
2	A	403	SFU	C4-C3-C2	-2.67	106.12	110.84
2	B	405	SFU	C4-C3-C2	-2.67	106.12	110.84
2	B	401	SFU	O2-C2-C3	-2.66	104.57	110.36
2	A	405	SFU	C4-C3-C2	-2.63	106.19	110.84
2	B	404	SFU	O2-C2-C3	-2.61	104.67	110.36
2	A	405	SFU	O2-C2-C3	-2.51	104.89	110.36
2	B	404	SFU	C4-C3-C2	-2.47	106.48	110.84
2	A	403	SFU	O2-C2-C3	-2.41	105.11	110.36
2	C	405	SFU	O2-C2-C3	-2.38	105.18	110.36
2	A	404	SFU	O2-C2-C3	-2.33	105.28	110.36
2	B	401	SFU	C4-C3-C2	-2.26	106.84	110.84
2	B	402	SFU	C4-C3-C2	-2.26	106.85	110.84
2	A	402	SFU	C4-C3-C2	-2.25	106.86	110.84
2	B	403	SFU	O2-C2-C3	-2.20	105.56	110.36
2	A	401	SFU	C4-C3-C2	-2.07	107.18	110.84
2	C	401	SFU	O5-C5-C6	2.00	110.92	106.69
2	A	401	SFU	C3-C4-C5	2.01	112.85	109.68
2	C	405	SFU	O5-C1-C2	2.10	113.72	110.16
2	C	403	SFU	C5-O5-C1	2.11	115.59	112.78
2	A	403	SFU	C5-O5-C1	2.11	115.60	112.78
2	A	402	SFU	O5-C5-C6	2.13	111.19	106.69
2	B	405	SFU	O5-C5-C6	2.13	111.19	106.69
2	C	404	SFU	O5-C5-C6	2.19	111.32	106.69
2	A	404	SFU	O5-C5-C6	2.20	111.34	106.69
2	B	404	SFU	C5-O5-C1	2.20	115.72	112.78
2	C	404	SFU	O5-C1-C2	2.21	113.91	110.16
2	B	401	SFU	C5-O5-C1	2.22	115.75	112.78
2	A	405	SFU	C5-O5-C1	2.30	115.85	112.78
2	C	401	SFU	O5-C1-C2	2.30	114.06	110.16
2	C	403	SFU	C3-C2-C1	2.31	112.95	109.22
2	C	404	SFU	O2-C2-C1	2.31	113.73	109.29
2	B	403	SFU	O5-C5-C6	2.32	111.60	106.69
2	A	405	SFU	O2-C2-C1	2.33	113.77	109.29

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	401	SFU	O5-C5-C6	2.38	111.72	106.69
2	C	402	SFU	C3-C4-C5	2.38	113.42	109.68
2	B	401	SFU	O2-C2-C1	2.40	113.89	109.29
2	A	403	SFU	O2-C2-C1	2.40	113.90	109.29
2	C	405	SFU	C5-O5-C1	2.41	116.00	112.78
2	C	404	SFU	C5-O5-C1	2.43	116.03	112.78
2	A	401	SFU	O2-C2-C1	2.44	113.97	109.29
2	A	403	SFU	O5-C5-C6	2.46	111.89	106.69
2	B	405	SFU	C5-O5-C1	2.46	116.07	112.78
2	B	402	SFU	C5-O5-C1	2.47	116.08	112.78
2	A	401	SFU	O5-C5-C4	2.48	113.71	109.62
2	C	401	SFU	O2-C2-C1	2.49	114.07	109.29
2	A	404	SFU	O5-C1-C2	2.52	114.42	110.16
2	B	402	SFU	O2-C2-C1	2.54	114.16	109.29
2	C	405	SFU	O5-C5-C6	2.55	112.08	106.69
2	A	404	SFU	O2-C2-C1	2.57	114.22	109.29
2	A	405	SFU	O5-C5-C6	2.61	112.22	106.69
2	A	402	SFU	O2-C2-C1	2.62	114.32	109.29
2	B	405	SFU	O2-C2-C1	2.62	114.32	109.29
2	C	402	SFU	C3-C2-C1	2.66	113.53	109.22
2	B	404	SFU	O5-C5-C6	2.67	112.33	106.69
2	B	401	SFU	O5-C1-C2	2.72	114.77	110.16
2	C	401	SFU	C5-O5-C1	2.74	116.43	112.78
2	A	403	SFU	O5-C1-C2	2.78	114.87	110.16
2	C	402	SFU	O5-C5-C6	2.80	112.62	106.69
2	B	402	SFU	O5-C5-C6	2.81	112.65	106.69
2	C	403	SFU	O2-C2-C1	2.82	114.70	109.29
2	A	405	SFU	O5-C1-C2	2.84	114.97	110.16
2	B	403	SFU	O2-C2-C1	2.88	114.81	109.29
2	C	403	SFU	C3-C4-C5	2.91	114.25	109.68
2	C	405	SFU	O2-C2-C1	2.92	114.89	109.29
2	B	402	SFU	O5-C1-C2	3.06	115.34	110.16
2	C	404	SFU	C3-C4-C5	3.09	114.53	109.68
2	A	402	SFU	C5-O5-C1	3.09	116.91	112.78
2	B	404	SFU	O2-C2-C1	3.12	115.28	109.29
2	C	402	SFU	C5-O5-C1	3.13	116.96	112.78
2	A	402	SFU	O5-C1-C2	3.13	115.47	110.16
2	B	405	SFU	O5-C1-C2	3.27	115.70	110.16
2	A	401	SFU	O5-C1-C2	3.50	116.09	110.16
2	A	401	SFU	C5-O5-C1	5.31	119.87	112.78

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	402	SFU	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å <sup>2</sup> )	Q < 0.9
1	A	310/310 (100%)	0.48	27 (8%)	11 15	30, 37, 51, 67	0
1	B	310/310 (100%)	0.36	11 (3%)	44 51	31, 41, 56, 70	0
1	C	310/310 (100%)	0.35	10 (3%)	48 55	30, 41, 54, 70	0
All	All	930/930 (100%)	0.40	48 (5%)	28 35	30, 40, 55, 70	0

All (48) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	196	THR	4.1
1	C	195	HIS	3.9
1	A	162	GLY	3.8
1	B	296	LYS	3.7
1	C	266	ILE	3.4
1	A	266	ILE	3.3
1	A	264	SER	3.2
1	A	265	GLU	3.2
1	A	15	ILE	3.1
1	C	55	LYS	3.0
1	A	161	ILE	3.0
1	A	214	ALA	2.9
1	B	91	GLY	2.9
1	A	111	ARG	2.9
1	A	13	THR	2.8
1	B	50	LYS	2.8
1	A	215	ILE	2.8
1	A	263	GLY	2.7
1	A	160	GLY	2.6
1	B	224	LYS	2.6
1	A	16	ALA	2.6
1	A	14	GLY	2.5
1	A	267	ALA	2.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	161	ILE	2.4
1	B	70	LYS	2.4
1	A	112	ILE	2.4
1	C	194	PRO	2.3
1	A	12	ARG	2.3
1	A	61	PRO	2.3
1	A	62	LEU	2.3
1	B	163	VAL	2.3
1	A	121	ASN	2.2
1	C	16	ALA	2.2
1	C	200	LYS	2.2
1	A	144	ASP	2.2
1	B	15	ILE	2.1
1	A	158	GLY	2.1
1	C	248	GLY	2.1
1	B	127	ILE	2.1
1	B	305	ARG	2.1
1	A	213	CYS	2.1
1	A	59	GLY	2.0
1	C	265	GLU	2.0
1	C	249	GLN	2.0
1	B	113	GLY	2.0
1	A	216	ALA	2.0
1	A	179	TRP	2.0
1	A	17	ALA	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	SFU	B	405	12/12	0.87	0.32	6.11	37,45,47,87	0
2	SFU	C	403	12/12	0.93	0.25	6.02	37,45,50,70	0
3	SO4	A	408	5/5	0.94	0.42	5.84	52,52,52,53	0
2	SFU	B	404	12/12	0.63	0.41	5.11	38,45,49,86	0
2	SFU	C	404	12/12	0.86	0.30	4.91	40,48,51,82	0
2	SFU	C	405	12/12	0.84	0.23	4.73	36,44,52,72	0
2	SFU	A	404	12/12	0.73	0.29	4.69	37,44,51,83	0
2	SFU	A	405	12/12	0.83	0.22	4.38	37,45,52,76	0
2	SFU	C	401	12/12	0.91	0.28	3.89	37,45,50,79	0
2	SFU	B	401	12/12	0.88	0.33	2.42	37,44,47,77	0
2	SFU	B	403	12/12	0.90	0.21	1.51	35,43,46,68	0
2	SFU	C	402	12/12	0.90	0.17	1.08	34,41,44,49	0
2	SFU	A	403	12/12	0.89	0.17	0.48	37,46,52,65	0
2	SFU	A	401	12/12	0.93	0.15	0.46	37,46,49,64	0
2	SFU	A	402	12/12	0.96	0.14	-0.46	40,48,50,60	0
2	SFU	B	402	12/12	0.97	0.10	-1.81	35,42,47,51	0
3	SO4	C	406	5/5	0.97	0.23	-	49,49,51,51	0
3	SO4	A	406	5/5	0.96	0.18	-	49,49,53,53	0
3	SO4	A	407	5/5	0.94	0.32	-	51,51,52,53	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.