



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 02:44 pm GMT

PDB ID : 3EQP  
Title : Crystal Structure of Ack1 with compound T95  
Authors : Liu, J.; Wang, Z.; Walker, N.P.C.  
Deposited on : 2008-10-01  
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	4130 (2.30-2.30)
Clashscore	112137	4751 (2.30-2.30)
Ramachandran outliers	110173	4705 (2.30-2.30)
Sidechain outliers	110143	4704 (2.30-2.30)
RSRZ outliers	101464	4156 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	276	<div> <div>5%</div> <div> <div></div> <div>75%</div> <div>18%</div> <div>• •</div> </div> </div>
1	B	276	<div> <div>4%</div> <div> <div></div> <div>80%</div> <div>14%</div> <div>• •</div> </div> </div>

## 2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 4580 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Activated CDC42 kinase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	270	Total	C	N	O	S	0	0	0
			2168	1380	385	388	15			
1	A	268	Total	C	N	O	S	0	0	0
			2153	1372	383	383	15			

- Molecule 2 is N-(2,6-DIMETHYLPHENYL)-4-(2-ETHOXYPHENOXY)-2-({4-[4-(2-HYDROXYETHYL)PIPERAZIN-1-YL]PHENYL}AMINO)PYRIMIDINE-5-CARBOXAMIDE (three-letter code: T95) (formula: C<sub>33</sub>H<sub>38</sub>N<sub>6</sub>O<sub>4</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	B	1	Total	C	N	O	0	0
			43	33	6	4		
2	A	1	Total	C	N	O	0	0
			43	33	6	4		

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total 1	Cl 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	105	Total 105	O 105	0	0
4	A	67	Total 67	O 67	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Activated CDC42 kinase 1



#### • Molecule 1: Activated CDC42 kinase 1



THR  
ASP

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	71.00Å 43.11Å 92.85Å 90.00° 99.87° 90.00°	Depositor
Resolution (Å)	60.86 – 2.30 60.82 – 2.30	Depositor EDS
% Data completeness (in resolution range)	94.2 (60.86-2.30) 94.2 (60.82-2.30)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.11	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.42 (at 2.29Å)	Xtriage
Refinement program	REFMAC 5.5.0044	Depositor
R, $R_{free}$	0.221 , 0.284 0.223 , 0.283	Depositor DCC
$R_{free}$ test set	1223 reflections (5.46%)	DCC
Wilson B-factor (Å <sup>2</sup> )	22.7	Xtriage
Anisotropy	0.279	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 45.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	4580	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.70% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: T95, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.56	0/2205	0.74	2/2983 (0.1%)
1	B	0.57	0/2220	0.70	0/3004
All	All	0.56	0/4425	0.72	2/5987 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	389	GLN	C-N-CD	-8.72	101.42	120.60
1	A	389	GLN	C-N-CA	6.26	148.31	122.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	389	GLN	Peptide

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2153	0	2147	39	0
1	B	2168	0	2158	30	0
2	A	43	0	38	2	0
2	B	43	0	38	4	0
3	B	1	0	0	0	0
4	A	67	0	0	5	0
4	B	105	0	0	2	0
All	All	4580	0	4381	74	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (74) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:147:ALA:HB1	1:B:148:PRO:HA	1.32	1.10
1:A:147:ALA:HB1	1:A:148:PRO:HA	1.29	1.06
1:A:359:ASP:HB2	4:A:395:HOH:O	1.62	1.00
1:A:388:ALA:HB2	4:A:397:HOH:O	1.63	0.98
1:A:359:ASP:HB3	1:A:388:ALA:HB1	1.54	0.86
1:A:147:ALA:HB1	1:A:148:PRO:CA	2.10	0.82
1:B:147:ALA:HB1	1:B:148:PRO:CA	2.09	0.81
1:A:120:LEU:HD11	1:A:197:LEU:HD22	1.63	0.79
1:B:235:GLN:HE22	1:B:266:VAL:H	1.33	0.76
1:B:120:LEU:HD11	1:B:197:LEU:HD22	1.69	0.75
1:B:235:GLN:NE2	1:B:266:VAL:H	1.84	0.75
1:B:147:ALA:CB	1:B:148:PRO:HA	2.17	0.71
1:A:377:THR:HG21	4:A:444:HOH:O	1.93	0.67
1:B:186:HIS:HE1	1:B:188:ASN:HD22	1.44	0.65
1:A:147:ALA:CB	1:A:148:PRO:HA	2.17	0.64
1:A:195:VAL:HG13	1:A:197:LEU:HD13	1.81	0.63
1:A:255:ALA:HA	1:A:258:LEU:HD22	1.82	0.61
1:B:177:GLU:HG3	1:B:272:GLY:HA2	1.83	0.61
1:A:353:PRO:HB2	1:A:356:CYS:HB2	1.84	0.58
2:A:1:T95:N3	2:A:1:T95:H15	2.19	0.57
2:B:1:T95:H15	2:B:1:T95:N3	2.19	0.57
1:A:235:GLN:NE2	1:A:266:VAL:H	2.03	0.57
1:B:291:LYS:NZ	4:B:429:HOH:O	2.38	0.55
1:A:352:ARG:NH1	1:A:356:CYS:O	2.40	0.55

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:320:LEU:HD12	1:B:323:MET:HE3	1.90	0.54
1:B:235:GLN:HE22	1:B:265:LEU:HA	1.73	0.53
1:B:288:GLU:CD	1:B:288:GLU:H	2.12	0.53
1:A:288:GLU:H	1:A:288:GLU:CD	2.12	0.52
1:B:186:HIS:CE1	1:B:188:ASN:HD22	2.27	0.51
1:A:222:GLY:O	1:A:224:PHE:N	2.44	0.50
1:B:297:CYS:HB2	1:B:302:LEU:HD13	1.92	0.50
1:A:222:GLY:C	1:A:224:PHE:N	2.65	0.49
1:A:186:HIS:HE1	1:A:188:ASN:HD22	1.61	0.49
1:B:186:HIS:HB3	1:B:189:LEU:HD22	1.93	0.49
1:A:230:SER:HB3	1:A:389:GLN:HG2	1.94	0.49
1:B:230:SER:HB2	1:B:389:GLN:HG3	1.96	0.48
1:A:383:ASP:O	1:A:387:GLU:HG2	2.13	0.48
1:B:137:PHE:O	1:B:158:LYS:HE3	2.13	0.47
1:B:256:ARG:NH2	1:B:293:PRO:HG2	2.30	0.47
1:A:147:ALA:HA	1:A:150:GLY:H	1.80	0.47
1:B:177:GLU:OE2	2:B:1:T95:H31	2.15	0.47
1:A:389:GLN:OE1	4:A:447:HOH:O	2.20	0.47
1:B:298:ALA:HB1	1:B:300:GLU:OE2	2.15	0.46
1:B:288:GLU:HA	1:B:305:ARG:CZ	2.45	0.46
1:B:341:HIS:CD2	1:B:345:LYS:HG3	2.51	0.46
1:B:147:ALA:HA	1:B:150:GLY:H	1.81	0.45
1:A:222:GLY:O	1:A:223:HIS:C	2.53	0.45
2:B:1:T95:C15	2:B:1:T95:N3	2.80	0.45
1:A:284:TYR:OH	1:A:287:GLN:NE2	2.31	0.45
1:B:314:TRP:C	1:B:314:TRP:CD1	2.90	0.45
1:B:297:CYS:CB	1:B:302:LEU:HD13	2.48	0.44
1:A:120:LEU:HD12	1:A:120:LEU:HA	1.87	0.44
2:B:1:T95:H40A	4:B:398:HOH:O	2.19	0.43
1:A:235:GLN:HE22	1:A:266:VAL:H	1.66	0.43
1:A:302:LEU:HD12	1:A:302:LEU:HA	1.94	0.43
1:A:222:GLY:C	1:A:224:PHE:H	2.21	0.43
1:A:261:ALA:HB3	1:A:265:LEU:HD23	2.01	0.42
1:A:147:ALA:CB	1:A:148:PRO:CA	2.85	0.42
1:A:206:GLU:OE2	1:A:267:LYS:HE3	2.19	0.42
1:B:261:ALA:HB2	1:B:267:LYS:HD2	2.01	0.42
1:A:303:LYS:HB3	1:A:304:THR:HG23	2.00	0.42
1:A:377:THR:HG23	1:A:380:ALA:H	1.84	0.42
1:A:359:ASP:CB	4:A:395:HOH:O	2.41	0.42
1:A:232:TYR:O	1:A:236:VAL:HG23	2.20	0.42
2:A:1:T95:N3	2:A:1:T95:C15	2.81	0.41

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:319:THR:O	1:B:323:MET:HG3	2.21	0.41
1:A:308:SER:H	1:A:311:SER:HG	1.63	0.41
1:B:227:GLY:H	1:B:392:ASP:HB3	1.86	0.41
1:A:256:ARG:NH2	1:A:293:PRO:HG2	2.36	0.40
1:A:325:THR:HB	1:A:328:GLN:HG3	2.03	0.40
1:B:319:THR:HG22	1:B:323:MET:CE	2.51	0.40
1:A:137:PHE:O	1:A:158:LYS:HE3	2.22	0.40
1:A:377:THR:O	1:A:381:LEU:HD22	2.21	0.40
1:B:354:GLU:O	1:B:355:ASP:HB2	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	264/276 (96%)	252 (96%)	9 (3%)	3 (1%)	17	18
1	B	266/276 (96%)	255 (96%)	9 (3%)	2 (1%)	22	26
All	All	530/552 (96%)	507 (96%)	18 (3%)	5 (1%)	20	23

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	221	GLN
1	B	147	ALA
1	A	147	ALA
1	A	270	ASP
1	B	390	PRO

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	232/240 (97%)	210 (90%)	22 (10%)	10	12
1	B	234/240 (98%)	216 (92%)	18 (8%)	15	18
All	All	466/480 (97%)	426 (91%)	40 (9%)	12	14

All (40) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	160	LEU
1	B	189	LEU
1	B	191	ARG
1	B	197	LEU
1	B	258	LEU
1	B	265	LEU
1	B	279	GLN
1	B	281	ASP
1	B	283	HIS
1	B	288	GLU
1	B	292	VAL
1	B	300	GLU
1	B	302	LEU
1	B	320	LEU
1	B	354	GLU
1	B	359	ASP
1	B	371	LYS
1	B	382	ARG
1	A	132	LEU
1	A	151	LYS
1	A	160	LEU
1	A	183	SER
1	A	195	VAL
1	A	197	LEU
1	A	206	GLU
1	A	223	HIS
1	A	258	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	273	LEU
1	A	279	GLN
1	A	283	HIS
1	A	288	GLU
1	A	291	LYS
1	A	292	VAL
1	A	303	LYS
1	A	354	GLU
1	A	357	PRO
1	A	371	LYS
1	A	373	GLU
1	A	377	THR
1	A	381	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	188	ASN
1	B	221	GLN
1	B	235	GLN
1	B	257	ASN
1	A	182	HIS
1	A	188	ASN
1	A	235	GLN
1	A	287	GLN
1	A	389	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	T95	A	1	-	47,47,47	0.83	2 (4%)	60,64,64	2.01	11 (18%)
2	T95	B	1	-	47,47,47	0.92	2 (4%)	60,64,64	1.84	9 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	T95	A	1	-	-	0/26/36/36	0/5/5/5
2	T95	B	1	-	-	0/26/36/36	0/5/5/5

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1	T95	C9-N8	-3.34	1.36	1.43
2	B	1	T95	C9-N8	-2.98	1.37	1.43
2	A	1	T95	C4-N3	2.12	1.35	1.32
2	B	1	T95	C4-N3	3.47	1.37	1.32

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1	T95	N1-C2-N3	-8.03	118.95	126.68
2	B	1	T95	N1-C2-N3	-8.00	118.98	126.68
2	A	1	T95	C5-C4-N3	-4.74	119.76	124.44
2	B	1	T95	C5-C4-N3	-4.66	119.83	124.44
2	A	1	T95	C5-C6-N1	-4.36	118.71	124.41
2	A	1	T95	C36-C37-N38	-4.03	102.48	110.63
2	A	1	T95	C40-C39-N38	-3.32	103.92	110.63

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1	T95	C40-C39-N38	-2.73	105.11	110.63
2	B	1	T95	C36-C37-N38	-2.64	105.30	110.63
2	B	1	T95	C5-C6-N1	-2.60	121.00	124.41
2	A	1	T95	O26-C25-C24	-2.31	119.03	124.00
2	A	1	T95	C27-O26-C25	2.04	122.79	118.04
2	B	1	T95	C27-O26-C25	2.05	122.81	118.04
2	A	1	T95	O26-C25-C11	2.45	120.86	115.76
2	B	1	T95	C6-N1-C2	3.29	121.32	115.88
2	A	1	T95	C2-N3-C4	3.33	121.14	115.18
2	B	1	T95	C2-N3-C4	3.66	121.74	115.18
2	A	1	T95	C6-N1-C2	4.00	122.50	115.88
2	A	1	T95	C36-N20-C40	4.97	122.11	111.57
2	B	1	T95	C36-N20-C40	5.70	123.65	111.57

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1	T95	2	0
2	B	1	T95	4	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	268/276 (97%)	0.28	15 (5%) 25 32	10, 24, 50, 60	0
1	B	270/276 (97%)	0.20	11 (4%) 38 45	10, 24, 49, 60	0
All	All	538/552 (97%)	0.24	26 (4%) 31 38	10, 24, 50, 60	0

All (26) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	390	PRO	11.1
1	A	389	GLN	4.5
1	A	292	VAL	4.2
1	B	160	LEU	4.0
1	A	284	TYR	3.9
1	A	388	ALA	3.9
1	A	287	GLN	3.6
1	A	289	HIS	3.3
1	A	223	HIS	3.2
1	B	168	PRO	2.9
1	B	279	GLN	2.8
1	A	124	LYS	2.7
1	B	284	TYR	2.7
1	B	292	VAL	2.6
1	B	117	LEU	2.5
1	A	151	LYS	2.4
1	B	167	GLN	2.4
1	B	287	GLN	2.4
1	B	134	ASP	2.3
1	A	286	MET	2.2
1	B	288	GLU	2.2
1	A	231	ARG	2.2
1	A	294	PHE	2.1
1	B	169	GLU	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	152	THR	2.1
1	A	134	ASP	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	T95	A	1	43/43	0.94	0.14	0.14	18,21,36,41	0
2	T95	B	1	43/43	0.93	0.14	-0.21	17,22,36,39	0
3	CL	B	393	1/1	0.99	0.07	-3.04	16,16,16,16	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.