



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 16, 2017 – 03:10 PM EDT

PDB ID : 3EQV
Title : Crystal structure of penicillin-binding protein 2 from *Neisseria gonorrhoeae* containing four mutations associated with penicillin resistance
Authors : Powell, A.J.; Deacon, A.M.; Nicholas, R.A.; Davies, C.
Deposited on : unknown
Resolution : 2.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20030345
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20030345


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	3166 (2.40-2.40)
Clashscore	112137	3674 (2.40-2.40)
Ramachandran outliers	110173	3616 (2.40-2.40)
Sidechain outliers	110143	3617 (2.40-2.40)
RSRZ outliers	101464	3195 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	542	<div> <div>4%</div> <div>69%</div> <div>11%</div> <div>18%</div> </div>
1	B	542	<div> <div>6%</div> <div>68%</div> <div>9%</div> <div>21%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	SO4	B	6	-	-	-	X

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 6805 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Penicillin-binding protein 2.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	443	Total	C	N	O	P	S	0	1	0
			3395	2126	613	647	1	8			
1	B	426	Total	C	N	O	P	S	0	0	0
			3244	2034	583	618	1	8			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	40	GLY	-	EXPRESSION TAG	UNP P08149
A	41	SER	-	EXPRESSION TAG	UNP P08149
A	42	GLY	-	EXPRESSION TAG	UNP P08149
A	43	GLY	-	EXPRESSION TAG	UNP P08149
A	504	LEU	PHE	ENGINEERED	UNP P08149
A	510	VAL	ALA	ENGINEERED	UNP P08149
A	516	GLY	ALA	ENGINEERED	UNP P08149
A	551	SER	PRO	ENGINEERED	UNP P08149
B	40	GLY	-	EXPRESSION TAG	UNP P08149
B	41	SER	-	EXPRESSION TAG	UNP P08149
B	42	GLY	-	EXPRESSION TAG	UNP P08149
B	43	GLY	-	EXPRESSION TAG	UNP P08149
B	504	LEU	PHE	ENGINEERED	UNP P08149
B	510	VAL	ALA	ENGINEERED	UNP P08149
B	516	GLY	ALA	ENGINEERED	UNP P08149
B	551	SER	PRO	ENGINEERED	UNP P08149

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	40	Total	O	0	0
			40	40		
4	B	39	Total	O	0	0
			39	39		

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Chain A:

4% 69% 11% 18%

Amino Acid	Frequency (%)
GLY	4%
SER	69%
GLY	69%
GLY	69%
ALA	11%
ARG	1%
GLY	69%
LEU	69%
TYR	69%
GLN	69%
LEU	69%
THR	69%
VAL	15%
TS3	11%
K58	1%
G61	1%
D62	1%
T74	1%
D80	1%
L86	1%
S89	1%
T92	1%
E93	1%
SER	69%
LEU	69%
PHE	69%
ALA	11%
VAL	15%
PRO	1%
LYS	69%
ASP	69%
MET	69%
LYS	69%
GLU	69%
MET	69%
PRO	1%
SER	69%
ALA	11%
ALA	11%
GLN	69%
LEU	69%
GLU	69%
ARG	69%
LEU	69%
SER	69%
GLU	69%
VAL	15%
ASP	69%
VAL	15%
ASP	69%
LEU	69%
ARG	69%
GLN	69%
LEU	69%
ASP	69%
PRO	1%
LYS	69%
VAL	15%
ALA	11%
GLU	69%
GLU	69%
VAL	15%
LYS	69%
ALA	11%
LEU	69%
GLY	69%
LEU	69%
GLU	69%
GLU	69%
ASN	69%
PHE	69%
VAL	15%
PHE	69%
VAL	15%
GLU	69%
LYS	69%
E164	1%
L185	1%
K166	1%
R167	1%
M171	1%
G172	1%
N173	1%
L174	1%
T182	1%
D185	1%
E193	1%
L194	1%
L424	1%
Q425	1%
L426	1%
E203	1%
S427	1%
P213	1%

Chain B:

6% 68% 9% 21%

GLY SER GLY GLY ALA ARG GLY LEU TYR LEU GLN THR VAL THR VAL THR TYR ASN PHE LEU LYS GLU GLN GLY ASP ASN ARG ASN ILE V66 R67 T68 Q69 D80 P91 T92 E93 SER PHE LEU ALA VAL PRO LYS ASP MET LYS GLU MET PRO SER ALA ALA GLN LEU GLU ARG LEU SER GLU VAL

VAL ASP VAL PRO VAL ASP VAL LEU ARG ASN LYS LEU GLN THR VAL GLN LYS TYR LYS GLY ASN LYS SER PHE ILE TRP ILE LYS ARG LYS GLN ASP LEU ASP PRO LYS VAL GLU VAL LYS ALA LEU GLY LEU GLU ASN PHE VAL PHE ASP LYS GLU L465 M171 G172 N173 L174 D183 I184 D185 Q189

L194 A206 V208 V209 L210 R211 D212 ARG GLN LYS M216 L217 V218 D219 S220 L237 Q241 L246 E250 E256 R271 R298 V302 I306 G309 A318 R331 L332 R345 V349 Y350 K361 S362 S371 F374 E378 N379 Y380 V390

L403 W407 T417 G421 L424 L428 L438 L453 W454 R468 E469 V470 R471 M472 L473 M474 G487 D490 K497 T498 K503 LEU VAL ASN GLY ARG TYR VAL ASP M512 K513 G516 T517 F518 F521 P527 T534 E537 V547 V555

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	57.60Å 137.20Å 229.80Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.10 – 2.40 48.21 – 2.39	Depositor EDS
% Data completeness (in resolution range)	97.7 (48.10-2.40) 97.6 (48.21-2.39)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.23 (at 2.39Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.217 , 0.253 0.255 , 0.286	Depositor DCC
R_{free} test set	3528 reflections (4.98%)	DCC
Wilson B-factor (Å ²)	43.1	Xtriage
Anisotropy	0.797	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 38.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	6805	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.38% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, SO4, SEP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.58	1/3452 (0.0%)	0.71	0/4671
1	B	0.52	0/3293	0.65	1/4457 (0.0%)
All	All	0.55	1/6745 (0.0%)	0.68	1/9128 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	333	ASN	CG-ND2	9.17	1.55	1.32

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	332	LEU	CA-CB-CG	6.79	130.91	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3395	0	3392	35	0
1	B	3244	0	3252	26	0
2	A	30	0	0	0	0
2	B	45	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	6	0	8	0	0
3	B	6	0	8	0	0
4	A	40	0	0	0	0
4	B	39	0	0	0	0
All	All	6805	0	6660	61	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 61 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:92:THR:HB	1:B:93:GLU:HA	1.49	0.94
1:A:251:LEU:HG	1:A:265:VAL:HG23	1.49	0.94
1:A:86:LEU:CD2	1:A:171:MET:CE	2.65	0.74
1:A:86:LEU:CD2	1:A:171:MET:HE3	2.18	0.74
1:A:522:ALA:HB3	1:A:557:MET:HE2	1.69	0.73

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	437/542 (81%)	429 (98%)	7 (2%)	1 (0%)	51	67
1	B	417/542 (77%)	406 (97%)	11 (3%)	0	100	100
All	All	854/1084 (79%)	835 (98%)	18 (2%)	1 (0%)	55	72

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	480	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	356/438 (81%)	323 (91%)	33 (9%)	10	15
1	B	340/438 (78%)	317 (93%)	23 (7%)	18	29
All	All	696/876 (80%)	640 (92%)	56 (8%)	14	21

5 of 56 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	431	LEU
1	A	551	SER
1	B	390	VAL
1	A	460	ARG
1	A	473	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	425	GLN
1	B	562	ASN
1	B	173	ASN
1	A	329	ASN
1	B	69	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
1	SEP	A	465	1	9,9,10	1.82	2 (22%)	9,12,14	1.82	2 (22%)
1	SEP	B	465	1	9,9,10	1.88	2 (22%)	9,12,14	2.67	3 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	SEP	A	465	1	-	0/5/8/10	0/0/0/0
1	SEP	B	465	1	-	0/5/8/10	0/0/0/0

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	465	SEP	P-O1P	3.05	1.61	1.50
1	A	465	SEP	P-O1P	3.11	1.61	1.50
1	B	465	SEP	CA-C	3.41	1.54	1.50
1	A	465	SEP	CA-C	3.43	1.54	1.50

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	465	SEP	O3P-P-OG	2.09	112.30	106.73
1	A	465	SEP	OG-P-O1P	2.25	112.78	106.47
1	B	465	SEP	O2P-P-OG	2.27	112.78	106.73
1	A	465	SEP	OG-CB-CA	4.07	112.18	108.17
1	B	465	SEP	OG-CB-CA	6.82	114.90	108.17

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

17 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	A	11	-	4,4,4	0.17	0	6,6,6	0.33	0
2	SO4	A	13	-	4,4,4	0.16	0	6,6,6	0.14	0
2	SO4	A	15	-	4,4,4	0.18	0	6,6,6	0.18	0
3	GOL	A	2	-	5,5,5	0.39	0	5,5,5	0.37	0
2	SO4	A	3	-	4,4,4	0.17	0	6,6,6	0.15	0
2	SO4	A	4	-	4,4,4	0.14	0	6,6,6	0.39	0
2	SO4	A	8	-	4,4,4	0.13	0	6,6,6	0.08	0
2	SO4	B	1	-	4,4,4	0.21	0	6,6,6	0.17	0
2	SO4	B	10	-	4,4,4	0.17	0	6,6,6	0.08	0
2	SO4	B	12	-	4,4,4	0.18	0	6,6,6	0.25	0
2	SO4	B	14	-	4,4,4	0.20	0	6,6,6	0.35	0
2	SO4	B	2	-	4,4,4	0.13	0	6,6,6	0.23	0
2	SO4	B	5	-	4,4,4	0.11	0	6,6,6	0.33	0
3	GOL	B	582	-	5,5,5	0.41	0	5,5,5	0.34	0
2	SO4	B	6	-	4,4,4	0.15	0	6,6,6	0.16	0
2	SO4	B	7	-	4,4,4	0.17	0	6,6,6	0.12	0
2	SO4	B	9	-	4,4,4	0.09	0	6,6,6	0.83	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SO4	A	11	-	-	0/0/0/0	0/0/0/0
2	SO4	A	13	-	-	0/0/0/0	0/0/0/0
2	SO4	A	15	-	-	0/0/0/0	0/0/0/0
3	GOL	A	2	-	-	0/4/4/4	0/0/0/0
2	SO4	A	3	-	-	0/0/0/0	0/0/0/0
2	SO4	A	4	-	-	0/0/0/0	0/0/0/0
2	SO4	A	8	-	-	0/0/0/0	0/0/0/0
2	SO4	B	1	-	-	0/0/0/0	0/0/0/0
2	SO4	B	10	-	-	0/0/0/0	0/0/0/0
2	SO4	B	12	-	-	0/0/0/0	0/0/0/0
2	SO4	B	14	-	-	0/0/0/0	0/0/0/0
2	SO4	B	2	-	-	0/0/0/0	0/0/0/0
2	SO4	B	5	-	-	0/0/0/0	0/0/0/0
3	GOL	B	582	-	-	0/4/4/4	0/0/0/0
2	SO4	B	6	-	-	0/0/0/0	0/0/0/0
2	SO4	B	7	-	-	0/0/0/0	0/0/0/0
2	SO4	B	9	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	442/542 (81%)	0.38	21 (4%) 31 30	33, 46, 56, 65	0
1	B	425/542 (78%)	0.65	32 (7%) 15 13	33, 49, 61, 71	0
All	All	867/1084 (79%)	0.52	53 (6%) 22 20	33, 48, 60, 71	0

The worst 5 of 53 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	218	VAL	6.1
1	B	217	ILE	5.9
1	A	543	TYR	5.1
1	B	209	VAL	4.5
1	B	211	ARG	4.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	SEP	B	465	10/11	0.76	0.22	-	52,53,57,58	5
1	SEP	A	465	10/11	0.86	0.18	-	46,47,50,50	5

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	SO4	B	6	5/5	0.70	0.22	4.27	97,97,98,98	5
3	GOL	A	2	6/6	0.91	0.18	0.88	53,54,55,56	0
3	GOL	B	582	6/6	0.92	0.17	0.75	62,63,64,64	0
2	SO4	B	5	5/5	0.98	0.14	-0.25	63,65,66,66	0
2	SO4	B	1	5/5	0.96	0.13	-1.75	72,73,73,74	0
2	SO4	B	9	5/5	0.99	0.12	-1.94	42,45,49,49	0
2	SO4	B	12	5/5	0.67	0.43	-	59,59,60,61	5
2	SO4	B	7	5/5	0.86	0.26	-	62,62,63,63	5
2	SO4	A	13	5/5	0.90	0.13	-	65,65,65,66	5
2	SO4	A	3	5/5	0.88	0.18	-	64,65,66,66	5
2	SO4	A	8	5/5	0.94	0.27	-	64,64,65,65	5
2	SO4	B	14	5/5	0.88	0.27	-	42,42,42,43	5
2	SO4	A	4	5/5	0.89	0.16	-	74,75,75,76	5
2	SO4	B	2	5/5	0.97	0.17	-	60,61,61,61	5
2	SO4	A	11	5/5	0.88	0.19	-	61,64,64,65	5
2	SO4	B	10	5/5	0.75	0.15	-	73,73,73,73	5
2	SO4	A	15	5/5	0.83	0.20	-	59,60,60,61	5

6.5 Other polymers [i](#)

There are no such residues in this entry.