



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 14, 2017 – 04:19 pm GMT

PDB ID : 5EQA
Title : Crystal structure of *Medicago truncatula* Histidinol-Phosphate Phosphatase (MtHPP) with intermolecular cross-link between Lys158 and Cys245
Authors : Ruszkowski, M.; Dauter, Z.
Deposited on : 2015-11-12
Resolution : 1.32 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.32 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	1164 (1.34-1.30)
Clashscore	112137	1223 (1.34-1.30)
Ramachandran outliers	110173	1179 (1.34-1.30)
Sidechain outliers	110143	1179 (1.34-1.30)
RSRZ outliers	101464	1167 (1.34-1.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	277	<div> <div>4%</div> <div>88%</div> <div>8%</div> </div>
1	B	277	<div> <div>%</div> <div>85%</div> <div>6%</div> <div>9%</div> </div>
1	C	277	<div> <div>13%</div> <div>85%</div> <div>7%</div> <div>8%</div> </div>
1	D	277	<div> <div>9%</div> <div>84%</div> <div>7%</div> <div>8%</div> </div>

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 9078 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Inositol monophosphatase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	254	Total	C	N	O	S	0	7	0
			2003	1288	331	379	5			
1	B	253	Total	C	N	O	S	0	7	0
			1994	1281	329	379	5			
1	C	256	Total	C	N	O	S	0	4	0
			2000	1282	333	380	5			
1	D	254	Total	C	N	O	S	0	4	0
			1997	1278	334	380	5			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	50	SER	-	expression tag	UNP G7J7Q5
A	51	ASN	-	expression tag	UNP G7J7Q5
A	52	ALA	-	expression tag	UNP G7J7Q5
B	50	SER	-	expression tag	UNP G7J7Q5
B	51	ASN	-	expression tag	UNP G7J7Q5
B	52	ALA	-	expression tag	UNP G7J7Q5
C	50	SER	-	expression tag	UNP G7J7Q5
C	51	ASN	-	expression tag	UNP G7J7Q5
C	52	ALA	-	expression tag	UNP G7J7Q5
D	50	SER	-	expression tag	UNP G7J7Q5
D	51	ASN	-	expression tag	UNP G7J7Q5
D	52	ALA	-	expression tag	UNP G7J7Q5

- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	3	Total	Cl	0	0
			3	3		
2	A	3	Total	Cl	0	0
			3	3		


Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	D	2	Total	Cl	0	0
			2	2		
2	C	1	Total	Cl	0	0
			1	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	308	Total	O	0	1
			309	309		
3	B	312	Total	O	0	3
			315	315		
3	C	211	Total	O	0	2
			213	213		
3	D	238	Total	O	0	0
			238	238		



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	62.24Å 89.39Å 92.24Å 90.00° 96.93° 90.00°	Depositor
Resolution (Å)	36.20 – 1.32 36.21 – 1.32	Depositor EDS
% Data completeness (in resolution range)	99.9 (36.20-1.32) 99.9 (36.21-1.32)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.11 (at 1.32Å)	Xtriage
Refinement program	REFMAC 5.8.0131	Depositor
R, R_{free}	0.124 , 0.165 0.125 , 0.165	Depositor DCC
R_{free} test set	2342 reflections (1.01%)	DCC
Wilson B-factor (Å ²)	14.6	Xtriage
Anisotropy	0.679	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 48.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	9078	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.09% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MLZ, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.04	0/2055	0.98	0/2793
1	B	1.07	0/2046	1.02	0/2782
1	C	0.98	0/2046	1.00	4/2780 (0.1%)
1	D	1.00	0/2040	0.95	2/2771 (0.1%)
All	All	1.02	0/8187	0.99	6/11126 (0.1%)

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	244	GLY	N-CA-C	8.56	134.49	113.10
1	C	245	CYS	N-CA-C	8.39	133.66	111.00
1	D	245	CYS	N-CA-C	8.09	132.85	111.00
1	C	244	GLY	N-CA-C	7.96	133.00	113.10
1	C	114[A]	SER	N-CA-C	6.24	127.86	111.00
1	C	114[B]	SER	N-CA-C	6.24	127.86	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2003	0	1998	11	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	1994	0	1986	14	0
1	C	2000	0	1985	15	0
1	D	1997	0	1973	14	0
2	A	3	0	0	0	0
2	B	3	0	0	0	0
2	C	1	0	0	0	0
2	D	2	0	0	0	0
3	A	309	0	0	2	0
3	B	315	0	0	9	0
3	C	213	0	0	0	0
3	D	238	0	0	3	0
All	All	9078	0	7942	48	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (48) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:214:TYR:CE2	1:C:240:ILE:HD11	2.04	0.92
1:B:218[B]:THR:HG21	3:B:712[B]:HOH:O	1.86	0.74
1:A:284[B]:VAL:HG21	1:A:314:LYS:CD	2.25	0.66
1:B:129:GLU:HG2	3:B:724:HOH:O	2.00	0.62
1:B:191[B]:THR:HG21	3:B:575:HOH:O	2.00	0.60
1:C:214:TYR:CZ	1:C:240:ILE:HD11	2.36	0.59
1:A:284[B]:VAL:HG21	1:A:314:LYS:HD3	1.85	0.58
1:D:207:CYS:SG	3:D:684:HOH:O	2.56	0.58
1:B:207:CYS:O	1:B:315:GLN:NE2	2.36	0.57
1:B:314:LYS:HE3	3:B:767:HOH:O	2.04	0.57
1:D:191[B]:THR:HG22	3:D:508:HOH:O	2.05	0.56
1:B:126:TYR:HE1	3:B:551:HOH:O	1.88	0.55
1:A:284[B]:VAL:CG2	1:A:314:LYS:HD3	2.37	0.55
1:A:207:CYS:HB3	1:A:316:ILE:HG13	1.88	0.54
1:B:191[B]:THR:HG22	3:B:519:HOH:O	2.07	0.54
1:A:284[B]:VAL:CG2	1:A:314:LYS:CD	2.85	0.54
1:C:214:TYR:CE2	1:C:240:ILE:CD1	2.85	0.54
1:B:218[A]:THR:HG23	3:B:540:HOH:O	2.08	0.53
1:B:284[B]:VAL:HG21	1:B:314:LYS:HG3	1.90	0.53
1:B:99:SER:HB2	3:B:755:HOH:O	2.09	0.53
1:C:216:TYR:CE1	1:D:158:MLZ:HCM2	2.45	0.52
1:A:284[B]:VAL:HG21	1:A:314:LYS:HD2	1.91	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:239:LYS:HG2	1:D:240:ILE:HG12	1.93	0.51
1:C:239:LYS:HG2	1:C:240:ILE:HD13	1.94	0.49
1:A:214:TYR:CE1	1:A:240:ILE:HD12	2.49	0.48
1:C:152:LYS:HE3	1:D:240:ILE:CD1	2.44	0.48
1:D:128:GLU:N	3:D:503:HOH:O	2.47	0.47
1:A:284[B]:VAL:CG2	1:A:314:LYS:HD2	2.44	0.46
1:C:158:MLZ:HCM2	1:D:216:TYR:CE1	2.52	0.45
1:D:239:LYS:HG2	1:D:240:ILE:CG1	2.46	0.45
1:C:224:SER:O	1:C:228:GLU:HB2	2.18	0.43
1:D:263[B]:GLU:HG2	1:D:266:LEU:HG	2.00	0.43
1:D:284:VAL:HG21	1:D:314:LYS:HG2	2.00	0.43
1:A:130:LYS:HA	3:A:503:HOH:O	2.18	0.43
1:A:263[B]:GLU:HG3	3:A:636:HOH:O	2.18	0.43
1:B:205:ARG:NH2	1:B:214:TYR:CD2	2.77	0.43
1:C:311:ALA:HB3	1:C:317:HIS:HB2	2.00	0.43
1:D:275:ILE:HB	1:D:276:PRO:HD3	2.00	0.42
1:B:174:ILE:HA	1:B:191[B]:THR:CG2	2.50	0.42
1:C:152:LYS:HE3	1:D:240:ILE:HD13	2.00	0.42
1:A:183:LEU:HD22	1:B:183:LEU:HD22	2.00	0.41
1:B:236:ASP:CG	3:B:502:HOH:O	2.58	0.41
1:C:224:SER:O	1:C:228:GLU:N	2.46	0.41
1:C:214:TYR:CD2	1:C:240:ILE:HD11	2.54	0.41
1:C:183:LEU:HD22	1:D:183:LEU:HD22	2.03	0.41
1:D:210:LEU:O	1:D:238:VAL:HA	2.21	0.40
1:C:294:ARG:HH11	1:C:294:ARG:HG2	1.86	0.40
1:C:146:ASP:OD2	1:C:270:ASP:OD1	2.39	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	256/277 (92%)	250 (98%)	6 (2%)	0	100	100
1	B	255/277 (92%)	248 (97%)	7 (3%)	0	100	100
1	C	255/277 (92%)	249 (98%)	6 (2%)	0	100	100
1	D	253/277 (91%)	244 (96%)	8 (3%)	1 (0%)	38	11
All	All	1019/1108 (92%)	991 (97%)	27 (3%)	1 (0%)	55	20

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	225	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	217/232 (94%)	214 (99%)	3 (1%)	71	35
1	B	216/232 (93%)	213 (99%)	3 (1%)	71	35
1	C	216/232 (93%)	211 (98%)	5 (2%)	56	15
1	D	214/232 (92%)	212 (99%)	2 (1%)	82	53
All	All	863/928 (93%)	850 (98%)	13 (2%)	70	32

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	130	LYS
1	A	180	GLN
1	A	314	LYS
1	B	87	LYS
1	B	103	ILE
1	B	180	GLN
1	C	135	LYS
1	C	136	GLN
1	C	180	GLN
1	C	240	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	314	LYS
1	D	180	GLN
1	D	240	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	315	GLN
1	C	136	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	MLZ	A	158	1	9,9,10	1.35	2 (22%)	6,9,11	1.03	0
1	MLZ	B	158	1	9,9,10	1.37	2 (22%)	6,9,11	1.12	1 (16%)
1	MLZ	C	158	1	9,9,10	1.49	2 (22%)	6,9,11	0.80	0
1	MLZ	D	158	1	9,9,10	0.93	1 (11%)	6,9,11	1.08	1 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	MLZ	A	158	1	-	0/6/8/10	0/0/0/0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	MLZ	B	158	1	-	0/6/8/10	0/0/0/0
1	MLZ	C	158	1	-	0/6/8/10	0/0/0/0
1	MLZ	D	158	1	-	0/6/8/10	0/0/0/0

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	158	MLZ	CB-CA	-2.64	1.50	1.53
1	C	158	MLZ	CB-CA	-2.25	1.50	1.53
1	B	158	MLZ	CB-CA	-2.24	1.50	1.53
1	D	158	MLZ	CA-C	2.37	1.53	1.50
1	A	158	MLZ	CA-C	2.56	1.53	1.50
1	B	158	MLZ	CA-C	2.63	1.53	1.50
1	C	158	MLZ	CA-C	2.66	1.53	1.50

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	158	MLZ	O-C-CA	-2.59	117.85	125.02
1	D	158	MLZ	O-C-CA	-2.07	119.31	125.02

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	C	158	MLZ	1	0
1	D	158	MLZ	1	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 9 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	253/277 (91%)	0.20	11 (4%) 36 36	10, 17, 37, 60	0
1	B	252/277 (90%)	0.14	4 (1%) 72 73	10, 16, 35, 50	0
1	C	255/277 (92%)	0.69	35 (13%) 3 4	13, 23, 48, 78	0
1	D	253/277 (91%)	0.38	24 (9%) 9 8	11, 20, 42, 59	0
All	All	1013/1108 (91%)	0.35	74 (7%) 16 15	10, 19, 41, 78	0

All (74) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	223	PHE	7.8
1	C	224	SER	7.7
1	C	63	LEU	7.6
1	C	98	LEU	7.0
1	D	223	PHE	6.4
1	B	208	ALA	6.0
1	A	208	ALA	5.7
1	A	63	LEU	5.6
1	D	222	LEU	5.5
1	C	64	ASN	5.3
1	C	135	LYS	5.1
1	C	208	ALA	5.1
1	C	221	HIS	5.0
1	C	137	ASP	5.0
1	C	222	LEU	5.0
1	C	218	THR	4.6
1	D	63	LEU	4.5
1	D	224	SER	4.3
1	C	132	TRP	4.2
1	D	221	HIS	4.1
1	B	99	SER	3.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	226	ASP	3.9
1	B	100	PRO	3.9
1	D	99	SER	3.9
1	D	208	ALA	3.8
1	C	231	PHE	3.7
1	D	225	GLY	3.7
1	D	64	ASN	3.7
1	C	88	ASN	3.6
1	A	64	ASN	3.6
1	C	299	PRO	3.6
1	D	220	PRO	3.6
1	C	225	GLY	3.6
1	C	226	ASP	3.5
1	B	64	ASN	3.5
1	D	326	TRP	3.4
1	D	302	ILE	3.4
1	C	62	GLN	3.3
1	C	240	ILE	3.3
1	C	227	ALA	3.2
1	D	244	GLY	3.2
1	C	220	PRO	3.1
1	A	100	PRO	3.1
1	D	243	TYR	3.1
1	C	100	PRO	3.0
1	C	129	GLU	3.0
1	A	99	SER	2.9
1	C	136	GLN	2.8
1	C	217	THR	2.7
1	D	100	PRO	2.7
1	D	137	ASP	2.6
1	D	129	GLU	2.6
1	C	103	ILE	2.5
1	C	99	SER	2.5
1	D	87	LYS	2.4
1	C	244	GLY	2.4
1	C	134	CYS	2.4
1	A	137	ASP	2.3
1	D	299	PRO	2.3
1	A	271	PHE	2.3
1	C	300	LEU	2.3
1	C	65	HIS	2.3
1	A	62	GLN	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	84	TYR	2.1
1	C	212	GLN	2.1
1	D	62	GLN	2.1
1	A	274	LEU	2.1
1	C	326	TRP	2.1
1	D	325	GLN	2.1
1	C	228	GLU	2.1
1	D	229	GLU	2.1
1	D	219	SER	2.1
1	A	207	CYS	2.1
1	A	178	ILE	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	MLZ	C	158	10/11	0.96	0.10	-	16,17,21,21	0
1	MLZ	B	158	10/11	0.96	0.11	-	10,12,12,13	0
1	MLZ	D	158	10/11	0.96	0.10	-	14,16,21,23	0
1	MLZ	A	158	10/11	0.97	0.11	-	11,12,12,13	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	CL	B	403	1/1	1.00	0.06	-0.96	22,22,22,22	0
2	CL	A	403	1/1	1.00	0.04	-1.91	20,20,20,20	0
2	CL	A	402	1/1	0.99	0.07	-	29,29,29,29	0
2	CL	A	401	1/1	1.00	0.04	-	32,32,32,32	0
2	CL	D	402	1/1	0.99	0.03	-	33,33,33,33	0
2	CL	C	401	1/1	0.99	0.04	-	31,31,31,31	0
2	CL	B	401	1/1	1.00	0.06	-	26,26,26,26	0
2	CL	D	401	1/1	1.00	0.07	-	29,29,29,29	0
2	CL	B	402	1/1	1.00	0.09	-	29,29,29,29	0

6.5 Other polymers [i](#)

There are no such residues in this entry.