



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 5, 2017 – 07:47 AM EST

PDB ID : 4EUU
Title : Structure of BX-795 Complexed with Human TBK1 Kinase Domain Phosphorylated on Ser172
Authors : Ma, X.; Helgason, E.; Phung, Q.T.; Quan, C.L.; Iyer, R.S.; Lee, M.W.; Bowman, K.K.; Starovasnik, M.A.; Dueber, E.C.
Deposited on : unknown
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20030345
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20030345


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	4827 (1.80-1.80)
Clashscore	112137	5742 (1.80-1.80)
Ramachandran outliers	110173	5676 (1.80-1.80)
Sidechain outliers	110143	5675 (1.80-1.80)
RSRZ outliers	101464	4906 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	319	<div> <div>5%</div> <div>87%</div> <div>9%</div> <div>.</div> </div>
1	B	319	<div> <div>5%</div> <div>88%</div> <div>8%</div> <div>.</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	IOD	A	402[B]	-	-	X	X
3	IOD	B	402[B]	-	-	X	-
4	SO4	A	403	-	-	X	-
5	GOL	A	405	-	-	-	X
5	GOL	A	406	-	-	-	X
5	GOL	B	404	-	-	-	X
5	GOL	B	405	-	-	-	X
5	GOL	B	406	-	-	X	X

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 5576 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Serine/threonine-protein kinase TBK1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	308	Total	C	N	O	P	S	0	0	0
			2447	1555	422	458	1	11			
1	B	308	Total	C	N	O	P	S	0	0	0
			2447	1555	422	458	1	11			

There are 26 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	MET	-	EXPRESSION TAG	UNP Q9UHD2
A	0	GLY	-	EXPRESSION TAG	UNP Q9UHD2
A	1	SER	-	EXPRESSION TAG	UNP Q9UHD2
A	135	ASN	ASP	ENGINEERED MUTATION	UNP Q9UHD2
A	309	GLY	-	EXPRESSION TAG	UNP Q9UHD2
A	310	ASN	-	EXPRESSION TAG	UNP Q9UHD2
A	311	SER	-	EXPRESSION TAG	UNP Q9UHD2
A	312	HIS	-	EXPRESSION TAG	UNP Q9UHD2
A	313	HIS	-	EXPRESSION TAG	UNP Q9UHD2
A	314	HIS	-	EXPRESSION TAG	UNP Q9UHD2
A	315	HIS	-	EXPRESSION TAG	UNP Q9UHD2
A	316	HIS	-	EXPRESSION TAG	UNP Q9UHD2
A	317	HIS	-	EXPRESSION TAG	UNP Q9UHD2
B	-1	MET	-	EXPRESSION TAG	UNP Q9UHD2
B	0	GLY	-	EXPRESSION TAG	UNP Q9UHD2
B	1	SER	-	EXPRESSION TAG	UNP Q9UHD2
B	135	ASN	ASP	ENGINEERED MUTATION	UNP Q9UHD2
B	309	GLY	-	EXPRESSION TAG	UNP Q9UHD2
B	310	ASN	-	EXPRESSION TAG	UNP Q9UHD2
B	311	SER	-	EXPRESSION TAG	UNP Q9UHD2
B	312	HIS	-	EXPRESSION TAG	UNP Q9UHD2
B	313	HIS	-	EXPRESSION TAG	UNP Q9UHD2
B	314	HIS	-	EXPRESSION TAG	UNP Q9UHD2
B	315	HIS	-	EXPRESSION TAG	UNP Q9UHD2
B	316	HIS	-	EXPRESSION TAG	UNP Q9UHD2

Continued on next page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	317	HIS	-	EXPRESSION TAG	UNP Q9UHD2

-
- Chemical structure of BX7, a pyridine derivative. The structure features a central pyridine ring (N11, N10) substituted with a 4-iodophenyl group (C28, C29, C30, I01), a 4-(2-((4-oxo-2-((4-iodophenyl)amino)ethyl)amino)butyl)phenyl group (C18, C17, C22, C25, C26, C27, C28, C29, C30, I01), and a 4-oxo-2-((4-iodophenyl)amino)ethyl group (C16, C17, C22, C25, C26, C27, C28, C29, C30, I01). The structure is labeled with atom IDs: N11, N10, N08, N06, N05, N07, N09, O03, O04, C13, C12, C15, C14, C18, C17, C22, C25, C26, C27, C28, C29, C30, I01, C19, C20, C21, C23, C31, C32, C33, C34, S02.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total 34	C 23	I 1	N 7	O 2	S 1	0	1
2	B	1	Total 34	C 23	I 1	N 7	O 2	S 1	0	1


- | Mol | Chain | Residues | Atoms | ZeroOcc | AltConf |
|-----|-------|----------|----------------|---------|---------|
| 3 | B | 1 | Total I
1 1 | 0 | 1 |
| 3 | A | 1 | Total I
1 1 | 0 | 1 |

- 



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total 6	C 3	O 3	0	0
5	A	1	Total 6	C 3	O 3	0	0
5	B	1	Total 6	C 3	O 3	0	0
5	B	1	Total 6	C 3	O 3	0	0
5	B	1	Total 6	C 3	O 3	0	0
5	B	1	Total 6	C 3	O 3	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	258	Total 258	O 258	0	0
6	B	297	Total 297	O 297	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Serine/threonine-protein kinase TBK1



- Molecule 1: Serine/threonine-protein kinase TBK1



4 Data and refinement statistics

Property	Value	Source
Space group	P 31	Depositor
Cell constants a, b, c, α , β , γ	76.09 Å 76.09 Å 130.94 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.00 – 1.80 32.95 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.8 (50.00-1.80) 99.8 (32.95-1.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.41 (at 1.81 Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.169 , 0.194 0.167 , 0.195	Depositor DCC
R_{free} test set	3949 reflections (5.30%)	DCC
Wilson B-factor (Å ²)	30.5	Xtriage
Anisotropy	0.490	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 50.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.011 for -h,-k,l 0.027 for h,-h-k,-l 0.016 for -k,-h,-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	5576	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.98% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, IOD, SO4, BX7, SEP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.65	0/2491	0.65	0/3366
1	B	0.66	0/2491	0.67	0/3366
All	All	0.66	0/4982	0.66	0/6732

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2447	0	2417	32	0
1	B	2447	0	2417	24	0
2	A	34	0	0	0	0
2	B	34	0	0	1	0
3	A	1	0	0	3	0
3	B	1	0	0	3	0
4	A	10	0	0	2	0
4	B	5	0	0	0	0
5	A	18	0	24	1	0
5	B	24	0	32	9	0
6	A	258	0	0	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	B	297	0	0	12	1
All	All	5576	0	4890	62	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (62) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:18:GLY:O	6:B:598:HOH:O	1.70	1.10
1:B:177:GLU:HG3	6:B:691:HOH:O	1.73	0.87
1:A:11:LEU:HB2	5:B:406:GOL:H2	1.54	0.87
1:B:186:GLU:OE2	1:B:235:TYR:OH	1.94	0.86
5:B:404:GOL:H12	6:B:673:HOH:O	1.75	0.86
1:A:232:GLU:OE1	1:B:191:ARG:NH1	2.10	0.83
5:B:407:GOL:H11	6:B:564:HOH:O	1.79	0.83
1:A:190:LEU:HD11	1:A:235:TYR:HB2	1.71	0.73
1:B:0:GLY:N	6:B:720:HOH:O	2.21	0.73
1:A:191:ARG:NH2	4:A:403:SO4:O1	2.23	0.71
1:A:190:LEU:HD11	1:A:235:TYR:CB	2.21	0.71
1:A:41:ASN:HB3	6:B:641:HOH:O	1.90	0.70
3:B:402[B]:IOD:I	6:B:519:HOH:O	2.79	0.69
1:B:294:GLY:HA3	5:B:404:GOL:H11	1.75	0.69
1:B:41:ASN:HB3	6:B:748:HOH:O	1.94	0.68
1:A:11:LEU:CB	5:B:406:GOL:H2	2.26	0.66
1:B:157:ASP:HA	3:B:402[B]:IOD:I	2.66	0.66
1:A:191:ARG:NH2	4:A:403:SO4:S	2.70	0.64
5:B:404:GOL:C1	6:B:673:HOH:O	2.38	0.64
1:A:0:GLY:N	6:A:641:HOH:O	2.30	0.63
1:B:86:MET:HE1	2:B:401[A]:BX7:I01	2.69	0.63
1:B:25:ARG:NH2	6:B:563:HOH:O	2.35	0.59
1:A:127:ARG:NH2	1:A:166:ASP:HB3	2.18	0.58
1:A:157:ASP:HA	3:A:402[B]:IOD:I	2.74	0.58
1:A:190:LEU:HD23	1:B:191:ARG:NH2	2.20	0.56
1:A:190:LEU:CD1	1:A:235:TYR:CG	2.88	0.56
1:B:190:LEU:HD13	1:B:192:LYS:HE2	1.87	0.56
1:B:278:THR:OG1	1:B:279:PRO:HD3	2.06	0.55
1:B:235:TYR:CZ	1:B:239:THR:HG21	2.42	0.54
1:A:258:ASP:OD1	6:A:706:HOH:O	2.19	0.52
1:A:53:MET:SD	1:A:82:LYS:NZ	2.82	0.52
1:A:58:VAL:HG22	1:A:131:ILE:HD12	1.90	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:190:LEU:CD1	1:A:235:TYR:HB2	2.40	0.48
1:B:187:ARG:O	1:B:191:ARG:HA	2.15	0.47
1:A:193:ASP:C	1:A:193:ASP:OD1	2.53	0.47
1:A:235:TYR:CE1	1:A:239:THR:HG21	2.50	0.47
1:A:190:LEU:CD1	1:A:235:TYR:CB	2.93	0.46
1:A:270:SER:O	1:A:274:GLN:HG3	2.17	0.45
1:A:43:ILE:CD1	1:A:46:LEU:HD12	2.47	0.45
1:A:1:SER:OG	1:A:10:LEU:HD12	2.16	0.45
1:B:229:ARG:HD3	1:B:229:ARG:HA	1.83	0.43
1:A:11:LEU:HD12	5:B:406:GOL:H31	2.00	0.43
1:A:43:ILE:HD11	1:A:46:LEU:HD12	2.00	0.43
1:B:213:HIS:HE1	6:B:722:HOH:O	2.02	0.43
1:A:86:MET:HE1	3:A:402[B]:IOD:I	2.90	0.42
1:B:225:GLU:OE1	1:B:225:GLU:N	2.50	0.42
1:B:253:GLU:HG2	1:B:254:ASN:OD1	2.20	0.42
1:B:92:GLY:HA3	5:B:406:GOL:H12	2.03	0.41
1:A:191:ARG:HG2	1:A:191:ARG:HH11	1.84	0.41
1:B:193:ASP:OD1	1:B:193:ASP:C	2.58	0.41
1:A:271:ARG:HH21	5:A:407:GOL:H32	1.86	0.41
1:B:41:ASN:ND2	6:B:647:HOH:O	2.53	0.41
1:A:280:VAL:O	1:A:284:ILE:HG12	2.21	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:B:787:HOH:O	6:B:790:HOH:O[2_555]	1.85	0.35

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	305/319 (96%)	299 (98%)	6 (2%)	0	100	100
1	B	305/319 (96%)	300 (98%)	5 (2%)	0	100	100
All	All	610/638 (96%)	599 (98%)	11 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	265/275 (96%)	264 (100%)	1 (0%)	93	91
1	B	265/275 (96%)	265 (100%)	0	100	100
All	All	530/550 (96%)	529 (100%)	1 (0%)	94	93

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	61	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	SEP	A	172	1	9,9,10	1.90	2 (22%)	9,12,14	1.01	0
1	SEP	B	172	1	9,9,10	1.77	2 (22%)	9,12,14	1.13	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	SEP	A	172	1	-	0/5/8/10	0/0/0/0
1	SEP	B	172	1	-	0/5/8/10	0/0/0/0

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	172	SEP	P-O1P	2.90	1.60	1.50
1	A	172	SEP	P-O1P	2.91	1.60	1.50
1	B	172	SEP	CA-C	3.67	1.55	1.50
1	A	172	SEP	CA-C	4.18	1.55	1.50

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 14 ligands modelled in this entry, 2 are monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	BX7	A	401[A]	-	35,37,37	2.74	11 (31%)	42,49,49	2.21	13 (30%)
4	SO4	A	403	-	4,4,4	0.28	0	6,6,6	0.32	0
4	SO4	A	404	-	4,4,4	0.20	0	6,6,6	0.16	0
5	GOL	A	405	-	5,5,5	0.23	0	5,5,5	0.42	0
5	GOL	A	406	-	5,5,5	0.38	0	5,5,5	0.81	0
5	GOL	A	407	-	5,5,5	0.28	0	5,5,5	0.40	0
2	BX7	B	401[A]	-	35,37,37	2.82	9 (25%)	42,49,49	2.60	13 (30%)
4	SO4	B	403	-	4,4,4	0.26	0	6,6,6	0.37	0
5	GOL	B	404	-	5,5,5	0.81	0	5,5,5	0.85	0
5	GOL	B	405	-	5,5,5	0.54	0	5,5,5	1.05	0
5	GOL	B	406	-	5,5,5	0.47	0	5,5,5	1.05	0
5	GOL	B	407	-	5,5,5	0.41	0	5,5,5	0.31	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BX7	A	401[A]	-	-	0/22/31/31	0/4/4/4
4	SO4	A	403	-	-	0/0/0/0	0/0/0/0
4	SO4	A	404	-	-	0/0/0/0	0/0/0/0
5	GOL	A	405	-	-	0/4/4/4	0/0/0/0
5	GOL	A	406	-	-	0/4/4/4	0/0/0/0
5	GOL	A	407	-	-	0/4/4/4	0/0/0/0
2	BX7	B	401[A]	-	-	0/22/31/31	0/4/4/4
4	SO4	B	403	-	-	0/0/0/0	0/0/0/0
5	GOL	B	404	-	-	0/4/4/4	0/0/0/0
5	GOL	B	405	-	-	0/4/4/4	0/0/0/0
5	GOL	B	406	-	-	0/4/4/4	0/0/0/0
5	GOL	B	407	-	-	0/4/4/4	0/0/0/0

All (20) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	401[A]	BX7	C31-S02	-8.05	1.62	1.72
2	A	401[A]	BX7	C31-S02	-6.78	1.64	1.72
2	B	401[A]	BX7	C14-N05	-3.71	1.38	1.47
2	A	401[A]	BX7	C15-N05	-3.42	1.39	1.47
2	B	401[A]	BX7	C15-N05	-3.33	1.39	1.47
2	A	401[A]	BX7	C14-N05	-3.22	1.39	1.47
2	A	401[A]	BX7	C27-N11	-2.23	1.31	1.34
2	B	401[A]	BX7	C30-C28	2.41	1.43	1.39
2	A	401[A]	BX7	C27-N08	2.55	1.41	1.36
2	A	401[A]	BX7	C16-N05	2.59	1.41	1.36
2	B	401[A]	BX7	C27-N08	3.06	1.42	1.36
2	A	401[A]	BX7	C30-C28	3.59	1.44	1.39
2	B	401[A]	BX7	C29-N09	3.94	1.42	1.33
2	A	401[A]	BX7	C29-N09	3.99	1.42	1.33
2	A	401[A]	BX7	C33-C32	4.44	1.54	1.39
2	B	401[A]	BX7	C33-C32	4.83	1.55	1.39
2	A	401[A]	BX7	C16-N06	5.05	1.44	1.36
2	B	401[A]	BX7	C16-N06	5.63	1.45	1.36
2	B	401[A]	BX7	C26-N07	8.42	1.47	1.34
2	A	401[A]	BX7	C26-N07	9.16	1.48	1.34

All (26) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	401[A]	BX7	C28-C30-N11	-10.84	114.41	123.15
2	A	401[A]	BX7	N11-C27-N10	-6.02	120.89	126.68
2	A	401[A]	BX7	C28-C30-N11	-5.12	119.02	123.15
2	B	401[A]	BX7	C24-C20-C17	-4.75	113.97	119.72
2	B	401[A]	BX7	C30-C28-I01	-3.84	114.10	118.94
2	A	401[A]	BX7	C23-N09-C29	-3.57	113.96	122.11
2	B	401[A]	BX7	N07-C26-N10	-3.02	115.01	118.65
2	A	401[A]	BX7	C21-N07-C26	-2.73	120.00	123.35
2	A	401[A]	BX7	C19-C23-N09	-2.46	105.09	112.18
2	B	401[A]	BX7	C23-N09-C29	-2.45	116.51	122.11
2	A	401[A]	BX7	C17-N06-C16	-2.38	121.45	125.80
2	A	401[A]	BX7	O03-C16-N06	-2.31	119.59	124.24
2	A	401[A]	BX7	C28-C26-N07	-2.24	117.85	120.80
2	B	401[A]	BX7	C19-C23-N09	-2.16	105.97	112.18
2	A	401[A]	BX7	C19-C21-N07	-2.05	105.99	111.46
2	B	401[A]	BX7	C18-C22-N08	2.14	127.21	120.36
2	B	401[A]	BX7	C20-C17-C18	2.81	122.97	119.66
2	B	401[A]	BX7	N06-C16-N05	3.03	119.31	115.91
2	B	401[A]	BX7	C20-C24-C25	3.23	124.77	120.24

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	401[A]	BX7	C30-N11-C27	3.44	121.58	115.88
2	A	401[A]	BX7	C20-C17-C18	3.46	123.73	119.66
2	B	401[A]	BX7	C26-C28-I01	3.79	125.03	120.94
2	A	401[A]	BX7	C26-C28-I01	4.31	125.59	120.94
2	A	401[A]	BX7	C30-N11-C27	4.32	123.03	115.88
2	A	401[A]	BX7	C14-N05-C15	4.51	118.04	111.36
2	B	401[A]	BX7	C14-N05-C15	4.60	118.18	111.36

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6 monomers are involved in 13 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	403	SO4	2	0
5	A	407	GOL	1	0
2	B	401[A]	BX7	1	0
5	B	404	GOL	3	0
5	B	406	GOL	5	0
5	B	407	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	307/319 (96%)	0.03	15 (4%) 30 25	25, 42, 70, 91	0
1	B	307/319 (96%)	-0.02	15 (4%) 30 25	23, 37, 68, 86	0
All	All	614/638 (96%)	0.00	30 (4%) 30 25	23, 40, 70, 91	0

All (30) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	194	HIS	5.0
1	A	195	GLN	4.5
1	B	195	GLN	4.3
1	A	290	GLU	4.2
1	A	147	GLU	4.0
1	A	194	HIS	3.9
1	B	193	ASP	3.2
1	A	193	ASP	3.0
1	B	147	GLU	3.0
1	B	254	ASN	2.9
1	A	289	GLN	2.9
1	B	240	GLY	2.8
1	A	167	ASP	2.8
1	B	192	LYS	2.8
1	A	307	HIS	2.7
1	B	207	ILE	2.6
1	B	232	GLU	2.6
1	A	192	LYS	2.5
1	B	167	ASP	2.5
1	A	50	ASP	2.4
1	B	252	ALA	2.4
1	B	251	LYS	2.4
1	B	148	ASP	2.3
1	A	166	ASP	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	304	ASP	2.3
1	B	136	ILE	2.3
1	A	19	ALA	2.2
1	B	224	PHE	2.1
1	A	191	ARG	2.1
1	A	179	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	SEP	A	172	10/11	0.99	0.07	-	45,47,52,52	0
1	SEP	B	172	10/11	0.99	0.08	-	38,40,42,46	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
5	GOL	A	405	6/6	0.75	0.23	7.75	57,58,60,64	0
5	GOL	B	405	6/6	0.93	0.20	4.96	33,50,53,56	0
3	IOD	A	402[B]	1/1	0.94	0.27	4.95	38,38,38,38	1
5	GOL	A	406	6/6	0.89	0.20	4.72	41,57,59,64	0
5	GOL	B	406	6/6	0.78	0.18	4.69	40,47,53,54	0
5	GOL	B	404	6/6	0.88	0.18	3.92	42,47,48,52	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	IOD	B	402[B]	1/1	0.93	0.14	1.70	43,43,43,43	1
4	SO4	A	403	5/5	0.83	0.20	0.48	78,84,89,92	0
2	BX7	A	401[A]	34/34	0.99	0.11	0.34	20,26,45,46	1
4	SO4	B	403	5/5	0.97	0.13	0.29	58,59,59,65	0
2	BX7	B	401[A]	34/34	0.99	0.10	0.00	17,23,45,46	1
4	SO4	A	404	5/5	0.86	0.18	-	87,89,91,92	0
5	GOL	A	407	6/6	0.84	0.23	-	66,70,75,79	0
5	GOL	B	407	6/6	0.77	0.16	-	56,58,62,66	0

6.5 Other polymers

There are no such residues in this entry.