



Full wwPDB NMR Structure Validation Report ⓘ

Feb 12, 2017 – 06:00 pm GMT

PDB ID : 1EVO
Title : NMR OBSERVATION OF A NOVEL C-TETRAD
Authors : Patel, P.K.; Bhavesh, N.S.; Hosur, R.V.
Deposited on : 2000-04-20

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/NMRValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)
NmrClust : Kelley et al. (1996)
MolProbity : 4.02b-467
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : trunk28760
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR





The overall completeness of chemical shifts assignment is 35%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	125131	11601

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	O	8	
1	P	8	
1	Q	8	
1	R	8	

2 Ensemble composition and analysis ⓘ

This entry contains 10 models. This entry does not contain polypeptide chains, therefore identification of well-defined residues and clustering analysis are not possible. All residues are included in the validation scores.

3 Entry composition [i](#)

There is only 1 type of molecule in this entry. The entry contains 1032 atoms, of which 368 are hydrogens and 0 are deuteriums.

- Molecule 1 is a DNA chain called DNA (5'-D(*TP*GP*GP*GP*CP*GP*GP*T)-3').

Mol	Chain	Residues	Atoms							Trace
1	O	8	Total	C	H	N	O	P		0
			258	79	92	32	48	7		
1	P	8	Total	C	H	N	O	P		0
			258	79	92	32	48	7		
1	Q	8	Total	C	H	N	O	P		0
			258	79	92	32	48	7		
1	R	8	Total	C	H	N	O	P		0
			258	79	92	32	48	7		


4 Residue-property plots [i](#)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: DNA (5'-D(*TP*GP*GP*GP*CP*GP*GP*T)-3')

Chain O: 



- Molecule 1: DNA (5'-D(*TP*GP*GP*GP*CP*GP*GP*T)-3')

Chain P: 



- Molecule 1: DNA (5'-D(*TP*GP*GP*GP*CP*GP*GP*T)-3')

Chain Q: 



- Molecule 1: DNA (5'-D(*TP*GP*GP*GP*CP*GP*GP*T)-3')

Chain R: 



4.2 Scores per residue for each member of the ensemble

Colouring as in section [4.1](#) above.

4.2.1 Score per residue for model 1

- Molecule 1: DNA (5'-D(*TP*GP*GP*GP*CP*GP*GP*T)-3')

Chain O:  25% 75%



- Molecule 1: DNA (5'-D(*TP*GP*GP*GP*CP*GP*GP*T)-3')

Chain P:  25% 75%



- Molecule 1: DNA (5'-D(*TP*GP*GP*GP*CP*GP*GP*T)-3')

Chain Q:  25% 75%



- Molecule 1: DNA (5'-D(*TP*GP*GP*GP*CP*GP*GP*T)-3')

Chain R:  25% 75%



4.2.2 Score per residue for model 2

- Molecule 1: DNA (5'-D(*TP*GP*GP*GP*CP*GP*GP*T)-3')

Chain O:  38% 63%




- Molecule 1: DNA (5'-D(*TP*GP*GP*GP*CP*GP*GP*T)-3')

Chain P:  25% 75%




- Molecule 1: DNA (5'-D(*TP*GP*GP*GP*CP*GP*GP*T)-3')

Chain Q:  38% 63%



- Molecule 1: DNA (5'-D(*TP*GP*GP*GP*CP*GP*GP*T)-3')

Chain R: 38% 63%



4.2.3 Score per residue for model 3

- Molecule 1: DNA (5'-D(*TP*GP*GP*GP*CP*GP*GP*T)-3')

Chain O: 38% 63%



- Molecule 1: DNA (5'-D(*TP*GP*GP*GP*CP*GP*GP*T)-3')

Chain P: 25% 75%




- Molecule 1: DNA (5'-D(*TP*GP*GP*GP*CP*GP*GP*T)-3')

Chain Q: 25% 75%



- Molecule 1: DNA (5'-D(*TP*GP*GP*GP*CP*GP*GP*T)-3')


Chain R: 38% 63%



4.2.4 Score per residue for model 4

- Molecule 1: DNA (5'-D(*TP*GP*GP*GP*CP*GP*GP*T)-3')

Chain O: 38% 63%



- Molecule 1: DNA (5'-D(*TP*GP*GP*GP*CP*GP*GP*T)-3')

Chain P:  38% 63%

T101
G102
G103
G104
C105
G106
G107
T108

- Molecule 1: DNA (5'-D(*TP*GP*GP*GP*CP*GP*GP*T)-3')

Chain Q:  38% 63%

T201
G202
G203
G204
C205
G206
G207
T208

- Molecule 1: DNA (5'-D(*TP*GP*GP*GP*CP*GP*GP*T)-3')

Chain R:  25% 75%

T401
G402
G403
G404
C405
G406
G407
T408

4.2.5 Score per residue for model 5

- Molecule 1: DNA (5'-D(*TP*GP*GP*GP*CP*GP*GP*T)-3')

Chain O:  38% 63%

T301
G302
G303
G304
C305
G306
G307
T308

- Molecule 1: DNA (5'-D(*TP*GP*GP*GP*CP*GP*GP*T)-3')

Chain P:  25% 75%

T101
G102
G103
G104
C105
G106
G107
T108

- Molecule 1: DNA (5'-D(*TP*GP*GP*GP*CP*GP*GP*T)-3')

Chain Q:  25% 75%

T201
G202
G203
G204
C205
G206
G207
T208



- Molecule 1: DNA (5'-D(*TP*GP*GP*GP*CP*GP*GP*T)-3')

Chain R:  25% 75%



4.2.6 Score per residue for model 6


- Molecule 1: DNA (5'-D(*TP*GP*GP*GP*CP*GP*GP*T)-3')




- Molecule 1: DNA (5'-D(*TP*GP*GP*GP*CP*GP*GP*T)-3')



- Molecule 1: DNA (5'-D(*TP*GP*GP*GP*CP*GP*GP*T)-3')



- Molecule 1: DNA (5'-D(*TP*GP*GP*GP*CP*GP*GP*T)-3')



4.2.7 Score per residue for model 7

- Molecule 1: DNA (5'-D(*TP*GP*GP*GP*CP*GP*GP*T)-3')



- Molecule 1: DNA (5'-D(*TP*GP*GP*GP*CP*GP*GP*T)-3')



- Molecule 1: DNA (5'-D(*TP*GP*GP*GP*CP*GP*GP*T)-3')

Chain Q:  38% 63%



- Molecule 1: DNA (5'-D(*TP*GP*GP*GP*CP*GP*GP*T)-3')


Chain R:  25% 75%



4.2.8 Score per residue for model 8

- Molecule 1: DNA (5'-D(*TP*GP*GP*GP*CP*GP*GP*T)-3')

Chain O:  38% 63%



- Molecule 1: DNA (5'-D(*TP*GP*GP*GP*CP*GP*GP*T)-3')

Chain P:  25% 75%



- Molecule 1: DNA (5'-D(*TP*GP*GP*GP*CP*GP*GP*T)-3')

Chain Q:  38% 63%



- Molecule 1: DNA (5'-D(*TP*GP*GP*GP*CP*GP*GP*T)-3')

Chain R:  25% 75%



4.2.9 Score per residue for model 9

- Molecule 1: DNA (5'-D(*TP*GP*GP*GP*CP*GP*GP*T)-3')

Chain O:  25% 75%

T301
G302
G303
G304
C305
G306
G307
T308

- Molecule 1: DNA (5'-D(*TP*GP*GP*GP*CP*GP*GP*T)-3')

Chain P:  25% 75%

T101
G102
G103
G104
C105
G106
G107
T108

- Molecule 1: DNA (5'-D(*TP*GP*GP*GP*CP*GP*GP*T)-3')

Chain Q:  38% 63%

T201
G202
G203
G204
C205
G206
G207
T208

- Molecule 1: DNA (5'-D(*TP*GP*GP*GP*CP*GP*GP*T)-3')

Chain R:  25% 75%

T401
G402
G403
G404
C405
G406
G407
T408

4.2.10 Score per residue for model 10

- Molecule 1: DNA (5'-D(*TP*GP*GP*GP*CP*GP*GP*T)-3')

Chain O:  38% 63%

T301
G302
G303
G304
C305
G306
G307
T308

- Molecule 1: DNA (5'-D(*TP*GP*GP*GP*CP*GP*GP*T)-3')

Chain P:  38% 63%

T101
G102
G103
G104
C105
G106
G107
T108

- Molecule 1: DNA (5'-D(*TP*GP*GP*GP*CP*GP*GP*T)-3')

Chain Q:  25% 75%

T201
G202
G203
G204
C205
G206
G207
T208

- Molecule 1: DNA (5'-D(*TP*GP*GP*GP*CP*GP*GP*T)-3')

Chain R:



T401	G402	G403	G404	C405	G406	G407	T408
------	------	------	------	------	------	------	------

5 Refinement protocol and experimental data overview

The models were refined using the following method: *Relaxation matrix refinement*.

Of the 10 calculated structures, 10 were deposited, based on the following criterion: *all calculated structures submitted*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
DISCOVER	structure solution	3.1
IRMA	refinement	2.3

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	BMRB entry 4612
Number of chemical shift lists	1
Total number of shifts	248
Number of shifts mapped to atoms	248
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	35%

No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality

6.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	O	1.55±0.00	1±0/186 (0.5±0.0%)	2.12±0.02	7±1/287 (2.6±0.3%)
1	P	1.56±0.01	1±0/186 (0.4±0.2%)	2.13±0.02	8±1/287 (2.7±0.3%)
1	Q	1.56±0.00	1±0/186 (0.5±0.2%)	2.12±0.03	7±1/287 (2.6±0.3%)
1	R	1.56±0.01	1±0/186 (0.5±0.2%)	2.14±0.03	8±1/287 (2.9±0.4%)
All	All	1.56	36/7440 (0.5%)	2.13	309/11480 (2.7%)

All unique bond outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
1	R	408	DT	C4'-O4'	-5.48	1.39	1.45	6	9
1	O	308	DT	C4'-O4'	-5.46	1.39	1.45	4	10
1	Q	208	DT	C4'-O4'	-5.40	1.39	1.45	1	9
1	P	108	DT	C4'-O4'	-5.37	1.39	1.45	1	8

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	R	408	DT	C6-C5-C7	-8.12	118.03	122.90	1	10
1	Q	208	DT	C6-C5-C7	-7.92	118.15	122.90	4	10
1	P	108	DT	C6-C5-C7	-7.73	118.26	122.90	1	10
1	O	308	DT	C6-C5-C7	-7.50	118.40	122.90	4	10
1	R	401	DT	O4'-C1'-C2'	-6.50	100.70	105.90	4	10
1	O	305	DC	O4'-C1'-N1	6.46	112.52	108.00	10	2
1	R	404	DG	O4'-C1'-C2'	-6.41	100.77	105.90	10	10
1	O	302	DG	O4'-C1'-C2'	-6.41	100.77	105.90	4	10
1	P	104	DG	O4'-C1'-C2'	-6.41	100.78	105.90	1	10
1	Q	202	DG	O4'-C1'-C2'	-6.38	100.80	105.90	2	10
1	R	405	DC	O4'-C1'-N1	6.33	112.43	108.00	1	2
1	R	402	DG	O4'-C1'-C2'	-6.33	100.84	105.90	9	10

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	P	101	DT	O4'-C1'-C2'	-6.28	100.87	105.90	4	8
1	O	304	DG	O4'-C1'-C2'	-6.28	100.88	105.90	5	10
1	P	102	DG	O4'-C1'-C2'	-6.27	100.88	105.90	8	10
1	Q	204	DG	O4'-C1'-C2'	-6.27	100.89	105.90	6	10
1	O	301	DT	O4'-C1'-C2'	-6.05	101.06	105.90	9	10
1	P	105	DC	O4'-C1'-C2'	-5.94	101.15	105.90	8	9
1	Q	205	DC	O4'-C1'-N1	5.90	112.13	108.00	10	1
1	O	305	DC	O4'-C1'-C2'	-5.83	101.24	105.90	1	4
1	O	301	DT	C6-C5-C7	-5.80	119.42	122.90	10	5
1	P	101	DT	C6-C5-C7	-5.69	119.48	122.90	4	6
1	Q	205	DC	O4'-C1'-C2'	-5.69	101.35	105.90	3	4
1	R	405	DC	O4'-C1'-C2'	-5.67	101.36	105.90	8	4
1	R	401	DT	C6-C5-C7	-5.65	119.51	122.90	4	9
1	R	408	DT	C1'-O4'-C4'	-5.64	104.45	110.10	1	8
1	O	306	DG	C4'-C3'-C2'	-5.64	98.03	103.10	9	8
1	P	106	DG	C4'-C3'-C2'	-5.63	98.03	103.10	1	9
1	Q	206	DG	C4'-C3'-C2'	-5.63	98.03	103.10	1	9
1	R	405	DC	O4'-C4'-C3'	-5.50	102.30	104.50	1	1
1	Q	201	DT	O4'-C1'-C2'	-5.47	101.53	105.90	1	10
1	O	308	DT	C1'-O4'-C4'	-5.46	104.64	110.10	2	8
1	P	108	DT	C1'-O4'-C4'	-5.45	104.65	110.10	3	9
1	R	406	DG	C4'-C3'-C2'	-5.42	98.22	103.10	10	10
1	R	401	DT	N1-C1'-C2'	5.38	122.81	112.60	4	9
1	Q	201	DT	N1-C1'-C2'	5.35	122.77	112.60	1	7
1	P	101	DT	N1-C1'-C2'	5.35	122.77	112.60	7	7
1	O	301	DT	N1-C1'-C2'	5.32	122.71	112.60	4	7
1	Q	201	DT	C6-C5-C7	-5.30	119.72	122.90	2	6
1	Q	208	DT	C1'-O4'-C4'	-5.26	104.84	110.10	4	7

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
All	All	6640	3680	3680	-

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is -.

There are no clashes.

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

There are no protein molecules in this entry.

6.3.2 Protein sidechains [i](#)

There are no protein molecules in this entry.

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues

There are no chain breaks in this entry.

7 Chemical shift validation [i](#)

The completeness of assignment taking into account all chemical shift lists is 35% for the well-defined parts and 35% for the entire structure.

7.1 Chemical shift list 1

File name: BMRB entry 4612

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	248
Number of shifts mapped to atoms	248
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.1.2 Chemical shift referencing [i](#)

No chemical shift referencing corrections were calculated (not enough data).

7.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 35%, i.e. 224 atoms were assigned a chemical shift out of a possible 644. 0 out of 0 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	0/0 (—%)	0/0 (—%)	0/0 (—%)	0/0 (—%)
Sidechain	0/0 (—%)	0/0 (—%)	0/0 (—%)	0/0 (—%)
Aromatic	0/0 (—%)	0/0 (—%)	0/0 (—%)	0/0 (—%)
Overall	224/644 (35%)	224/388 (58%)	0/204 (0%)	0/52 (0%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 35%, i.e. 224 atoms were assigned a chemical shift out of a possible 644. 0 out of 0 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹H	¹³C	¹⁵N
Backbone	0/0 (—%)	0/0 (—%)	0/0 (—%)	0/0 (—%)
Sidechain	0/0 (—%)	0/0 (—%)	0/0 (—%)	0/0 (—%)
Aromatic	0/0 (—%)	0/0 (—%)	0/0 (—%)	0/0 (—%)
Overall	224/644 (35%)	224/388 (58%)	0/204 (0%)	0/52 (0%)

7.1.4 Statistically unusual chemical shifts ⓘ

There are no statistically unusual chemical shifts.

7.1.5 Random Coil Index (RCI) plots ⓘ

No *random coil index* (RCI) plot could be generated from the current chemical shift list (assigned_chem_shift_list_1). RCI is only applicable to proteins.