



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 08:02 am GMT

PDB ID : 5EWX
Title : Fusion protein of T4 lysozyme and B4 domain of protein A from staphylococcal aureus with chemical cross-linker EY-CBS
Authors : Jeong, W.H.; Lee, H.; Song, D.H.; Lee, J.O.
Deposited on : 2015-11-22
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	2542 (2.60-2.60)
Clashscore	112137	2895 (2.60-2.60)
Ramachandran outliers	110173	2848 (2.60-2.60)
Sidechain outliers	110143	2848 (2.60-2.60)
RSRZ outliers	101464	2550 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	227	<div> <div>4%</div> <div> <div></div> <div>79%</div> <div>11%</div> <div>• 7%</div> </div> </div>
1	B	227	<div> <div></div> <div> <div>75%</div> <div>17%</div> <div>• 6%</div> </div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3472 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Endolysin,Immunoglobulin G-binding protein A,Endolysin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	210	Total	C	N	O	S	0	0	0
			1666	1043	301	315	7			
1	B	214	Total	C	N	O	S	0	0	0
			1682	1051	305	319	7			

There are 42 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	12	GLY	ARG	see sequence details	UNP P00720
A	1209	GLY	-	linker	UNP P00720
A	1210	GLY	-	linker	UNP P00720
A	1211	GLY	-	linker	UNP P00720
A	1212	GLY	-	linker	UNP P00720
A	1213	SER	-	linker	UNP P00720
A	1214	GLY	-	linker	UNP P00720
A	1215	GLY	-	linker	UNP P00720
A	1216	GLY	-	linker	UNP P00720
A	1217	GLY	-	linker	UNP P00720
A	1218	SER	-	linker	UNP P00720
A	1218A	VAL	ALA	engineered mutation	UNP P38507
A	1240	ALA	GLY	engineered mutation	UNP P38507
A	1258	CYS	GLU	engineered mutation	UNP P38507
A	1261	ALA	LYS	engineered mutation	UNP P38507
A	1262	ALA	LEU	engineered mutation	UNP P38507
A	40	CYS	ASN	engineered mutation	UNP P00720
A	54	THR	CYS	engineered mutation	UNP P00720
A	97	ALA	CYS	engineered mutation	UNP P00720
A	137	ARG	ILE	see sequence details	UNP P00720
A	162	ALA	LYS	engineered mutation	UNP P00720
B	12	GLY	ARG	see sequence details	UNP P00720
B	1209	GLY	-	linker	UNP P00720
B	1210	GLY	-	linker	UNP P00720
B	1211	GLY	-	linker	UNP P00720

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	1212	GLY	-	linker	UNP P00720
B	1213	SER	-	linker	UNP P00720
B	1214	GLY	-	linker	UNP P00720
B	1215	GLY	-	linker	UNP P00720
B	1216	GLY	-	linker	UNP P00720
B	1217	GLY	-	linker	UNP P00720
B	1218	SER	-	linker	UNP P00720
B	1218A	VAL	ALA	engineered mutation	UNP P38507
B	1240	ALA	GLY	engineered mutation	UNP P38507
B	1258	CYS	GLU	engineered mutation	UNP P38507
B	1261	ALA	LYS	engineered mutation	UNP P38507
B	1262	ALA	LEU	engineered mutation	UNP P38507
B	40	CYS	ASN	engineered mutation	UNP P00720
B	54	THR	CYS	engineered mutation	UNP P00720
B	97	ALA	CYS	engineered mutation	UNP P00720
B	137	ARG	ILE	see sequence details	UNP P00720
B	162	ALA	LYS	engineered mutation	UNP P00720

- Molecule 2 is 2,2'-ethyne-1,2-diylbis{5-[(chloroacetyl)amino]benzenesulfonic acid} (three-letter code: EYC) (formula: C₁₈H₁₄Cl₂N₂O₈S₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			30	18	2	8	2		
2	B	1	Total	C	N	O	S	0	0
			30	18	2	8	2		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	29	Total 29	O 29	0	0
3	B	35	Total 35	O 35	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	89.52Å 149.85Å 55.84Å 90.00° 122.69° 90.00°	Depositor
Resolution (Å)	42.74 – 2.60 42.74 – 2.59	Depositor EDS
% Data completeness (in resolution range)	94.1 (42.74-2.60) 94.8 (42.74-2.59)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.67 (at 2.58Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.9_1692)	Depositor
R, R_{free}	0.203 , 0.253 0.204 , 0.253	Depositor DCC
R_{free} test set	3536 reflections (9.99%)	DCC
Wilson B-factor (Å ²)	40.1	Xtrriage
Anisotropy	0.644	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 41.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	3472	wwPDB-VP
Average B, all atoms (Å ²)	56.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.03% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: EYC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.46	0/1690	0.63	0/2279
1	B	0.50	0/1706	0.65	1/2299 (0.0%)
All	All	0.48	0/3396	0.64	1/4578 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	7	LEU	CA-CB-CG	-5.32	103.07	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1666	0	1669	33	0
1	B	1682	0	1681	34	0
2	A	30	0	0	4	0
2	B	30	0	0	4	0
3	A	29	0	0	15	1
3	B	35	0	0	9	1
All	All	3472	0	3350	63	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (63) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:40:CYS:SG	2:A:1301:EYC:C19	2.15	1.34
1:A:40:CYS:SG	2:A:1301:EYC:C17	2.29	1.19
1:A:40:CYS:SG	2:A:1301:EYC:O18	2.12	1.07
1:A:76:ARG:HD3	3:A:1402:HOH:O	1.60	0.99
1:B:1221:GLN:NE2	3:B:1404:HOH:O	1.98	0.97
1:A:64:GLU:OE2	3:A:1403:HOH:O	1.84	0.95
1:A:1:MET:N	3:A:1405:HOH:O	2.01	0.91
1:B:1221:GLN:OE1	3:B:1402:HOH:O	1.87	0.91
1:B:1219:GLU:N	3:B:1402:HOH:O	2.07	0.88
1:A:76:ARG:NH1	3:A:1402:HOH:O	1.83	0.83
1:A:154:ARG:NE	3:A:1401:HOH:O	1.81	0.81
1:A:163:ASN:HB3	1:B:136:SER:HB2	1.65	0.78
1:A:139:TYR:OH	3:A:1404:HOH:O	2.01	0.78
1:B:40:CYS:SG	2:B:1301:EYC:C17	2.74	0.76
1:B:26:THR:HG22	1:B:32:LEU:HA	1.66	0.76
1:B:40:CYS:SG	2:B:1301:EYC:O18	2.43	0.75
1:A:164:LEU:H	1:B:114:PHE:HA	1.52	0.75
1:B:115:THR:OG1	3:B:1403:HOH:O	1.90	0.74
1:B:40:CYS:SG	2:B:1301:EYC:C19	2.79	0.71
1:B:1221:GLN:OE1	3:B:1404:HOH:O	2.08	0.71
1:A:1221:GLN:CA	3:A:1408:HOH:O	2.38	0.71
1:B:1221:GLN:CD	3:B:1404:HOH:O	2.26	0.71
1:A:1221:GLN:HB3	3:A:1408:HOH:O	1.91	0.70
1:A:65:LYS:O	3:A:1406:HOH:O	2.13	0.66
1:A:40:CYS:HG	2:A:1301:EYC:C17	1.92	0.66
1:B:105:GLN:OE1	3:B:1405:HOH:O	2.14	0.65
1:A:163:ASN:HB2	1:B:114:PHE:CE2	2.32	0.65
1:A:163:ASN:CB	1:B:136:SER:HB2	2.29	0.60
1:B:88:TYR:CZ	1:B:96:ARG:HD3	2.37	0.60
1:A:114:PHE:O	1:A:118:LEU:HB2	2.01	0.59
1:B:80:ARG:NH1	3:B:1407:HOH:O	2.32	0.57
1:B:76:ARG:NH1	3:B:1401:HOH:O	1.82	0.57
1:A:1221:GLN:CB	3:A:1408:HOH:O	2.50	0.57
1:A:1221:GLN:HA	3:A:1408:HOH:O	2.01	0.56
1:B:2:ASN:ND2	1:B:5:GLU:HG2	2.20	0.55
1:B:1219:GLU:N	1:B:1221:GLN:OE1	2.38	0.55
1:A:157:THR:O	3:A:1407:HOH:O	2.19	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:87:VAL:O	1:A:91:LEU:HG	2.11	0.50
1:B:1209:GLY:O	1:B:1210:GLY:C	2.49	0.49
1:A:76:ARG:CD	3:A:1402:HOH:O	2.35	0.49
1:B:163:ASN:HA	1:B:164:LEU:C	2.35	0.48
1:B:105:GLN:HB2	1:B:145:ARG:NH2	2.29	0.47
1:B:1232:ASN:HB2	1:B:1263:ASN:OD1	2.14	0.47
1:A:1:MET:HG2	1:A:158:TRP:HB3	1.97	0.46
1:A:154:ARG:CD	3:A:1401:HOH:O	2.49	0.46
1:A:121:LEU:HD13	1:A:129:ALA:HB1	1.98	0.45
1:A:81:ASN:HD22	1:A:108:GLU:HG2	1.82	0.45
1:A:85:LYS:NZ	1:A:89:ASP:OD2	2.49	0.43
1:B:118:LEU:HA	1:B:118:LEU:HD12	1.91	0.43
1:B:52:ARG:NH2	1:B:62:GLU:OE1	2.40	0.43
1:A:129:ALA:O	1:A:133:LEU:HD22	2.19	0.43
1:B:163:ASN:O	1:B:163:ASN:ND2	2.52	0.43
1:B:1258:CYS:SG	2:B:1301:EYC:C03	3.08	0.42
1:B:1245:LEU:HA	1:B:1255:LEU:HD23	2.01	0.42
1:B:14:ARG:HG3	1:B:18:TYR:CE1	2.55	0.42
1:A:163:ASN:O	1:A:163:ASN:ND2	2.50	0.41
1:A:1221:GLN:C	3:A:1408:HOH:O	2.57	0.41
1:A:79:LEU:HA	1:A:79:LEU:HD12	1.83	0.41
1:A:2:ASN:ND2	1:A:5:GLU:HG2	2.36	0.41
1:B:85:LYS:HB3	1:B:86:PRO:HD3	2.01	0.41
1:B:9:ILE:HD13	1:B:164:LEU:HD23	2.02	0.41
1:B:87:VAL:O	1:B:91:LEU:HG	2.21	0.41
1:B:79:LEU:O	1:B:85:LYS:HD3	2.21	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:1421:HOH:O	3:B:1417:HOH:O[1_556]	1.73	0.47

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	206/227 (91%)	202 (98%)	4 (2%)	0	100	100
1	B	210/227 (92%)	206 (98%)	4 (2%)	0	100	100
All	All	416/454 (92%)	408 (98%)	8 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	174/183 (95%)	165 (95%)	9 (5%)	27	52
1	B	174/183 (95%)	164 (94%)	10 (6%)	24	47
All	All	348/366 (95%)	329 (94%)	19 (6%)	25	49

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1252	SER
1	A	1255	LEU
1	A	38	SER
1	A	79	LEU
1	A	94	VAL
1	A	118	LEU
1	A	121	LEU
1	A	133	LEU
1	A	163	ASN
1	B	35	LYS
1	B	1228	LEU
1	B	1252	SER
1	B	1255	LEU
1	B	38	SER
1	B	79	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	106	MET
1	B	118	LEU
1	B	121	LEU
1	B	163	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	EYC	A	1301	-	31,31,33	2.64	8 (25%)	42,46,48	3.36	19 (45%)
2	EYC	B	1301	-	31,31,33	2.25	8 (25%)	42,46,48	2.18	19 (45%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means

no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EYC	A	1301	-	-	0/25/25/29	0/2/2/2
2	EYC	B	1301	-	-	0/25/25/29	0/2/2/2

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1301	EYC	C21-C22	-2.46	1.35	1.39
2	B	1301	EYC	O01-C02	-2.25	1.18	1.23
2	B	1301	EYC	C17-N16	2.85	1.40	1.35
2	A	1301	EYC	C17-N16	3.02	1.41	1.35
2	A	1301	EYC	C06-N05	3.22	1.47	1.41
2	A	1301	EYC	C13-C14	3.26	1.44	1.38
2	B	1301	EYC	C02-N05	3.58	1.42	1.35
2	A	1301	EYC	O26-S23	4.08	1.65	1.43
2	B	1301	EYC	O26-S23	4.18	1.65	1.43
2	A	1301	EYC	O31-S28	4.20	1.65	1.43
2	B	1301	EYC	O31-S28	4.25	1.66	1.43
2	B	1301	EYC	C12-C11	5.17	1.52	1.43
2	B	1301	EYC	C09-C10	5.48	1.53	1.43
2	A	1301	EYC	C09-C10	5.59	1.53	1.43
2	A	1301	EYC	C12-C11	5.65	1.53	1.43
2	A	1301	EYC	C02-N05	7.42	1.48	1.35

All (38) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1301	EYC	O26-S23-C22	-13.03	91.15	106.17
2	A	1301	EYC	C21-C15-N16	-4.77	104.81	120.16
2	A	1301	EYC	C21-C22-S23	-4.63	107.91	116.92
2	A	1301	EYC	C09-C27-S28	-4.55	117.08	122.92
2	B	1301	EYC	C21-C15-N16	-4.47	105.77	120.16
2	A	1301	EYC	O18-C17-N16	-4.06	117.78	123.06
2	B	1301	EYC	C13-C14-C15	-3.86	115.78	120.30
2	B	1301	EYC	C32-C06-N05	-3.68	108.33	120.16
2	B	1301	EYC	O18-C17-N16	-3.66	118.31	123.06
2	A	1301	EYC	O01-C02-C03	-3.65	115.42	122.06
2	A	1301	EYC	C22-C12-C11	-3.53	113.53	121.57
2	A	1301	EYC	C13-C14-C15	-3.01	116.77	120.30
2	B	1301	EYC	C19-C17-N16	-3.00	110.62	115.02
2	B	1301	EYC	C22-C12-C11	-2.86	115.04	121.57
2	B	1301	EYC	O01-C02-N05	-2.77	119.47	123.06

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1301	EYC	C19-C17-N16	-2.71	111.03	115.02
2	A	1301	EYC	C07-C06-C32	-2.57	116.64	119.66
2	B	1301	EYC	O31-S28-C27	-2.54	103.24	106.17
2	B	1301	EYC	C27-C09-C10	-2.46	115.95	121.57
2	B	1301	EYC	C21-C22-S23	-2.17	112.70	116.92
2	B	1301	EYC	C13-C12-C11	2.20	124.46	120.20
2	A	1301	EYC	C32-C27-S28	2.26	121.32	116.92
2	B	1301	EYC	O18-C17-C19	2.44	126.49	122.06
2	B	1301	EYC	C08-C09-C10	2.49	125.03	120.20
2	B	1301	EYC	O30-S28-C27	2.65	109.22	106.17
2	A	1301	EYC	C15-C21-C22	2.74	122.13	119.25
2	B	1301	EYC	C12-C22-S23	2.96	126.72	122.92
2	A	1301	EYC	O30-S28-C27	3.02	109.65	106.17
2	B	1301	EYC	C07-C06-N05	3.03	130.56	120.41
2	B	1301	EYC	C14-C15-N16	3.06	130.67	120.41
2	B	1301	EYC	C14-C15-C21	3.32	123.56	119.66
2	B	1301	EYC	O24-S23-C22	3.41	110.10	106.17
2	A	1301	EYC	O18-C17-C19	3.63	128.66	122.06
2	A	1301	EYC	C14-C15-N16	4.13	134.25	120.41
2	A	1301	EYC	C03-C02-N05	4.42	121.52	115.02
2	A	1301	EYC	O24-S23-C22	4.67	111.55	106.17
2	A	1301	EYC	C13-C12-C11	4.94	129.79	120.20
2	A	1301	EYC	C12-C22-S23	5.76	130.32	122.92

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1301	EYC	4	0
2	B	1301	EYC	4	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	210/227 (92%)	0.02	9 (4%) 36 28	33, 52, 111, 130	0
1	B	214/227 (94%)	-0.17	0 100 100	31, 48, 85, 101	0
All	All	424/454 (93%)	-0.08	9 (2%) 64 58	31, 50, 97, 130	0

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1227	ILE	3.2
1	A	1247	ASP	3.1
1	A	1249	PRO	3.0
1	A	1225	TYR	2.9
1	A	1224	PHE	2.8
1	A	35	LYS	2.8
1	A	1248	ASP	2.5
1	A	34	THR	2.4
1	A	1228	LEU	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron

density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	EYC	B	1301	30/32	0.95	0.19	1.31	22,55,76,102	0
2	EYC	A	1301	30/32	0.94	0.21	1.29	15,58,104,111	0

6.5 Other polymers [i](#)

There are no such residues in this entry.