



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 09:32 am GMT

PDB ID : 4EXJ
Title : Crystal structure of glutathione s-transferase like protein lelg_03239 (target efi-501752) from lodderomyces elongisporus
Authors : Patskovsky, Y.; Toro, R.; Bhosle, R.; Zencheck, W.D.; Hillerich, B.; Seidel, R.D.; Washington, E.; Scott Glenn, A.; Chowdhury, S.; Evans, B.; Hammonds, J.; Imker, H.J.; Armstrong, R.N.; Gerlt, J.A.; Almo, S.C.; Enzyme Function Initiative (EFI)
Deposited on : 2012-04-30
Resolution : 1.64 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.64 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	2211 (1.66-1.62)
Clashscore	112137	2356 (1.66-1.62)
Ramachandran outliers	110173	2315 (1.66-1.62)
Sidechain outliers	110143	2315 (1.66-1.62)
RSRZ outliers	101464	2219 (1.66-1.62)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	238	
1	B	238	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4096 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called uncharacterized protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	221	Total	C	N	O	S	0	5	0
			1851	1203	304	341	3			
1	B	222	Total	C	N	O	S	0	3	0
			1858	1212	306	337	3			

There are 48 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	MET	-	EXPRESSION TAG	UNP A5E0V2
A	0	VAL	-	EXPRESSION TAG	UNP A5E0V2
A	215	ALA	-	EXPRESSION TAG	UNP A5E0V2
A	216	GLU	-	EXPRESSION TAG	UNP A5E0V2
A	217	ASN	-	EXPRESSION TAG	UNP A5E0V2
A	218	LEU	-	EXPRESSION TAG	UNP A5E0V2
A	219	TYR	-	EXPRESSION TAG	UNP A5E0V2
A	220	PHE	-	EXPRESSION TAG	UNP A5E0V2
A	221	GLN	-	EXPRESSION TAG	UNP A5E0V2
A	222	SER	-	EXPRESSION TAG	UNP A5E0V2
A	223	HIS	-	EXPRESSION TAG	UNP A5E0V2
A	224	HIS	-	EXPRESSION TAG	UNP A5E0V2
A	225	HIS	-	EXPRESSION TAG	UNP A5E0V2
A	226	HIS	-	EXPRESSION TAG	UNP A5E0V2
A	227	HIS	-	EXPRESSION TAG	UNP A5E0V2
A	228	HIS	-	EXPRESSION TAG	UNP A5E0V2
A	229	TRP	-	EXPRESSION TAG	UNP A5E0V2
A	230	SER	-	EXPRESSION TAG	UNP A5E0V2
A	231	HIS	-	EXPRESSION TAG	UNP A5E0V2
A	232	PRO	-	EXPRESSION TAG	UNP A5E0V2
A	233	GLN	-	EXPRESSION TAG	UNP A5E0V2
A	234	PHE	-	EXPRESSION TAG	UNP A5E0V2
A	235	GLU	-	EXPRESSION TAG	UNP A5E0V2
A	236	LYS	-	EXPRESSION TAG	UNP A5E0V2
B	-1	MET	-	EXPRESSION TAG	UNP A5E0V2

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	VAL	-	EXPRESSION TAG	UNP A5E0V2
B	215	ALA	-	EXPRESSION TAG	UNP A5E0V2
B	216	GLU	-	EXPRESSION TAG	UNP A5E0V2
B	217	ASN	-	EXPRESSION TAG	UNP A5E0V2
B	218	LEU	-	EXPRESSION TAG	UNP A5E0V2
B	219	TYR	-	EXPRESSION TAG	UNP A5E0V2
B	220	PHE	-	EXPRESSION TAG	UNP A5E0V2
B	221	GLN	-	EXPRESSION TAG	UNP A5E0V2
B	222	SER	-	EXPRESSION TAG	UNP A5E0V2
B	223	HIS	-	EXPRESSION TAG	UNP A5E0V2
B	224	HIS	-	EXPRESSION TAG	UNP A5E0V2
B	225	HIS	-	EXPRESSION TAG	UNP A5E0V2
B	226	HIS	-	EXPRESSION TAG	UNP A5E0V2
B	227	HIS	-	EXPRESSION TAG	UNP A5E0V2
B	228	HIS	-	EXPRESSION TAG	UNP A5E0V2
B	229	TRP	-	EXPRESSION TAG	UNP A5E0V2
B	230	SER	-	EXPRESSION TAG	UNP A5E0V2
B	231	HIS	-	EXPRESSION TAG	UNP A5E0V2
B	232	PRO	-	EXPRESSION TAG	UNP A5E0V2
B	233	GLN	-	EXPRESSION TAG	UNP A5E0V2
B	234	PHE	-	EXPRESSION TAG	UNP A5E0V2
B	235	GLU	-	EXPRESSION TAG	UNP A5E0V2
B	236	LYS	-	EXPRESSION TAG	UNP A5E0V2

- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total Cl 1 1	0	0
2	A	2	Total Cl 2 2	0	0

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	165	Total	O	0	1
			166	166		
4	B	208	Total	O	0	0
			208	208		

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	54.67Å 87.19Å 55.35Å 90.00° 115.17° 90.00°	Depositor
Resolution (Å)	50.00 – 1.64 31.78 – 1.64	Depositor EDS
% Data completeness (in resolution range)	98.9 (50.00-1.64) 98.9 (31.78-1.64)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.33 (at 1.64Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.184 , 0.225 0.186 , 0.223	Depositor DCC
R_{free} test set	1765 reflections (3.21%)	DCC
Wilson B-factor (Å ²)	26.9	Xtriage
Anisotropy	0.341	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 45.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.019 for l,-k,h	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	4096	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.05% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.54	0/1913	0.57	0/2595
1	B	0.51	0/1920	0.60	0/2607
All	All	0.52	0/3833	0.58	0/5202

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1851	0	1814	21	0
1	B	1858	0	1822	14	0
2	A	2	0	0	0	0
2	B	1	0	0	0	0
3	A	5	0	0	0	0
3	B	5	0	0	1	0
4	A	166	0	0	3	0
4	B	208	0	0	5	0
All	All	4096	0	3636	35	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:6[A]:THR:HG23	1:A:7:GLY:O	1.78	0.83
1:A:32:HIS:O	1:A:35:THR:HB	1.81	0.79
1:B:189:ASN:HB2	4:B:481:HOH:O	1.86	0.76
1:A:195:LYS:HD2	1:A:199:ARG:HH12	1.57	0.67
1:B:195:LYS:HG2	1:B:199:ARG:NH2	2.10	0.67
1:A:6[A]:THR:HG23	1:A:7:GLY:N	2.14	0.63
1:B:195:LYS:NZ	1:B:199:ARG:HH22	1.98	0.62
1:B:20:LYS:HE2	4:B:570:HOH:O	2.00	0.60
1:A:81:HIS:CE1	1:A:86:SER:HB3	2.39	0.58
1:B:14:LYS:HD3	1:B:165:ALA:HA	1.87	0.56
1:A:13:ARG:NH2	1:A:212:GLU:O	2.40	0.54
1:A:90[B]:ASP:OD2	1:A:93:LEU:HD12	2.07	0.54
1:B:187:LYS:HE3	4:B:468:HOH:O	2.07	0.54
1:B:167:LEU:C	1:B:167:LEU:HD12	2.29	0.52
1:A:167:LEU:C	1:A:167:LEU:HD12	2.31	0.51
1:B:195:LYS:HZ2	1:B:199:ARG:HH22	1.58	0.50
1:A:8:PRO:HB3	1:A:13:ARG:NH2	2.28	0.49
1:A:7:GLY:C	4:A:538:HOH:O	2.51	0.48
1:B:39:GLN:NE2	4:B:572:HOH:O	2.44	0.48
1:B:117:ASN:ND2	3:B:302:SO4:O3	2.47	0.47
1:A:32:HIS:O	1:A:35:THR:CB	2.57	0.47
1:B:14:LYS:HB2	1:B:15:PRO:HD3	1.97	0.46
1:A:147:LYS:O	1:A:151:ARG:HD3	2.15	0.46
1:B:180:SER:HB2	1:B:181:PRO:HD2	1.98	0.46
1:B:91:PRO:HG3	4:B:422:HOH:O	2.15	0.46
1:A:51:ILE:HD12	4:A:436:HOH:O	2.17	0.45
1:A:13:ARG:HD2	1:A:16:LEU:HD23	1.99	0.44
1:A:81:HIS:HE1	1:A:86:SER:HB3	1.81	0.43
1:A:13:ARG:HA	1:A:13:ARG:HD3	1.65	0.43
1:A:14:LYS:HB2	1:A:15:PRO:CD	2.50	0.42
1:A:4:LEU:HG	1:A:6[B]:THR:CG2	2.51	0.41
1:A:151:ARG:HB3	1:A:151:ARG:HE	1.31	0.41
1:A:153:TRP:CH2	1:A:199:ARG:HG2	2.55	0.41
1:A:12:GLY:HA3	4:A:538:HOH:O	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	222/238 (93%)	214 (96%)	7 (3%)	1 (0%)	32	11
1	B	223/238 (94%)	216 (97%)	7 (3%)	0	100	100
All	All	445/476 (94%)	430 (97%)	14 (3%)	1 (0%)	51	27

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	8	PRO

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	201/214 (94%)	194 (96%)	7 (4%)	41	12
1	B	200/214 (94%)	198 (99%)	2 (1%)	80	63
All	All	401/428 (94%)	392 (98%)	9 (2%)	60	28

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	6[A]	THR
1	A	6[B]	THR
1	A	36	LYS
1	A	52	VAL
1	A	89	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	151	ARG
1	A	216	GLU
1	B	62	PRO
1	B	167	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	81	HIS
1	A	107	GLN
1	A	109	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 3 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SO4	A	302	-	4,4,4	0.15	0	6,6,6	0.17	0
3	SO4	B	302	-	4,4,4	0.69	0	6,6,6	0.31	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SO4	A	302	-	-	0/0/0/0	0/0/0/0
3	SO4	B	302	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	302	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	221/238 (92%)	0.18	16 (7%) 16 14	24, 35, 58, 71	0
1	B	222/238 (93%)	-0.13	4 (1%) 69 71	20, 30, 48, 66	0
All	All	443/476 (93%)	0.02	20 (4%) 34 32	20, 33, 57, 71	0

All (20) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	0	VAL	4.7
1	A	7	GLY	4.6
1	A	124	HIS	4.2
1	B	102	PHE	2.8
1	A	122	ASN	2.7
1	A	121	GLN	2.6
1	A	125	ILE	2.6
1	A	105	ALA	2.5
1	A	101	LEU	2.4
1	B	105	ALA	2.4
1	A	0	VAL	2.3
1	B	32	HIS	2.2
1	A	37	ASP	2.2
1	A	189	ASN	2.2
1	A	10	GLY	2.1
1	A	9	THR	2.1
1	A	178	HIS	2.1
1	A	102	PHE	2.0
1	A	123	GLY	2.0
1	A	213	HIS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	SO4	A	302	5/5	0.72	0.22	1.75	39,41,49,52	5
3	SO4	B	302	5/5	0.97	0.08	-0.91	31,36,44,47	0
2	CL	A	303	1/1	0.98	0.03	-1.83	47,47,47,47	0
2	CL	A	301	1/1	0.94	0.09	-1.89	46,46,46,46	0
2	CL	B	301	1/1	0.96	0.05	-4.92	39,39,39,39	0

6.5 Other polymers [i](#)

There are no such residues in this entry.