



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 08:36 am GMT

PDB ID : 4EXP  
Title : Structure of mouse Interleukin-34 in complex with mouse FMS  
Authors : Liu, H.; Leo, C.; Chen, X.; Wong, B.R.; Williams, L.T.; Lin, H.; He, X.  
Deposited on : 2012-04-30  
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	2583 (2.80-2.80)
Clashscore	112137	3033 (2.80-2.80)
Ramachandran outliers	110173	2983 (2.80-2.80)
Sidechain outliers	110143	2985 (2.80-2.80)
RSRZ outliers	101464	2610 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	177	<div> <div> <div></div> <div>62%</div> <div>31%</div> <div>...</div> </div> <div>11%</div> </div>
2	X	283	<div> <div> <div></div> <div>66%</div> <div>31%</div> <div>.</div> </div> <div>11%</div> </div>

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3838 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Interleukin-34.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	171	Total	C	N	O	S	0	0	0
			1391	894	238	249	10			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	195	ALA	-	EXPRESSION TAG	UNP Q8R1R4
A	196	ALA	-	EXPRESSION TAG	UNP Q8R1R4
A	197	ALA	-	EXPRESSION TAG	UNP Q8R1R4

- Molecule 2 is a protein called Macrophage colony-stimulating factor 1 receptor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	X	283	Total	C	N	O	S	0	0	0
			2193	1381	377	425	10			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
X	17	ALA	-	EXPRESSION TAG	UNP P09581
X	18	ASP	-	EXPRESSION TAG	UNP P09581
X	19	PRO	-	EXPRESSION TAG	UNP P09581
X	299	HIS	-	EXPRESSION TAG	UNP P09581

- Molecule 3 is SUGAR (N-ACETYL-D-GLUCOSAMINE) (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	X	1	Total	C	N	O	0	0
			14	8	1	5		
3	X	1	Total	C	N	O	0	0
			14	8	1	5		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	103	Total	O	0	0
			103	103		
4	X	95	Total	O	0	0
			95	95		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Interleukin-34



#### • Molecule 2: Macrophage colony-stimulating factor 1 receptor



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	44.71Å 135.34Å 203.49Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.80 47.91 – 2.80	Depositor EDS
% Data completeness (in resolution range)	99.3 (50.00-2.80) 99.6 (47.91-2.80)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.44 (at 2.81Å)	Xtriage
Refinement program	CNS	Depositor
R, $R_{free}$	0.248 , 0.279 0.251 , 0.285	Depositor DCC
$R_{free}$ test set	772 reflections (4.95%)	DCC
Wilson B-factor (Å <sup>2</sup> )	54.4	Xtriage
Anisotropy	0.353	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.27 , 67.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.45$ , $\langle L^2 \rangle = 0.27$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	3838	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	85.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.12% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.54	0/1421	0.70	2/1928 (0.1%)
2	X	0.41	0/2241	0.58	0/3055
All	All	0.47	0/3662	0.63	2/4983 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	188	TRP	CB-CA-C	-8.14	94.11	110.40
1	A	188	TRP	N-CA-C	7.00	129.89	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1391	0	1415	37	0
2	X	2193	0	2168	70	0
3	A	28	0	26	1	0
3	X	28	0	26	5	0
4	A	103	0	0	9	0
4	X	95	0	0	12	0
All	All	3838	0	3635	109	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

All (109) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:X:212:LEU:HD12	2:X:214:ARG:HH21	1.39	0.87
2:X:75:THR:H	2:X:78:ASN:HD22	1.22	0.87
2:X:202:PRO:HB2	2:X:287:ARG:HG2	1.59	0.82
2:X:73:ASN:HB2	3:X:1001:NAG:H82	1.63	0.81
1:A:113:HIS:HD2	1:A:115:SER:H	1.29	0.79
1:A:138:ILE:HG23	4:A:2182:HOH:O	1.84	0.76
2:X:298:SER:O	2:X:299:HIS:HB2	1.86	0.75
2:X:206:LYS:HD2	4:X:1181:HOH:O	1.89	0.72
1:A:117:LYS:O	1:A:121:GLU:HG3	1.89	0.71
2:X:118:VAL:HG12	2:X:198:VAL:HG22	1.70	0.71
1:A:74:VAL:HG23	4:A:2115:HOH:O	1.88	0.71
2:X:17:ALA:HB1	3:X:1000:NAG:H2	1.73	0.71
2:X:218:GLU:HA	2:X:267:ALA:HA	1.72	0.71
2:X:134:LEU:HG	2:X:181:VAL:HG11	1.73	0.70
2:X:54:ILE:HG12	2:X:58:TRP:HD1	1.55	0.69
2:X:110:LEU:HD12	2:X:113:GLN:HE21	1.55	0.69
2:X:246:ILE:HD11	2:X:261:ARG:HB3	1.73	0.69
2:X:109:ASN:HD22	2:X:128:LEU:HD12	1.57	0.68
1:A:88:ARG:NH1	1:A:136:VAL:HG22	2.09	0.68
2:X:119:GLU:HB3	2:X:198:VAL:HG23	1.77	0.67
2:X:203:PRO:HD3	2:X:282:ASN:OD1	1.95	0.67
1:A:91:ARG:O	1:A:95:VAL:HG23	1.96	0.64
2:X:110:LEU:H	2:X:110:LEU:HD23	1.63	0.64
2:X:109:ASN:HB3	2:X:128:LEU:HB2	1.81	0.63
1:A:74:VAL:O	1:A:78:THR:HG23	1.98	0.62
2:X:212:LEU:HD12	2:X:214:ARG:NH2	2.14	0.61
1:A:77:ILE:O	1:A:81:GLN:HG3	2.01	0.61
4:A:2117:HOH:O	2:X:251:ASP:HB2	2.01	0.60
2:X:77:LYS:HD3	2:X:109:ASN:OD1	2.01	0.60
1:A:38:THR:OG1	1:A:188:TRP:HH2	1.85	0.59
2:X:59:THR:HG21	4:X:1168:HOH:O	2.02	0.58
2:X:146:ARG:HG2	2:X:146:ARG:HH21	1.68	0.58
1:A:38:THR:HG1	1:A:188:TRP:HH2	1.49	0.57
2:X:202:PRO:HA	2:X:282:ASN:OD1	2.04	0.57
1:A:79:ARG:HA	1:A:82:LYS:HE3	1.87	0.57
1:A:33:LYS:HE3	4:A:2116:HOH:O	2.05	0.56
2:X:77:LYS:HD2	4:X:1136:HOH:O	2.06	0.55

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:X:228:ASN:ND2	2:X:230:GLU:H	2.04	0.55
2:X:75:THR:HG22	2:X:78:ASN:ND2	2.22	0.55
2:X:291:MET:HE3	2:X:293:PHE:HB2	1.89	0.54
2:X:170:LEU:H	2:X:170:LEU:HD12	1.72	0.54
1:A:182:GLN:HB2	4:A:2106:HOH:O	2.08	0.54
1:A:98:SER:O	1:A:102:THR:HG23	2.08	0.54
1:A:85:VAL:HG13	1:A:142:VAL:HG22	1.89	0.54
1:A:123:GLN:O	1:A:127:GLU:HG3	2.08	0.54
2:X:83:ARG:HG3	2:X:96:THR:HG22	1.90	0.53
2:X:73:ASN:CB	3:X:1001:NAG:H82	2.37	0.53
1:A:147:SER:O	1:A:151:THR:HG23	2.08	0.53
2:X:75:THR:H	2:X:78:ASN:ND2	2.01	0.52
1:A:113:HIS:CD2	1:A:115:SER:H	2.17	0.52
2:X:110:LEU:HD13	2:X:189:THR:OG1	2.10	0.51
1:A:42:ARG:NH2	4:A:2101:HOH:O	2.43	0.51
1:A:79:ARG:HD3	4:A:2198:HOH:O	2.11	0.51
2:X:212:LEU:HA	4:X:1163:HOH:O	2.10	0.50
2:X:296:VAL:HG12	2:X:297:GLU:N	2.26	0.50
1:A:192:GLU:OE1	1:A:192:GLU:HA	2.12	0.49
1:A:35:CYS:O	1:A:188:TRP:HZ3	1.95	0.49
2:X:110:LEU:HB3	4:X:1118:HOH:O	2.11	0.49
1:A:38:THR:CB	1:A:188:TRP:HH2	2.26	0.49
1:A:45:LEU:CD2	1:A:165:LEU:HD12	2.43	0.49
1:A:94:TRP:CZ2	1:A:172:MET:HB2	2.48	0.49
2:X:226:ALA:HB3	2:X:235:VAL:CG2	2.43	0.49
1:A:79:ARG:HH12	3:A:2000:NAG:HN2	1.61	0.48
2:X:287:ARG:HG3	2:X:287:ARG:NH1	2.28	0.48
2:X:50:TRP:CZ3	2:X:84:CYS:HB2	2.48	0.48
2:X:216:ARG:HG3	2:X:218:GLU:O	2.13	0.48
2:X:215:ILE:HG12	2:X:215:ILE:O	2.13	0.48
2:X:56:PRO:HD3	4:X:1122:HOH:O	2.14	0.47
1:A:113:HIS:HD2	1:A:115:SER:N	2.06	0.47
2:X:172:SER:OG	2:X:194:LYS:HA	2.15	0.47
2:X:110:LEU:HA	4:X:1118:HOH:O	2.16	0.46
2:X:291:MET:CE	2:X:293:PHE:HB2	2.45	0.46
1:A:41:LEU:HD11	1:A:126:LEU:HD23	1.97	0.46
2:X:56:PRO:CD	4:X:1122:HOH:O	2.64	0.46
1:A:60:ILE:HG22	1:A:61:ASN:N	2.31	0.45
2:X:146:ARG:NH2	2:X:146:ARG:HG2	2.32	0.45
2:X:54:ILE:HD11	2:X:57:TYR:HB2	1.98	0.45
2:X:287:ARG:HG3	2:X:287:ARG:HH11	1.80	0.45

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:161:PRO:HG2	4:A:2141:HOH:O	2.17	0.44
2:X:197:ARG:NH2	4:X:1162:HOH:O	2.50	0.44
1:A:145:VAL:O	1:A:149:LEU:HG	2.18	0.44
2:X:50:TRP:CD1	2:X:67:SER:HB3	2.53	0.44
2:X:141:MET:HG3	2:X:147:GLN:HA	1.99	0.43
2:X:179:THR:HG22	2:X:181:VAL:HG13	1.99	0.43
2:X:152:THR:HG23	2:X:171:ASP:CG	2.39	0.43
1:A:151:THR:N	1:A:152:PRO:HA	2.34	0.43
2:X:118:VAL:CG1	2:X:198:VAL:HG22	2.46	0.43
1:A:109:LEU:HG	1:A:113:HIS:CG	2.54	0.43
2:X:180:MET:HG3	2:X:185:GLU:HG3	2.00	0.43
2:X:47:SER:HB2	2:X:87:LEU:HB2	2.00	0.42
3:X:1001:NAG:H83	4:X:1167:HOH:O	2.18	0.42
1:A:85:VAL:CG1	1:A:90:LEU:HD13	2.50	0.42
3:X:1001:NAG:C8	4:X:1167:HOH:O	2.67	0.42
1:A:38:THR:OG1	1:A:188:TRP:CH2	2.65	0.42
2:X:134:LEU:HB3	2:X:179:THR:HG21	2.01	0.42
2:X:147:GLN:NE2	2:X:147:GLN:H	2.18	0.42
2:X:239:ARG:HB2	2:X:244:LEU:HD11	2.02	0.42
1:A:108:VAL:O	1:A:108:VAL:HG12	2.18	0.41
2:X:110:LEU:HD12	2:X:113:GLN:NE2	2.30	0.41
2:X:127:CYS:H	2:X:161:GLY:HA3	1.86	0.41
2:X:40:LEU:HB2	2:X:69:LEU:HB2	2.01	0.41
1:A:28:THR:HG21	4:A:2158:HOH:O	2.20	0.41
2:X:189:THR:HG23	4:X:1173:HOH:O	2.21	0.41
2:X:253:GLN:HE21	2:X:253:GLN:HB2	1.61	0.41
2:X:23:ILE:HD11	2:X:95:THR:HG22	2.02	0.41
2:X:141:MET:CE	2:X:178:LYS:HD2	2.51	0.40
2:X:233:PHE:CZ	2:X:259:LYS:HG3	2.56	0.40
2:X:215:ILE:HA	2:X:296:VAL:HB	2.02	0.40
2:X:70:THR:HG22	2:X:71:THR:N	2.37	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	169/177 (96%)	154 (91%)	8 (5%)	7 (4%)	3	11
2	X	281/283 (99%)	246 (88%)	29 (10%)	6 (2%)	8	27
All	All	450/460 (98%)	400 (89%)	37 (8%)	13 (3%)	5	18

All (13) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	194	PRO
2	X	65	PRO
2	X	89	ASP
2	X	219	ALA
1	A	184	PRO
1	A	193	LEU
2	X	28	PRO
1	A	83	ALA
2	X	88	GLU
1	A	185	ILE
2	X	126	PRO
1	A	60	ILE
1	A	152	PRO

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	156/162 (96%)	141 (90%)	15 (10%)	10	28
2	X	247/247 (100%)	224 (91%)	23 (9%)	10	30
All	All	403/409 (98%)	365 (91%)	38 (9%)	10	29

All (38) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	28	THR
1	A	31	GLN
1	A	32	ASP
1	A	55	LYS
1	A	60	ILE
1	A	90	LEU
1	A	100	ASN
1	A	129	VAL
1	A	135	ASP
1	A	137	GLU
1	A	150	SER
1	A	155	SER
1	A	160	ARG
1	A	188	TRP
1	A	192	GLU
2	X	30	LEU
2	X	55	SER
2	X	60	LEU
2	X	61	ASP
2	X	63	GLU
2	X	110	LEU
2	X	122	GLU
2	X	141	MET
2	X	146	ARG
2	X	147	GLN
2	X	152	THR
2	X	160	ARG
2	X	168	LYS
2	X	173	ASN
2	X	192	TRP
2	X	207	LEU
2	X	211	LYS
2	X	212	LEU
2	X	218	GLU
2	X	224	CYS
2	X	253	GLN
2	X	263	LEU
2	X	291	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	52	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	113	HIS
1	A	120	GLN
1	A	141	HIS
2	X	78	ASN
2	X	98	HIS
2	X	113	GLN
2	X	147	GLN
2	X	173	ASN
2	X	221	GLN
2	X	228	ASN
2	X	253	GLN
2	X	255	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	NAG	A	2000	1	14,14,15	0.62	0	15,19,21	1.18	1 (6%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	NAG	A	2001	1	14,14,15	1.04	1 (7%)	15,19,21	0.83	0
3	NAG	X	1000	2	14,14,15	1.11	1 (7%)	15,19,21	0.53	0
3	NAG	X	1001	2	14,14,15	1.26	2 (14%)	15,19,21	0.77	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	2000	1	-	0/6/23/26	0/1/1/1
3	NAG	A	2001	1	-	0/6/23/26	0/1/1/1
3	NAG	X	1000	2	-	0/6/23/26	0/1/1/1
3	NAG	X	1001	2	-	0/6/23/26	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	X	1001	NAG	C3-C2	2.07	1.57	1.52
3	X	1000	NAG	C1-C2	3.44	1.57	1.52
3	A	2001	NAG	C1-C2	3.50	1.57	1.52
3	X	1001	NAG	C1-C2	3.68	1.57	1.52

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	2000	NAG	C4-C3-C2	2.10	114.10	111.02

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	2000	NAG	1	0
3	X	1000	NAG	1	0
3	X	1001	NAG	4	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	171/177 (96%)	0.09	2 (1%) 79 72	21, 50, 127, 141	0
2	X	283/283 (100%)	0.55	32 (11%) 6 3	34, 105, 147, 151	0
All	All	454/460 (98%)	0.38	34 (7%) 15 8	21, 83, 145, 151	0

All (34) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	29	LEU	5.9
2	X	53	PRO	5.4
2	X	215	ILE	4.4
2	X	159	TRP	4.1
2	X	161	GLY	4.0
2	X	99	LEU	4.0
2	X	49	GLU	3.9
2	X	134	LEU	3.9
2	X	65	PRO	3.7
2	X	52	GLY	3.6
2	X	86	GLU	3.6
2	X	105	ALA	3.3
2	X	30	LEU	3.1
2	X	40	LEU	3.0
2	X	133	ALA	2.8
2	X	19	PRO	2.8
2	X	88	GLU	2.6
2	X	54	ILE	2.6
2	X	219	ALA	2.6
2	X	128	LEU	2.5
2	X	41	ARG	2.5
2	X	35	GLY	2.3
2	X	100	TYR	2.3
2	X	126	PRO	2.2

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	X	157	SER	2.2
2	X	295	VAL	2.2
2	X	156	PHE	2.2
2	X	110	LEU	2.1
2	X	129	ILE	2.1
2	X	48	VAL	2.1
2	X	31	VAL	2.1
2	X	58	TRP	2.0
2	X	47	SER	2.0
1	A	28	THR	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	NAG	A	2001	14/15	0.82	0.29	1.53	134,141,141,141	0
3	NAG	X	1000	14/15	0.74	0.27	-0.01	148,148,148,148	0
3	NAG	A	2000	14/15	0.90	0.20	-0.07	112,112,112,112	0
3	NAG	X	1001	14/15	0.70	0.30	-	132,136,136,136	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.