



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 02:32 pm GMT

PDB ID : 5EY0  
Title : Crystal structure of CodY from Staphylococcus aureus with GTP and Ile  
Authors : Han, A.; Hwang, K.Y.  
Deposited on : 2015-11-24  
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	2696 (1.60-1.60)
Clashscore	112137	2967 (1.60-1.60)
Ramachandran outliers	110173	2887 (1.60-1.60)
Sidechain outliers	110143	2886 (1.60-1.60)
RSRZ outliers	101464	2714 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	274	<div> <div>13%</div> <div> <div></div> <div>83%</div> <div>8%</div> <div>8%</div> </div> </div>
1	B	274	<div> <div>9%</div> <div> <div></div> <div>80%</div> <div>11%</div> <div>8%</div> </div> </div>

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 4502 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GTP-sensing transcriptional pleiotropic repressor CodY.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	253	Total	C	N	O	S	0	0	0
			1991	1254	345	387	5			
1	A	252	Total	C	N	O	S	0	0	0
			1983	1248	344	386	5			

There are 34 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	-16	HIS	-	expression tag	UNP A7X1N2
B	-15	HIS	-	expression tag	UNP A7X1N2
B	-14	HIS	-	expression tag	UNP A7X1N2
B	-13	HIS	-	expression tag	UNP A7X1N2
B	-12	HIS	-	expression tag	UNP A7X1N2
B	-11	HIS	-	expression tag	UNP A7X1N2
B	-10	SER	-	expression tag	UNP A7X1N2
B	-9	SER	-	expression tag	UNP A7X1N2
B	-8	GLY	-	expression tag	UNP A7X1N2
B	-7	LEU	-	expression tag	UNP A7X1N2
B	-6	VAL	-	expression tag	UNP A7X1N2
B	-5	PRO	-	expression tag	UNP A7X1N2
B	-4	ARG	-	expression tag	UNP A7X1N2
B	-3	GLY	-	expression tag	UNP A7X1N2
B	-2	SER	-	expression tag	UNP A7X1N2
B	-1	HIS	-	expression tag	UNP A7X1N2
B	0	MET	-	expression tag	UNP A7X1N2
A	-16	HIS	-	expression tag	UNP A7X1N2
A	-15	HIS	-	expression tag	UNP A7X1N2
A	-14	HIS	-	expression tag	UNP A7X1N2
A	-13	HIS	-	expression tag	UNP A7X1N2
A	-12	HIS	-	expression tag	UNP A7X1N2
A	-11	HIS	-	expression tag	UNP A7X1N2
A	-10	SER	-	expression tag	UNP A7X1N2
A	-9	SER	-	expression tag	UNP A7X1N2

*Continued on next page...*

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
A	-8	GLY	-	expression tag	UNP A7X1N2
A	-7	LEU	-	expression tag	UNP A7X1N2
A	-6	VAL	-	expression tag	UNP A7X1N2
A	-5	PRO	-	expression tag	UNP A7X1N2
A	-4	ARG	-	expression tag	UNP A7X1N2
A	-3	GLY	-	expression tag	UNP A7X1N2
A	-2	SER	-	expression tag	UNP A7X1N2
A	-1	HIS	-	expression tag	UNP A7X1N2
A	0	MET	-	expression tag	UNP A7X1N2

- Molecule 2 is ISOLEUCINE (three-letter code: ILE) (formula:  $C_6H_{13}NO_2$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	B	1	Total	C	N	O	0	0
			9	6	1	2		
2	A	1	Total	C	N	O	0	0
			9	6	1	2		

- Molecule 3 is GUANOSINE-5'-TRIPHOSPHATE (three-letter code: GTP) (formula:  $C_{10}H_{16}N_5O_{14}P_3$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	B	1	Total	C	N	O	P	0	0
			32	10	5	14	3		
3	A	1	Total	C	N	O	P	0	0
			32	10	5	14	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	211	Total	O	0	0
			211	211		
4	A	235	Total	O	0	0
			235	235		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: GTP-sensing transcriptional pleiotropic repressor CodY



- Molecule 1: GTP-sensing transcriptional pleiotropic repressor CodY



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	45.89Å 76.03Å 158.02Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.94 – 1.60 29.94 – 1.60	Depositor EDS
% Data completeness (in resolution range)	94.0 (29.94-1.60) 94.1 (29.94-1.60)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.15 (at 1.60Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.9_1692)	Depositor
R, $R_{free}$	0.178 , 0.209 0.177 , 0.209	Depositor DCC
$R_{free}$ test set	3416 reflections (4.91%)	DCC
Wilson B-factor (Å <sup>2</sup> )	15.0	Xtriage
Anisotropy	0.389	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 55.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	4502	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.52% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GTP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.36	0/2002	0.52	0/2689
1	B	0.35	0/2010	0.52	1/2700 (0.0%)
All	All	0.35	0/4012	0.52	1/5389 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	235	LEU	CA-CB-CG	5.16	127.17	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1983	0	2056	19	0
1	B	1991	0	2067	23	0
2	A	9	0	10	0	0
2	B	9	0	10	0	0
3	A	32	0	12	0	0
3	B	32	0	12	0	0
4	A	235	0	0	5	3
4	B	211	0	0	7	1

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	4502	0	4167	41	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:132:HIS:O	4:A:401:HOH:O	1.93	0.84
1:B:231:GLU:HG2	1:B:245:LYS:HE2	1.60	0.82
1:B:138:ASN:OD1	4:B:402:HOH:O	2.00	0.80
1:A:104:GLU:OE2	4:A:402:HOH:O	2.03	0.76
1:B:241:PHE:HE2	1:B:243:LYS:HD3	1.51	0.74
1:B:231:GLU:CG	1:B:245:LYS:HE2	2.23	0.67
1:B:26:ASP:OD2	4:B:403:HOH:O	2.13	0.67
1:B:61:ARG:HD3	1:B:101:GLU:OE2	1.98	0.63
1:A:37:THR:HG22	1:A:130:ARG:HG3	1.81	0.62
1:A:58:LYS:NZ	4:A:408:HOH:O	2.33	0.62
1:B:190:ILE:HD11	1:B:217:ILE:HG23	1.80	0.62
1:A:60:GLN:HG3	4:A:414:HOH:O	1.98	0.62
1:B:37:THR:HG22	1:B:130:ARG:HG3	1.83	0.59
1:B:251:ASP:OD2	4:B:404:HOH:O	2.17	0.58
1:A:61:ARG:HD3	1:A:101:GLU:OE2	2.04	0.57
1:B:183:GLU:OE2	1:B:219:ASN:ND2	2.32	0.55
1:A:231:GLU:HG3	1:A:243:LYS:HB3	1.89	0.55
1:A:246:LYS:O	1:A:249:PHE:N	2.41	0.54
1:A:1:MET:HE3	4:A:520:HOH:O	2.08	0.53
1:B:81:GLU:HB2	4:B:422:HOH:O	2.08	0.52
1:B:19:GLY:O	4:B:405:HOH:O	2.20	0.48
1:B:190:ILE:HG23	1:B:201:LEU:HD13	1.95	0.47
1:B:232:SER:HB3	1:B:242:ILE:HG22	1.97	0.47
1:B:234:SER:OG	1:B:234:SER:O	2.19	0.47
1:A:198:GLU:CD	1:A:243:LYS:HE3	2.35	0.46
1:A:80:MET:HE2	1:A:80:MET:HA	1.97	0.46
1:B:205:LYS:CE	1:B:209:ARG:HH22	2.29	0.46
1:B:78:ARG:NH2	4:B:413:HOH:O	2.49	0.46
1:A:247:GLU:C	1:A:249:PHE:H	2.20	0.45
1:B:194:LEU:HD23	1:B:194:LEU:HA	1.86	0.44
1:A:189:HIS:O	1:A:193:GLU:HB3	2.18	0.43
1:B:1:MET:N	4:B:401:HOH:O	1.89	0.42
1:B:207:ALA:HB2	1:B:217:ILE:HD12	2.02	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:222:ARG:HH12	1:A:177:ASN:ND2	2.18	0.42
1:B:44:ARG:HD3	1:B:122:ARG:O	2.20	0.42
1:A:117:LEU:HD23	1:A:122:ARG:HA	2.03	0.41
1:A:250:LEU:HD23	1:A:250:LEU:HA	1.90	0.41
1:A:233:ARG:HG2	1:A:234:SER:N	2.36	0.41
1:A:231:GLU:HG3	1:A:243:LYS:CB	2.51	0.41
1:B:202:ILE:HD13	1:B:205:LYS:HD3	2.02	0.40
1:A:247:GLU:C	1:A:249:PHE:N	2.74	0.40

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:420:HOH:O	4:A:591:HOH:O[4_545]	2.05	0.15
4:B:527:HOH:O	4:A:508:HOH:O[1_655]	2.12	0.08
4:A:402:HOH:O	4:A:420:HOH:O[4_445]	2.19	0.01

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	248/274 (90%)	240 (97%)	7 (3%)	1 (0%)	38	16
1	B	249/274 (91%)	243 (98%)	6 (2%)	0	100	100
All	All	497/548 (91%)	483 (97%)	13 (3%)	1 (0%)	51	27

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	252	GLU

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	223/242 (92%)	220 (99%)	3 (1%)	73	55
1	B	224/242 (93%)	221 (99%)	3 (1%)	73	55
All	All	447/484 (92%)	441 (99%)	6 (1%)	73	55

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	208	ASP
1	B	234	SER
1	B	235	LEU
1	A	193	GLU
1	A	231	GLU
1	A	243	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	177	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	ILE	A	301	-	5,8,8	0.39	0	4,10,10	0.17	0
3	GTP	A	302	-	27,34,34	0.94	1 (3%)	27,54,54	1.68	5 (18%)
2	ILE	B	301	-	5,8,8	0.44	0	4,10,10	0.25	0
3	GTP	B	302	-	27,34,34	0.94	1 (3%)	27,54,54	1.81	5 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ILE	A	301	-	-	0/6/10/10	0/0/0/0
3	GTP	A	302	-	-	0/18/38/38	0/3/3/3
2	ILE	B	301	-	-	0/6/10/10	0/0/0/0
3	GTP	B	302	-	-	0/18/38/38	0/3/3/3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	302	GTP	C6-N1	3.21	1.38	1.33
3	A	302	GTP	C6-N1	3.24	1.38	1.33

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	302	GTP	N3-C2-N1	-5.29	119.73	127.46
3	A	302	GTP	N3-C2-N1	-4.69	120.61	127.46
3	B	302	GTP	C5-C6-N1	-3.44	118.59	123.48
3	A	302	GTP	C5-C6-N1	-3.19	118.94	123.48
3	B	302	GTP	C4-C5-N7	-2.16	107.32	109.41

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	302	GTP	C4-C5-N7	-2.13	107.35	109.41
3	A	302	GTP	C6-N1-C2	2.90	120.23	116.06
3	B	302	GTP	C6-N1-C2	3.60	121.24	116.06
3	B	302	GTP	C2-N3-C4	3.78	119.58	115.16
3	A	302	GTP	C2-N3-C4	3.97	119.80	115.16

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	252/274 (91%)	0.60	35 (13%) <b>3</b> <b>3</b>	10, 25, 96, 114	0
1	B	253/274 (92%)	0.42	24 (9%) <b>9</b> <b>8</b>	11, 26, 82, 115	0
All	All	505/548 (92%)	0.51	59 (11%) <b>5</b> <b>5</b>	10, 25, 94, 115	0

All (59) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	195	GLY	11.6
1	A	196	GLY	9.2
1	A	200	LEU	8.7
1	A	241	PHE	8.6
1	B	197	THR	8.3
1	A	195	GLY	7.3
1	A	244	VAL	6.4
1	A	197	THR	6.0
1	A	249	PHE	6.0
1	B	235	LEU	5.9
1	A	233	ARG	5.8
1	A	181	TYR	5.8
1	A	247	GLU	5.5
1	A	242	ILE	5.2
1	B	241	PHE	5.2
1	B	196	GLY	5.1
1	B	233	ARG	4.9
1	B	234	SER	4.8
1	A	192	GLU	4.8
1	B	239	GLY	4.8
1	A	239	GLY	4.8
1	A	234	SER	4.6
1	A	254	GLU	4.6
1	A	232	SER	4.5

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	200	LEU	4.5
1	A	250	LEU	4.5
1	A	251	ASP	4.4
1	A	238	LYS	4.3
1	B	247	GLU	4.3
1	A	240	THR	4.3
1	A	194	LEU	4.2
1	B	198	GLU	4.2
1	A	202	ILE	4.0
1	A	248	LYS	3.9
1	B	245	LYS	3.8
1	B	238	LYS	3.7
1	B	243	LYS	3.6
1	A	201	LEU	3.5
1	B	228	GLY	3.4
1	A	191	PHE	3.4
1	B	231	GLU	3.4
1	A	198	GLU	3.2
1	B	229	VAL	3.1
1	A	246	LYS	3.0
1	B	232	SER	2.9
1	A	199	GLY	2.9
1	A	231	GLU	2.8
1	A	243	LYS	2.8
1	A	210	VAL	2.7
1	B	251	ASP	2.7
1	B	242	ILE	2.6
1	A	211	GLY	2.4
1	B	254	GLU	2.4
1	B	248	LYS	2.4
1	A	186	ALA	2.2
1	A	255	LYS	2.1
1	A	204	SER	2.1
1	B	162	VAL	2.1
1	B	250	LEU	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	ILE	B	301	9/9	0.96	0.09	0.76	16,20,22,22	0
2	ILE	A	301	9/9	0.96	0.08	-0.16	14,19,20,23	0
3	GTP	A	302	32/32	0.94	0.09	-0.18	10,21,55,61	0
3	GTP	B	302	32/32	0.96	0.07	-0.29	16,22,47,61	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.