



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 14, 2017 – 03:01 am GMT

PDB ID : 2F2U  
Title : crystal structure of the Rho-kinase kinase domain  
Authors : Yamaguchi, H.; Hakoshima, T.  
Deposited on : 2005-11-18  
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	3166 (2.40-2.40)
Clashscore	112137	3674 (2.40-2.40)
Ramachandran outliers	110173	3616 (2.40-2.40)
Sidechain outliers	110143	3617 (2.40-2.40)
RSRZ outliers	101464	3195 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	402	<div> <div>4%</div> <div> <div></div> <div>78%</div> <div>16%</div> <div>• •</div> </div> </div>
1	B	402	<div> <div>5%</div> <div> <div></div> <div>73%</div> <div>20%</div> <div>• 5%</div> </div> </div>

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 6413 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Rho-associated protein kinase 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	386	Total	C	N	O	S	0	0	0
			3118	2001	518	578	21			
1	B	383	Total	C	N	O	S	0	0	0
			3093	1986	515	571	21			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	16	GLY	-	CLONING ARTIFACT	UNP Q28021
A	17	ALA	-	CLONING ARTIFACT	UNP Q28021
A	52	SER	PRO	SEE REMARK 999	UNP Q28021
B	16	GLY	-	CLONING ARTIFACT	UNP Q28021
B	17	ALA	-	CLONING ARTIFACT	UNP Q28021
B	52	SER	PRO	SEE REMARK 999	UNP Q28021

- Molecule 2 is 5-(1,4-DIAZEPAN-1-SULFONYL)ISOQUINOLINE (three-letter code: M77) (formula: C<sub>14</sub>H<sub>17</sub>N<sub>3</sub>O<sub>2</sub>S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			20	14	3	2	1		
2	B	1	Total	C	N	O	S	0	0
			20	14	3	2	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	75	Total	O	0	0
			75	75		
3	B	87	Total	O	0	0
			87	87		


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Rho-associated protein kinase 2



#### • Molecule 1: Rho-associated protein kinase 2



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	102.53Å 102.53Å 257.19Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 2.40 29.86 – 2.40	Depositor EDS
% Data completeness (in resolution range)	95.9 (30.00-2.40) 95.9 (29.86-2.40)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.07	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.93 (at 2.39Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.195 , 0.235 0.194 , 0.234	Depositor DCC
$R_{free}$ test set	2662 reflections (5.35%)	DCC
Wilson B-factor (Å <sup>2</sup> )	42.2	Xtriage
Anisotropy	0.101	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 40.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	6413	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	47.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 13.75% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: M77

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.60	0/3194	0.81	1/4315 (0.0%)
1	B	0.64	0/3169	0.80	1/4281 (0.0%)
All	All	0.62	0/6363	0.80	2/8596 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	365	ARG	NE-CZ-NH2	-5.62	117.49	120.30
1	B	142	ASP	CB-CG-OD1	5.22	123.00	118.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3118	0	3051	52	0
1	B	3093	0	3030	70	0
2	A	20	0	17	0	0
2	B	20	0	17	1	0
3	A	75	0	0	3	0
3	B	87	0	0	9	0
All	All	6413	0	6115	122	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (122) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:126:PHE:HD1	1:B:130:LYS:HZ2	0.99	0.93
1:B:60:PRO:HA	3:B:1568:HOH:O	1.69	0.93
1:B:342:ARG:HG2	3:B:1579:HOH:O	1.78	0.83
1:B:239:MET:HE3	1:B:243:GLY:O	1.79	0.82
1:A:240:ASP:HB3	1:A:242:THR:H	1.47	0.78
1:B:32:ALA:HA	1:B:35:ARG:HH12	1.48	0.78
1:A:151:TRP:CH2	1:A:365:ARG:HG3	2.18	0.78
1:A:121:LYS:HD2	1:A:167:MET:CE	2.17	0.74
1:B:126:PHE:O	1:B:130:LYS:HG2	1.88	0.72
1:A:41:ILE:O	1:A:41:ILE:HD12	1.91	0.70
1:B:30:LEU:HD12	3:B:1531:HOH:O	1.90	0.70
1:A:72:LEU:O	1:A:76:GLU:HB2	1.91	0.69
1:B:181:MET:HE2	1:B:286:GLU:HG2	1.75	0.68
1:A:317:GLU:CD	1:A:317:GLU:H	1.97	0.67
1:B:417:ARG:HG3	1:B:417:ARG:HH11	1.61	0.65
1:B:181:MET:HE3	1:B:287:MET:HA	1.79	0.64
1:B:263:LYS:HB3	3:B:1582:HOH:O	1.99	0.63
1:A:256:TYR:OH	1:A:286:GLU:OE1	2.15	0.63
1:B:330:CYS:O	1:B:334:THR:HG23	1.98	0.63
1:A:360:ASN:H	1:A:363:ASN:HD21	1.46	0.62
1:A:28:ARG:H	1:A:28:ARG:HD2	1.63	0.61
1:B:239:MET:HE1	1:B:273:ARG:HE	1.64	0.61
1:B:32:ALA:HA	1:B:35:ARG:NH1	2.14	0.60
1:A:76:GLU:O	1:A:80:LYS:HG2	2.01	0.59
1:B:126:PHE:HD1	1:B:130:LYS:NZ	1.86	0.59
1:B:35:ARG:HB2	1:B:35:ARG:HH11	1.67	0.59
1:A:148:ASN:HB3	3:A:552:HOH:O	2.02	0.59
1:B:27:GLN:HA	1:B:30:LEU:HD21	1.85	0.59
1:A:240:ASP:HB2	1:A:244:MET:H	1.67	0.58
1:A:209:GLY:HA2	1:A:239:MET:CE	2.34	0.58
1:A:248:ASP:OD2	1:A:267:GLY:HA2	2.04	0.57
1:A:209:GLY:HA2	1:A:239:MET:HE3	1.86	0.57
1:B:256:TYR:OH	1:B:286:GLU:OE1	2.23	0.57
1:A:363:ASN:C	1:A:363:ASN:HD22	2.08	0.56
1:B:62:LEU:HD11	1:B:403:PHE:N	2.21	0.56
1:B:268:ASP:HB3	3:B:1582:HOH:O	2.06	0.56
1:B:172:MET:SD	1:B:229:LYS:HD2	2.45	0.56

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:282:VAL:HG13	1:B:293:PRO:HD2	1.87	0.56
1:A:28:ARG:HH11	1:A:28:ARG:N	2.05	0.55
1:A:111:HIS:HE1	1:A:113:ALA:HB3	1.72	0.55
1:B:66:LYS:O	1:B:70:ASN:ND2	2.39	0.55
1:B:181:MET:CE	1:B:286:GLU:HG2	2.36	0.55
1:A:56:ASP:OD2	1:A:405:GLY:HA2	2.07	0.54
1:B:317:GLU:CD	1:B:317:GLU:H	2.11	0.54
1:B:220:MET:HB3	1:B:228:LEU:HD11	1.90	0.53
1:B:35:ARG:HB2	1:B:35:ARG:NH1	2.23	0.53
1:A:211:ILE:HG13	1:A:213:ARG:HD2	1.89	0.53
1:A:114:SER:O	1:A:115:GLN:HB2	2.09	0.53
1:B:250:ALA:HA	1:B:265:GLN:HE22	1.72	0.53
1:B:127:GLU:O	1:B:131:ARG:HB3	2.09	0.53
1:B:336:ARG:HD3	1:B:342:ARG:HH21	1.74	0.53
1:A:360:ASN:H	1:A:363:ASN:ND2	2.08	0.52
1:A:72:LEU:O	1:A:76:GLU:CB	2.58	0.52
1:A:29:LYS:O	1:A:32:ALA:HB3	2.09	0.52
1:A:70:ASN:O	1:A:73:ASN:HB2	2.10	0.52
1:A:70:ASN:CG	1:A:71:PHE:N	2.63	0.52
1:A:62:LEU:HD21	1:A:403:PHE:N	2.24	0.52
1:B:129:ILE:HG21	1:B:397:PHE:HE1	1.74	0.52
1:A:95:VAL:O	1:A:96:LYS:HB3	2.08	0.51
1:B:103:PHE:CZ	2:B:1501:M77:H211	2.45	0.51
1:B:30:LEU:C	1:B:32:ALA:N	2.61	0.51
1:B:141:ARG:HG3	1:B:142:ASP:N	2.25	0.50
1:B:111:HIS:HE1	1:B:113:ALA:HB3	1.76	0.49
1:B:28:ARG:H	1:B:29:LYS:NZ	2.09	0.49
1:A:264:SER:HB2	1:A:269:GLY:HA3	1.93	0.49
1:A:145:ALA:HB2	1:A:155:LEU:HD23	1.94	0.49
1:B:299:LEU:C	1:B:299:LEU:HD23	2.33	0.49
1:A:363:ASN:C	1:A:363:ASN:ND2	2.66	0.49
1:B:181:MET:CE	1:B:287:MET:HA	2.42	0.49
1:A:225:HIS:HD2	3:A:518:HOH:O	1.95	0.49
1:B:318:ASP:N	1:B:318:ASP:OD1	2.47	0.48
1:B:417:ARG:HG3	1:B:417:ARG:NH1	2.28	0.48
1:B:111:HIS:CE1	1:B:113:ALA:HB3	2.49	0.48
1:A:401:LYS:HA	1:A:401:LYS:HE2	1.95	0.47
1:B:70:ASN:O	1:B:74:ARG:HG3	2.12	0.47
1:B:214:ASP:OD1	1:B:216:LYS:HE3	2.14	0.47
1:B:30:LEU:C	1:B:32:ALA:H	2.17	0.47
1:B:363:ASN:O	1:B:367:THR:HG23	2.14	0.47

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:239:MET:CE	3:B:1516:HOH:O	2.61	0.47
1:A:130:LYS:HD3	1:A:131:ARG:N	2.30	0.47
1:A:178:VAL:HG23	1:A:217:PRO:HB2	1.98	0.46
1:B:216:LYS:HB2	1:B:217:PRO:HD2	1.97	0.46
1:B:129:ILE:HA	1:B:129:ILE:HD13	1.78	0.46
1:B:128:MET:O	1:B:134:SER:OG	2.32	0.45
1:B:95:VAL:O	1:B:96:LYS:HB3	2.16	0.45
1:B:237:MET:SD	1:B:247:CYS:HB2	2.56	0.45
1:A:321:ILE:HG22	1:A:325:ALA:HB3	1.98	0.45
1:B:30:LEU:H	1:B:30:LEU:HD23	1.82	0.45
1:A:317:GLU:CD	1:A:317:GLU:N	2.68	0.45
1:B:36:ASP:HA	1:B:37:PRO:HD3	1.77	0.44
1:B:356:ASN:HB2	1:B:358:GLN:HE21	1.82	0.44
1:A:27:GLN:N	1:A:28:ARG:HD2	2.33	0.44
1:A:100:ARG:NH1	1:A:387:ILE:HG23	2.32	0.44
1:B:128:MET:HB3	1:B:133:ASP:O	2.16	0.44
1:A:292:THR:HB	3:A:502:HOH:O	2.18	0.44
1:B:114:SER:O	1:B:115:GLN:HB2	2.18	0.43
1:B:128:MET:SD	1:B:136:PHE:CE1	3.11	0.43
1:A:111:HIS:CE1	1:A:113:ALA:HB3	2.52	0.43
1:B:124:SER:HB3	1:B:127:GLU:HB2	2.00	0.43
1:B:143:ILE:HD11	1:B:210:LEU:HD13	2.00	0.43
1:A:28:ARG:H	1:A:28:ARG:CD	2.28	0.43
1:A:399:ILE:H	1:A:399:ILE:HD12	1.84	0.43
1:A:277:TRP:CE3	1:A:277:TRP:HA	2.54	0.42
1:A:121:LYS:HD2	1:A:167:MET:HE2	2.00	0.42
1:A:36:ASP:HA	1:A:37:PRO:HD3	1.85	0.42
1:B:136:PHE:HA	3:B:1539:HOH:O	2.19	0.42
1:B:145:ALA:HB2	1:B:155:LEU:HD23	2.02	0.42
1:B:239:MET:O	3:B:1555:HOH:O	2.22	0.41
1:B:251:VAL:H	1:B:265:GLN:NE2	2.17	0.41
1:B:156:PHE:CE2	1:B:170:GLU:HB3	2.55	0.41
1:B:355:LYS:HA	1:B:355:LYS:HD3	1.78	0.41
1:B:39:SER:HA	1:B:40:PRO:HD3	1.86	0.41
1:B:239:MET:HE1	3:B:1516:HOH:O	2.19	0.41
1:A:34:ILE:HG23	1:A:43:VAL:HG22	2.02	0.41
1:A:330:CYS:O	1:A:334:THR:HG23	2.20	0.41
1:B:65:ASN:OD1	1:B:65:ASN:C	2.60	0.41
1:A:277:TRP:HE3	1:A:277:TRP:HA	1.86	0.41
1:B:28:ARG:H	1:B:29:LYS:HZ2	1.68	0.41
1:A:363:ASN:HD22	1:A:364:ILE:N	2.19	0.40

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:127:GLU:OE1	1:A:131:ARG:NH1	2.51	0.40
1:A:70:ASN:OD1	1:A:71:PHE:N	2.54	0.40
1:B:41:ILE:HD12	1:B:41:ILE:C	2.42	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	382/402 (95%)	369 (97%)	11 (3%)	2 (0%)	32	46
1	B	379/402 (94%)	359 (95%)	16 (4%)	4 (1%)	17	23
All	All	761/804 (95%)	728 (96%)	27 (4%)	6 (1%)	22	33

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	29	LYS
1	B	30	LEU
1	B	135	ALA
1	A	96	LYS
1	A	240	ASP
1	B	232	ASP

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	340/348 (98%)	324 (95%)	16 (5%)	30	48
1	B	337/348 (97%)	323 (96%)	14 (4%)	34	53
All	All	677/696 (97%)	647 (96%)	30 (4%)	33	51

All (30) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	28	ARG
1	A	29	LYS
1	A	63	ARG
1	A	70	ASN
1	A	105	GLU
1	A	130	LYS
1	A	131	ARG
1	A	142	ASP
1	A	163	LYS
1	A	165	LEU
1	A	185	ASP
1	A	218	ASP
1	A	237	MET
1	A	363	ASN
1	A	365	ARG
1	A	400	PRO
1	B	29	LYS
1	B	30	LEU
1	B	90	GLU
1	B	105	GLU
1	B	126	PHE
1	B	131	ARG
1	B	141	ARG
1	B	213	ARG
1	B	255	ASP
1	B	318	ASP
1	B	336	ARG
1	B	345	VAL
1	B	400	PRO
1	B	401	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	27	GLN
1	A	107	GLN
1	A	183	ASN
1	A	225	HIS
1	A	350	GLN
1	A	363	ASN
1	B	70	ASN
1	B	107	GLN
1	B	183	ASN
1	B	265	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	M77	A	501	-	20,22,22	2.11	8 (40%)	29,31,31	2.77	9 (31%)
2	M77	B	1501	-	20,22,22	2.38	9 (45%)	29,31,31	1.86	7 (24%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	M77	A	501	-	-	0/12/21/21	0/2/3/3
2	M77	B	1501	-	-	0/12/21/21	0/2/3/3

All (17) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1501	M77	S1-N4	-2.47	1.59	1.63
2	A	501	M77	S1-N4	-2.33	1.60	1.63
2	B	1501	M77	O1-S1	-2.08	1.41	1.43
2	A	501	M77	C7-C6	2.08	1.42	1.38
2	A	501	M77	C7-C8	2.09	1.41	1.36
2	B	1501	M77	C15-N4	2.24	1.50	1.47
2	A	501	M77	C15-N4	2.55	1.50	1.47
2	B	1501	M77	C7-C8	2.60	1.42	1.36
2	B	1501	M77	C7-C6	2.83	1.44	1.38
2	A	501	M77	C14-N13	2.89	1.37	1.32
2	B	1501	M77	C14-N13	3.36	1.38	1.32
2	A	501	M77	C10-C9	3.75	1.49	1.43
2	A	501	M77	C6-C5	3.80	1.42	1.37
2	B	1501	M77	C10-C9	3.96	1.50	1.43
2	B	1501	M77	C11-C12	4.13	1.42	1.36
2	A	501	M77	C11-C12	4.73	1.43	1.36
2	B	1501	M77	C6-C5	5.07	1.43	1.37

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	M77	O2-S1-O1	-4.84	111.32	119.46
2	A	501	M77	C6-C5-C10	-4.12	116.92	120.98
2	B	1501	M77	O2-S1-O1	-3.10	114.25	119.46
2	B	1501	M77	C6-C5-C10	-2.83	118.19	120.98
2	A	501	M77	C11-C12-N13	-2.29	120.68	123.73
2	B	1501	M77	C5-C10-C9	2.01	119.38	117.47
2	A	501	M77	C8-C9-C10	2.08	121.81	119.11
2	B	1501	M77	C12-N13-C14	2.09	122.24	117.27
2	A	501	M77	C15-N4-C22	2.36	122.88	115.19
2	B	1501	M77	O1-S1-N4	2.46	109.01	106.69
2	A	501	M77	C7-C6-C5	2.64	123.09	120.37
2	B	1501	M77	C22-C21-N17	3.70	118.08	112.87

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	M77	C21-C22-N4	4.56	119.61	112.27
2	B	1501	M77	O2-S1-N4	5.31	111.70	106.69
2	A	501	M77	C22-C21-N17	5.44	120.53	112.87
2	A	501	M77	O2-S1-N4	8.96	115.15	106.69

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1501	M77	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	386/402 (96%)	-0.08	16 (4%) 38 36	25, 43, 82, 95	0
1	B	383/402 (95%)	-0.06	21 (5%) 26 24	27, 42, 80, 95	0
All	All	769/804 (95%)	-0.07	37 (4%) 31 30	25, 42, 81, 95	0

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	394	VAL	4.3
1	A	268	ASP	4.3
1	B	27	GLN	4.2
1	B	35	ARG	4.0
1	B	318	ASP	3.8
1	B	133	ASP	3.6
1	A	395	GLU	3.4
1	A	29	LYS	3.2
1	A	64	LYS	3.1
1	B	132	SER	3.1
1	A	52	SER	3.0
1	A	49	GLY	3.0
1	A	27	GLN	2.9
1	B	131	ARG	2.9
1	B	134	SER	2.9
1	B	130	LYS	2.8
1	A	317	GLU	2.8
1	A	130	LYS	2.8
1	A	318	ASP	2.8
1	B	49	GLY	2.7
1	B	28	ARG	2.7
1	A	132	SER	2.6
1	B	30	LEU	2.5
1	B	52	SER	2.5

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	388	GLU	2.5
1	B	280	VAL	2.4
1	B	102	ALA	2.3
1	B	396	THR	2.3
1	B	64	LYS	2.3
1	A	396	THR	2.3
1	B	126	PHE	2.3
1	B	317	GLU	2.3
1	A	77	LYS	2.2
1	B	85	LEU	2.1
1	A	401	LYS	2.1
1	B	248	ASP	2.1
1	B	279	SER	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	M77	B	1501	20/20	0.97	0.14	0.13	31,39,54,56	0
2	M77	A	501	20/20	0.98	0.11	-0.56	27,36,55,56	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.