



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 14, 2017 – 01:31 pm GMT

PDB ID : 3F2I
Title : Crystal structure of the alr0221 protein from Nostoc, Northeast Structural Genomics Consortium Target NsR422.
Authors : Forouhar, F.; Lew, S.; Seetharaman, J.; Sahdev, S.; Xiao, R.; Foote, E.L.; Ciccocanti, C.; Belote, R.L.; Nair, R.; Everett, J.K.; Acton, T.B.; Rost, B.; Montelione, G.T.; Tong, L.; Hunt, J.F.; Northeast Structural Genomics Consortium (NESG)
Deposited on : 2008-10-29
Resolution : 2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	6609 (2.00-2.00)
Clashscore	112137	7775 (2.00-2.00)
Ramachandran outliers	110173	7679 (2.00-2.00)
Sidechain outliers	110143	7678 (2.00-2.00)
RSRZ outliers	101464	6696 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	172	<div> <div>10%</div> <div> <div></div> <div>72%</div> <div>20%</div> <div>• 6%</div> </div> </div>
1	B	172	<div> <div>3%</div> <div> <div></div> <div>73%</div> <div>22%</div> <div>• 5%</div> </div> </div>
1	C	172	<div> <div>4%</div> <div> <div></div> <div>65%</div> <div>27%</div> <div>• 6%</div> </div> </div>
1	D	172	<div> <div>2%</div> <div> <div></div> <div>71%</div> <div>24%</div> <div>• •</div> </div> </div>
1	E	172	<div> <div>2%</div> <div> <div></div> <div>75%</div> <div>20%</div> <div>• 5%</div> </div> </div>
1	F	172	<div> <div>4%</div> <div> <div></div> <div>69%</div> <div>28%</div> <div>• •</div> </div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 8330 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Alr0221 protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	162	Total	C	N	O	S	Se	0	0	0
			1300	839	223	233	2	3			
1	B	163	Total	C	N	O	S	Se	0	0	0
			1311	844	226	236	2	3			
1	C	161	Total	C	N	O	S	Se	0	0	0
			1292	833	222	232	2	3			
1	D	166	Total	C	N	O	S	Se	0	0	0
			1329	856	228	240	2	3			
1	E	164	Total	C	N	O	S	Se	0	0	0
			1312	845	226	236	2	3			
1	F	167	Total	C	N	O	S	Se	0	0	0
			1339	862	231	241	2	3			

There are 48 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	165	LEU	-	expression tag	UNP Q8Z077
A	166	GLU	-	expression tag	UNP Q8Z077
A	167	HIS	-	expression tag	UNP Q8Z077
A	168	HIS	-	expression tag	UNP Q8Z077
A	169	HIS	-	expression tag	UNP Q8Z077
A	170	HIS	-	expression tag	UNP Q8Z077
A	171	HIS	-	expression tag	UNP Q8Z077
A	172	HIS	-	expression tag	UNP Q8Z077
B	165	LEU	-	expression tag	UNP Q8Z077
B	166	GLU	-	expression tag	UNP Q8Z077
B	167	HIS	-	expression tag	UNP Q8Z077
B	168	HIS	-	expression tag	UNP Q8Z077
B	169	HIS	-	expression tag	UNP Q8Z077
B	170	HIS	-	expression tag	UNP Q8Z077
B	171	HIS	-	expression tag	UNP Q8Z077
B	172	HIS	-	expression tag	UNP Q8Z077
C	165	LEU	-	expression tag	UNP Q8Z077

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	166	GLU	-	expression tag	UNP Q8Z077
C	167	HIS	-	expression tag	UNP Q8Z077
C	168	HIS	-	expression tag	UNP Q8Z077
C	169	HIS	-	expression tag	UNP Q8Z077
C	170	HIS	-	expression tag	UNP Q8Z077
C	171	HIS	-	expression tag	UNP Q8Z077
C	172	HIS	-	expression tag	UNP Q8Z077
D	165	LEU	-	expression tag	UNP Q8Z077
D	166	GLU	-	expression tag	UNP Q8Z077
D	167	HIS	-	expression tag	UNP Q8Z077
D	168	HIS	-	expression tag	UNP Q8Z077
D	169	HIS	-	expression tag	UNP Q8Z077
D	170	HIS	-	expression tag	UNP Q8Z077
D	171	HIS	-	expression tag	UNP Q8Z077
D	172	HIS	-	expression tag	UNP Q8Z077
E	165	LEU	-	expression tag	UNP Q8Z077
E	166	GLU	-	expression tag	UNP Q8Z077
E	167	HIS	-	expression tag	UNP Q8Z077
E	168	HIS	-	expression tag	UNP Q8Z077
E	169	HIS	-	expression tag	UNP Q8Z077
E	170	HIS	-	expression tag	UNP Q8Z077
E	171	HIS	-	expression tag	UNP Q8Z077
E	172	HIS	-	expression tag	UNP Q8Z077
F	165	LEU	-	expression tag	UNP Q8Z077
F	166	GLU	-	expression tag	UNP Q8Z077
F	167	HIS	-	expression tag	UNP Q8Z077
F	168	HIS	-	expression tag	UNP Q8Z077
F	169	HIS	-	expression tag	UNP Q8Z077
F	170	HIS	-	expression tag	UNP Q8Z077
F	171	HIS	-	expression tag	UNP Q8Z077
F	172	HIS	-	expression tag	UNP Q8Z077

- Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	P	0	0
			5	4	1		
2	B	1	Total	O	P	0	0
			5	4	1		
2	C	1	Total	O	P	0	0
			5	4	1		
2	D	1	Total	O	P	0	0
			5	4	1		
2	E	1	Total	O	P	0	0
			5	4	1		
2	F	1	Total	O	P	0	0
			5	4	1		

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	D	1	Total	Cl	0	0
			1	1		
3	E	1	Total	Cl	0	0
			1	1		
3	B	1	Total	Cl	0	0
			1	1		
3	C	1	Total	Cl	0	0
			1	1		
3	A	1	Total	Cl	0	0
			1	1		
3	F	1	Total	Cl	0	0
			1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	37	Total 37	O 37	0	0
4	B	71	Total 71	O 71	0	0
4	C	62	Total 62	O 62	0	0
4	D	76	Total 76	O 76	0	0
4	E	82	Total 82	O 82	0	0
4	F	83	Total 83	O 83	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Alr0221 protein




• Molecule 1: Alr0221 protein



• Molecule 1: Alr0221 protein





• Molecule 1: Alr0221 protein






● Molecule 1: Alr0221 protein



● Molecule 1: Alr0221 protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	65.10Å 94.33Å 92.06Å 90.00° 93.64° 90.00°	Depositor
Resolution (Å)	19.93 – 2.00 29.67 – 1.99	Depositor EDS
% Data completeness (in resolution range)	85.5 (19.93-2.00) 97.4 (29.67-1.99)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.27 (at 2.00Å)	Xtriage
Refinement program	CNS 1.2 & XtalView	Depositor
R, R_{free}	0.196 , 0.240 0.215 , 0.258	Depositor DCC
R_{free} test set	3663 reflections (4.98%)	DCC
Wilson B-factor (Å ²)	17.2	Xtriage
Anisotropy	0.541	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 59.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	8330	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 32.61 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 9.1426e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.30	0/1326	0.52	0/1791
1	B	0.33	0/1338	0.55	0/1807
1	C	0.31	0/1318	0.54	0/1780
1	D	0.35	0/1356	0.58	0/1832
1	E	0.32	0/1339	0.55	0/1809
1	F	0.33	0/1367	0.56	0/1847
All	All	0.33	0/8044	0.55	0/10866

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1300	0	1334	40	0
1	B	1311	0	1336	40	0
1	C	1292	0	1323	41	0
1	D	1329	0	1364	41	0
1	E	1312	0	1347	25	0
1	F	1339	0	1371	44	0
2	A	5	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	5	0	0	0	0
2	C	5	0	0	0	0
2	D	5	0	0	0	0
2	E	5	0	0	0	0
2	F	5	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	1	0
3	E	1	0	0	1	0
3	F	1	0	0	0	0
4	A	37	0	0	2	0
4	B	71	0	0	4	0
4	C	62	0	0	2	0
4	D	76	0	0	1	0
4	E	82	0	0	2	0
4	F	83	0	0	4	0
All	All	8330	0	8075	221	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

The worst 5 of 221 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:4:TYR:HB3	1:D:137:MSE:HE2	1.26	1.15
1:B:4:TYR:HB3	1:B:137:MSE:HE2	1.20	1.11
1:F:4:TYR:HB3	1:F:137:MSE:HE2	1.27	1.09
1:F:6:ILE:HD11	1:F:137:MSE:HE3	1.43	0.99
1:A:118:THR:HG21	1:A:154:MSE:HE3	1.42	0.99

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	158/172 (92%)	155 (98%)	3 (2%)	0	100	100
1	B	159/172 (92%)	156 (98%)	3 (2%)	0	100	100
1	C	157/172 (91%)	153 (98%)	4 (2%)	0	100	100
1	D	164/172 (95%)	161 (98%)	3 (2%)	0	100	100
1	E	162/172 (94%)	160 (99%)	2 (1%)	0	100	100
1	F	165/172 (96%)	163 (99%)	2 (1%)	0	100	100
All	All	965/1032 (94%)	948 (98%)	17 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	140/146 (96%)	134 (96%)	6 (4%)	33	29
1	B	141/146 (97%)	140 (99%)	1 (1%)	87	90
1	C	139/146 (95%)	132 (95%)	7 (5%)	28	23
1	D	143/146 (98%)	139 (97%)	4 (3%)	49	49
1	E	141/146 (97%)	138 (98%)	3 (2%)	59	62
1	F	144/146 (99%)	139 (96%)	5 (4%)	41	39
All	All	848/876 (97%)	822 (97%)	26 (3%)	45	44

5 of 26 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	83	ASN
1	D	23	ARG
1	F	83	ASN
1	C	85	ASN
1	C	163	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 35 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	79	HIS
1	D	27	GLN
1	F	78	ASN
1	C	83	ASN
1	C	85	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 6 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PO4	A	201	-	4,4,4	1.37	0	6,6,6	0.40	0
2	PO4	B	201	-	4,4,4	1.30	0	6,6,6	0.41	0
2	PO4	C	201	-	4,4,4	1.31	0	6,6,6	0.40	0
2	PO4	D	201	-	4,4,4	1.29	0	6,6,6	0.40	0
2	PO4	E	201	-	4,4,4	1.33	0	6,6,6	0.41	0
2	PO4	F	201	-	4,4,4	1.24	0	6,6,6	0.42	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PO4	A	201	-	-	0/0/0/0	0/0/0/0
2	PO4	B	201	-	-	0/0/0/0	0/0/0/0
2	PO4	C	201	-	-	0/0/0/0	0/0/0/0
2	PO4	D	201	-	-	0/0/0/0	0/0/0/0
2	PO4	E	201	-	-	0/0/0/0	0/0/0/0
2	PO4	F	201	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	159/172 (92%)	0.72	18 (11%) 6 6	11, 28, 46, 52	0
1	B	160/172 (93%)	0.05	6 (3%) 41 41	8, 18, 40, 60	0
1	C	158/172 (91%)	0.16	7 (4%) 35 35	8, 20, 42, 60	0
1	D	163/172 (94%)	0.04	3 (1%) 69 68	6, 15, 35, 47	0
1	E	161/172 (93%)	0.10	3 (1%) 67 66	7, 19, 35, 47	0
1	F	164/172 (95%)	0.08	7 (4%) 36 36	8, 16, 38, 55	0
All	All	965/1032 (93%)	0.19	44 (4%) 33 33	6, 19, 42, 60	0

The worst 5 of 44 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	18	ILE	5.6
1	D	165	LEU	5.2
1	B	19	LYS	4.7
1	A	38	TYR	4.4
1	A	165	LEU	4.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	PO4	F	201	5/5	0.96	0.11	-0.04	19,20,35,39	0
2	PO4	A	201	5/5	0.98	0.08	-1.23	21,26,29,36	0
2	PO4	B	201	5/5	0.97	0.07	-1.43	18,19,32,33	0
2	PO4	C	201	5/5	0.98	0.07	-2.14	18,24,25,31	0
2	PO4	E	201	5/5	0.99	0.07	-2.16	12,12,16,22	0
2	PO4	D	201	5/5	0.98	0.07	-2.23	13,15,18,20	0
3	CL	D	202	1/1	0.95	0.13	-	31,31,31,31	0
3	CL	E	202	1/1	0.95	0.11	-	35,35,35,35	0
3	CL	A	202	1/1	0.97	0.07	-	37,37,37,37	0
3	CL	C	202	1/1	0.92	0.08	-	34,34,34,34	0
3	CL	F	202	1/1	0.97	0.14	-	24,24,24,24	0
3	CL	B	202	1/1	0.96	0.12	-	27,27,27,27	0

6.5 Other polymers [i](#)

There are no such residues in this entry.