



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 10, 2018 – 10:50 PM EST

PDB ID : 6F6N  
Title : CRYSTAL STRUCTURE OF EBOLAVIRUS GLYCOPROTEIN IN COMPLEX WITH SERTRALINE  
Authors : Ren, J.; Zhao, Y.; Fry, E.E.; Stuart, D.I.  
Deposited on : 2017-12-05  
Resolution : 2.15 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20030736
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20030736


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	1170 (2.16-2.16)
Clashscore	112137	1278 (2.16-2.16)
Ramachandran outliers	110173	1256 (2.16-2.16)
Sidechain outliers	110143	1255 (2.16-2.16)
RSRZ outliers	101464	1175 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	330	
2	B	168	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	NAG	A	504	-	-	-	X
4	GOL	A	505	-	-	-	X
4	GOL	A	506	-	-	-	X
4	GOL	A	507	-	-	-	X
4	GOL	B	707	-	-	-	X
4	GOL	B	708	-	-	-	X
5	SRE	A	508	-	-	-	X
8	DMS	B	706	-	-	-	X

## 2 Entry composition

There are 9 unique types of molecules in this entry. The entry contains 3308 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Envelope glycoprotein,Envelope glycoprotein,GP1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	253	Total	C	N	O	S	0	1	0
			1972	1259	331	377	5			

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	28	GLU	-	expression tag	UNP Q05320
A	29	THR	-	expression tag	UNP Q05320
A	30	GLY	-	expression tag	UNP Q05320
A	31	ARG	-	expression tag	UNP Q05320
A	42	ALA	THR	engineered mutation	UNP Q05320

- Molecule 2 is a protein called Envelope glycoprotein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	131	Total	C	N	O	S	0	0	0
			1035	660	177	192	6			

There are 38 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	613	ALA	HIS	engineered mutation	UNP A0A0U3BWW0
B	633	GLY	-	expression tag	UNP A0A0U3BWW0
B	634	SER	-	expression tag	UNP A0A0U3BWW0
B	635	GLY	-	expression tag	UNP A0A0U3BWW0
B	636	TYR	-	expression tag	UNP A0A0U3BWW0
B	637	ILE	-	expression tag	UNP A0A0U3BWW0
B	638	PRO	-	expression tag	UNP A0A0U3BWW0
B	639	GLU	-	expression tag	UNP A0A0U3BWW0
B	640	ALA	-	expression tag	UNP A0A0U3BWW0
B	641	PRO	-	expression tag	UNP A0A0U3BWW0
B	642	ARG	-	expression tag	UNP A0A0U3BWW0

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	643	ASP	-	expression tag	UNP A0A0U3BWW0
B	644	GLY	-	expression tag	UNP A0A0U3BWW0
B	645	GLN	-	expression tag	UNP A0A0U3BWW0
B	646	ALA	-	expression tag	UNP A0A0U3BWW0
B	647	TYR	-	expression tag	UNP A0A0U3BWW0
B	648	VAL	-	expression tag	UNP A0A0U3BWW0
B	649	ARG	-	expression tag	UNP A0A0U3BWW0
B	650	LYS	-	expression tag	UNP A0A0U3BWW0
B	651	ASP	-	expression tag	UNP A0A0U3BWW0
B	652	GLY	-	expression tag	UNP A0A0U3BWW0
B	653	GLU	-	expression tag	UNP A0A0U3BWW0
B	654	TRP	-	expression tag	UNP A0A0U3BWW0
B	655	VAL	-	expression tag	UNP A0A0U3BWW0
B	656	LEU	-	expression tag	UNP A0A0U3BWW0
B	657	LEU	-	expression tag	UNP A0A0U3BWW0
B	658	SER	-	expression tag	UNP A0A0U3BWW0
B	659	THR	-	expression tag	UNP A0A0U3BWW0
B	660	PHE	-	expression tag	UNP A0A0U3BWW0
B	661	LEU	-	expression tag	UNP A0A0U3BWW0
B	662	GLY	-	expression tag	UNP A0A0U3BWW0
B	663	THR	-	expression tag	UNP A0A0U3BWW0
B	664	HIS	-	expression tag	UNP A0A0U3BWW0
B	665	HIS	-	expression tag	UNP A0A0U3BWW0
B	666	HIS	-	expression tag	UNP A0A0U3BWW0
B	667	HIS	-	expression tag	UNP A0A0U3BWW0
B	668	HIS	-	expression tag	UNP A0A0U3BWW0
B	669	HIS	-	expression tag	UNP A0A0U3BWW0

- Molecule 3 is N-ACETYL-D-GLUCOSAMINE (three-letter code: NAG) (formula: C<sub>8</sub>H<sub>15</sub>NO<sub>6</sub>).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	B	1	Total	C	N	O	0	0
			14	8	1	5		
3	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is (1S,4S)-4-(3,4-dichlorophenyl)-N-methyl-1,2,3,4-tetrahydronaphthalen-1-amine (three-letter code: SRE) (formula: C<sub>17</sub>H<sub>17</sub>Cl<sub>2</sub>N).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	Cl	N	0	0
			20	17	2	1		


- Molecule 6 is BETA-D-MANNOSE (three-letter code: BMA) (formula:  $C_6H_{12}O_6$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	B	1	Total	C	O	0	0
			11	6	5		


- Molecule 7 is ALPHA-D-MANNOSE (three-letter code: MAN) (formula:  $C_6H_{12}O_6$ ).





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	B	1	Total	C	O	0	0
			11	6	5		
7	B	1	Total	C	O	0	0
			11	6	5		

- Molecule 8 is DIMETHYL SULFOXIDE (three-letter code: DMS) (formula:  $C_2H_6OS$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
8	B	1	Total	C	O	S	0	0
			4	2	1	1		


- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	85	Total 85	O 85	0	0
9	B	45	Total 45	O 45	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Envelope glycoprotein,Envelope glycoprotein,GP1



- Molecule 2: Envelope glycoprotein



## 4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	114.25Å 114.25Å 306.19Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	60.54 – 2.15 60.54 – 2.15	Depositor EDS
% Data completeness (in resolution range)	100.0 (60.54-2.15) 100.0 (60.54-2.15)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.67 (at 2.14Å)	Xtriage
Refinement program	PHENIX (dev_2940: ???)	Depositor
R, $R_{free}$	0.184 , 0.203 0.181 , 0.201	Depositor DCC
$R_{free}$ test set	1998 reflections (4.73%)	DCC
Wilson B-factor (Å <sup>2</sup> )	50.6	Xtriage
Anisotropy	0.325	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 59.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	3308	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	75.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.25% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, BMA, NAG, SRE, DMS, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.28	0/1986	0.47	0/2699
2	B	0.28	0/1060	0.44	0/1445
All	All	0.28	0/3046	0.46	0/4144

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	615	TRP	Peptide

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1972	0	1882	19	0
2	B	1035	0	1001	15	0
3	A	56	0	52	0	0
3	B	28	0	24	0	0
4	A	18	0	24	1	0
4	B	12	0	16	1	0
5	A	20	0	17	4	0
6	B	11	0	8	0	0
7	B	22	0	20	0	0
8	B	4	0	6	1	0
9	A	85	0	0	0	0
9	B	45	0	0	0	0
All	All	3308	0	3050	30	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 30 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:45:VAL:HG13	2:B:502:GLU:HG3	1.67	0.77
1:A:248:PHE:HA	1:A:283:THR:HG21	1.70	0.73
2:B:523:GLU:HG2	2:B:524:GLY:H	1.61	0.66
2:B:615:TRP:CD1	2:B:618:ASN:HB3	2.38	0.58
1:A:106:GLU:HG3	1:A:290:PHE:CD1	2.38	0.58

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	239/330 (72%)	232 (97%)	7 (3%)	0	<b>100</b> <b>100</b>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	B	129/168 (77%)	118 (92%)	10 (8%)	1 (1%)	22	14
All	All	368/498 (74%)	350 (95%)	17 (5%)	1 (0%)	44	41

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	525	ALA

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	210/273 (77%)	207 (99%)	3 (1%)	71	77
2	B	109/139 (78%)	109 (100%)	0	100	100
All	All	319/412 (77%)	316 (99%)	3 (1%)	82	87

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	44	GLN
1	A	66	VAL
1	A	268	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	255	GLN
2	B	551	GLN
2	B	508	GLN
1	A	139	HIS
2	B	550	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

16 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
3	NAG	A	501	1	14,14,15	0.21	0	15,19,21	0.57	0
3	NAG	A	502	1	14,14,15	1.48	1 (7%)	15,19,21	1.64	3 (20%)
3	NAG	A	503	1	14,14,15	0.38	0	15,19,21	0.47	0
3	NAG	A	504	1	14,14,15	0.78	1 (7%)	15,19,21	1.19	1 (6%)
4	GOL	A	505	-	5,5,5	0.87	0	5,5,5	1.00	0
4	GOL	A	506	-	5,5,5	0.90	0	5,5,5	0.93	0
4	GOL	A	507	-	5,5,5	0.76	0	5,5,5	0.89	0
5	SRE	A	508	-	22,22,22	2.27	5 (22%)	31,31,31	2.43	6 (19%)
3	NAG	B	701	3,2	14,14,15	0.23	0	15,19,21	0.62	0
3	NAG	B	702	3,6	14,14,15	0.19	0	15,19,21	0.57	0
6	BMA	B	703	3,7	11,11,12	0.82	0	13,15,17	1.23	1 (7%)
7	MAN	B	704	6	11,11,12	0.79	0	13,15,17	1.06	2 (15%)
7	MAN	B	705	6	11,11,12	1.17	1 (9%)	13,15,17	1.00	1 (7%)
8	DMS	B	706	-	3,3,3	0.63	0	3,3,3	0.45	0
4	GOL	B	707	-	5,5,5	0.82	0	5,5,5	1.03	0
4	GOL	B	708	-	5,5,5	0.67	0	5,5,5	1.12	0



In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	501	1	-	0/6/23/26	0/1/1/1
3	NAG	A	502	1	-	0/6/23/26	0/1/1/1
3	NAG	A	503	1	-	0/6/23/26	0/1/1/1
3	NAG	A	504	1	-	0/6/23/26	0/1/1/1
4	GOL	A	505	-	-	0/4/4/4	0/0/0/0
4	GOL	A	506	-	-	0/4/4/4	0/0/0/0
4	GOL	A	507	-	-	0/4/4/4	0/0/0/0
5	SRE	A	508	-	-	0/5/19/19	0/3/3/3
3	NAG	B	701	3,2	-	0/6/23/26	0/1/1/1
3	NAG	B	702	3,6	-	0/6/23/26	0/1/1/1
6	BMA	B	703	3,7	-	0/2/19/22	0/1/1/1
7	MAN	B	704	6	-	0/2/19/22	1/1/1/1
7	MAN	B	705	6	-	0/2/19/22	0/1/1/1
8	DMS	B	706	-	-	0/0/0/0	0/0/0/0
4	GOL	B	707	-	-	0/4/4/4	0/0/0/0
4	GOL	B	708	-	-	0/4/4/4	0/0/0/0

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	502	NAG	O5-C1	-4.82	1.35	1.43
5	A	508	SRE	C3-C4	-2.52	1.50	1.54
5	A	508	SRE	C16-CL19	2.11	1.78	1.73
7	B	705	MAN	C1-C2	2.18	1.57	1.52
5	A	508	SRE	C15-CL20	2.31	1.79	1.73

The worst 5 of 14 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	502	NAG	C1-O5-C5	-3.97	106.69	112.17
5	A	508	SRE	C3-C4-C13	-3.74	104.66	111.31
5	A	508	SRE	C13-C4-C5	-3.61	108.49	113.29
7	B	704	MAN	O2-C2-C3	-2.26	105.73	110.17
6	B	703	BMA	O2-C2-C3	-2.07	106.11	110.17

There are no chirality outliers.

There are no torsion outliers.

All (1) ring outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	B	704	MAN	C1-C2-C3-C4-C5-O5

4 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	506	GOL	1	0
5	A	508	SRE	4	0
8	B	706	DMS	1	0
4	B	708	GOL	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	246/330 (74%)	0.54	21 (8%) 11 15	40, 62, 130, 159	0
2	B	131/168 (77%)	1.30	21 (16%) 2 3	41, 56, 191, 204	0
All	All	377/498 (75%)	0.80	42 (11%) 6 8	40, 59, 159, 204	0

The worst 5 of 42 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	626	ILE	13.2
2	B	620	THR	12.3
2	B	630	PHE	10.4
2	B	631	VAL	9.1
2	B	627	ILE	9.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
8	DMS	B	706	4/4	0.87	0.35	20.87	138,140,140,141	0
4	GOL	A	506	6/6	0.94	0.30	9.16	49,63,67,67	0
4	GOL	B	707	6/6	0.95	0.23	8.13	63,71,76,80	0
4	GOL	B	708	6/6	0.74	0.23	4.69	91,101,102,102	0
5	SRE	A	508	20/20	0.82	0.22	3.68	94,103,117,120	0
4	GOL	A	505	6/6	0.79	0.23	3.59	95,105,107,109	0
3	NAG	A	504	14/15	0.66	0.48	2.96	119,134,143,143	0
4	GOL	A	507	6/6	0.89	0.16	2.39	94,101,108,110	0
3	NAG	A	501	14/15	0.96	0.13	-0.53	56,75,80,80	0
3	NAG	B	701	14/15	0.96	0.12	-1.63	45,52,59,65	0
7	MAN	B	704	11/12	0.78	0.38	-	140,145,147,147	0
7	MAN	B	705	11/12	0.69	0.32	-	117,135,140,142	0
3	NAG	B	702	14/15	0.92	0.14	-	72,81,91,101	0
6	BMA	B	703	11/12	0.83	0.21	-	106,111,125,135	0
3	NAG	A	502	14/15	0.71	0.46	-	129,140,144,146	0
3	NAG	A	503	14/15	0.88	0.18	-	96,108,111,113	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.