



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 9, 2017 – 03:06 AM EDT

PDB ID : 2F7O
Title : Golgi alpha-mannosidase II complex with mannostatin A
Authors : Kuntz, D.A.; Rose, D.R.
Deposited on : unknown
Resolution : 1.43 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20030345
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20030345


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.43 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	1367 (1.46-1.42)
Clashscore	112137	1425 (1.46-1.42)
Ramachandran outliers	110173	1405 (1.46-1.42)
Sidechain outliers	110143	1405 (1.46-1.42)
RSRZ outliers	101464	1372 (1.46-1.42)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1045	<div> <div style="width: 3%; height: 10px; background-color: red;"></div> <div style="width: 86%; height: 10px; background-color: green;"></div> <div style="width: 11%; height: 10px; background-color: yellow;"></div> <div style="width: 2%; height: 10px; background-color: grey;"></div> </div> <div> <div style="width: 3%;"></div> <div style="width: 86%;"></div> <div style="width: 11%;"></div> <div style="width: 2%;"></div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	PO4	A	5005	-	-	X	-
5	MSN	A	5002[A]	-	-	-	X

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	MSN	A	5002[B]	-	-	-	X
6	MPD	A	5003	X	-	-	-

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 9306 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called alpha-mannosidase II.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	1014	Total	C	N	O	S	0	11	0
			8218	5226	1438	1512	42			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	ARG	-	CLONING ATIFACT	UNP Q24451
A	2	SER	-	CLONING ATIFACT	UNP Q24451
A	3	SER	-	CLONING ATIFACT	UNP Q24451
A	4	HIS	-	EXPRESSION TAG	UNP Q24451
A	5	HIS	-	EXPRESSION TAG	UNP Q24451
A	6	HIS	-	EXPRESSION TAG	UNP Q24451
A	7	HIS	-	EXPRESSION TAG	UNP Q24451
A	8	HIS	-	EXPRESSION TAG	UNP Q24451
A	9	HIS	-	EXPRESSION TAG	UNP Q24451
A	10	GLY	-	CLONING ATIFACT	UNP Q24451
A	11	GLU	-	CLONING ATIFACT	UNP Q24451
A	12	PHE	-	CLONING ATIFACT	UNP Q24451

- Molecule 2 is N-ACETYL-D-GLUCOSAMINE (three-letter code: NAG) (formula: C₈H₁₅NO₆).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Zn	0	0
			1	1		

- Molecule 4 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	P	0	0
			5	4	1		

- Molecule 5 is (1R,2R,3R,4S,5R)-4-AMINO-5-(METHYLTHIO)CYCLOPENTANE-1,2,3-TRIOL (three-letter code: MSN) (formula: $C_6H_{13}NO_3S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	S	0	1
			12	7	1	3	1		

- Molecule 6 is (4S)-2-METHYL-2,4-PENTANEDIOL (three-letter code: MPD) (formula: $C_6H_{14}O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			8	6	2		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	1048	Total	O	0	0
			1048	1048		

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	69.00Å 109.30Å 138.35Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.75 – 1.43 29.75 – 1.43	Depositor EDS
% Data completeness (in resolution range)	95.2 (29.75-1.43) 95.4 (29.75-1.43)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.75 (at 1.43Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.206 , 0.235 0.207 , 0.237	Depositor DCC
R_{free} test set	4103 reflections (2.23%)	DCC
Wilson B-factor (Å ²)	11.8	Xtriage
Anisotropy	0.576	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 49.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.43$, $\langle L^2 \rangle = 0.26$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	9306	wwPDB-VP
Average B, all atoms (Å ²)	16.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.59% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, MPD, ZN, NAG, MSN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.92	3/8494 (0.0%)	0.98	10/11534 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	578	VAL	CB-CG2	-5.36	1.41	1.52
1	A	829	TYR	CE2-CZ	5.17	1.45	1.38
1	A	892	TYR	CE1-CZ	5.11	1.45	1.38

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	963	ARG	NE-CZ-NH2	-7.00	116.80	120.30
1	A	540	ARG	NE-CZ-NH1	6.84	123.72	120.30
1	A	818	ARG	NE-CZ-NH2	-6.58	117.01	120.30
1	A	868	ARG	NE-CZ-NH1	6.09	123.34	120.30
1	A	422	ARG	NE-CZ-NH2	5.91	123.25	120.30
1	A	167	MET	N-CA-C	-5.78	95.39	111.00
1	A	963	ARG	NE-CZ-NH1	5.76	123.18	120.30
1	A	565	ARG	NE-CZ-NH2	-5.73	117.44	120.30
1	A	781	ASP	CB-CG-OD1	5.64	123.38	118.30
1	A	270	ASP	CB-CG-OD1	5.44	123.19	118.30

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	102	TYR	Sidechain
1	A	53[B]	SER	Mainchain
1	A	979[B]	LEU	Mainchain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	8218	0	8016	78	0
2	A	14	0	13	0	0
3	A	1	0	0	0	0
4	A	5	0	0	2	0
5	A	12	0	6	2	0
6	A	8	0	14	0	0
7	A	1048	0	0	25	0
All	All	9306	0	8049	79	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (79) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:5002[A]:MSN:H73	7:A:5085:HOH:O	1.60	0.99
1:A:723:ARG:HH11	1:A:723:ARG:HG2	1.30	0.97
1:A:300:LYS:HE2	7:A:5067:HOH:O	1.78	0.84
1:A:742:GLN:HG3	7:A:5439:HOH:O	1.86	0.74
1:A:723:ARG:NH1	1:A:723:ARG:HG2	1.97	0.73
1:A:69:LYS:HG3	7:A:5665:HOH:O	1.88	0.71
1:A:989:THR:HG22	1:A:991:GLU:HG3	1.71	0.71
1:A:739:GLU:HG3	7:A:5416:HOH:O	1.91	0.69
1:A:290:MET:CG	1:A:303:PRO:HG2	2.25	0.67
1:A:770:ARG:NH1	4:A:5005:PO4:O1	2.31	0.64
1:A:904:ARG:HG2	1:A:985:MET:SD	2.39	0.62

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:32:GLN:HG2	7:A:5989:HOH:O	1.99	0.62
1:A:290:MET:HG3	1:A:303:PRO:HG2	1.81	0.61
1:A:954:ARG:HB3	7:A:5714:HOH:O	2.03	0.57
1:A:1008:ARG:NH1	1:A:1022:HIS:CE1	2.73	0.56
1:A:154:LYS:HE2	7:A:5892:HOH:O	2.06	0.56
1:A:290:MET:HG2	1:A:303:PRO:HG2	1.88	0.56
1:A:32:GLN:HG3	7:A:5991:HOH:O	2.05	0.56
1:A:703:SER:HB2	1:A:704:PRO:HD2	1.87	0.55
1:A:456:GLU:HB2	7:A:5619:HOH:O	2.07	0.54
1:A:256:ASP:HB2	7:A:5988:HOH:O	2.07	0.54
1:A:31:CYS:N	7:A:6015:HOH:O	2.39	0.54
1:A:62:TRP:CD2	1:A:65:GLY:HA3	2.43	0.53
1:A:53[B]:SER:OG	1:A:55:LYS:HG2	2.07	0.53
1:A:770:ARG:NH2	1:A:775:GLU:OE1	2.42	0.53
1:A:770:ARG:HH12	4:A:5005:PO4:P	2.32	0.52
1:A:141:LEU:O	1:A:146:LYS:HE3	2.10	0.52
1:A:294:GLY:C	7:A:5178:HOH:O	2.47	0.51
1:A:666:GLU:HG2	7:A:5577:HOH:O	2.09	0.51
1:A:1008:ARG:CZ	1:A:1022:HIS:CE1	2.94	0.51
1:A:96:ILE:O	1:A:96:ILE:HG22	2.11	0.50
1:A:981:ARG:NH1	7:A:5714:HOH:O	2.43	0.50
1:A:1008:ARG:CZ	1:A:1022:HIS:NE2	2.76	0.49
1:A:251:TRP:C	1:A:251:TRP:CD1	2.86	0.49
1:A:990:PRO:O	1:A:992:GLU:HG2	2.12	0.49
1:A:537:GLU:HG2	1:A:539:SER:HB3	1.95	0.49
1:A:116:ARG:HD2	1:A:117[A]:HIS:CD2	2.48	0.48
1:A:651:LEU:CD1	1:A:653:ARG:HG2	2.43	0.48
1:A:359:ARG:HD3	7:A:5237:HOH:O	2.13	0.48
1:A:693:LEU:HD13	7:A:5412:HOH:O	2.14	0.47
1:A:250:ILE:HB	1:A:910:PRO:HG2	1.96	0.47
1:A:580:VAL:HG22	1:A:634:LEU:HD22	1.95	0.47
1:A:174:HIS:CE1	1:A:176:ARG:HD3	2.49	0.47
1:A:636:ILE:HG12	1:A:637:SER:N	2.30	0.47
1:A:144:ASN:HB3	7:A:5195:HOH:O	2.15	0.47
1:A:515:TYR:CZ	1:A:517:PRO:HG3	2.50	0.47
1:A:583:LEU:HB2	7:A:5477:HOH:O	2.14	0.46
1:A:264[A]:MET:HE3	7:A:5284:HOH:O	2.15	0.46
1:A:975:VAL:HG21	1:A:1003:LEU:CD1	2.46	0.46
1:A:54:PHE:CG	1:A:822:LEU:HD21	2.51	0.46
1:A:388:GLN:HG3	1:A:391:ARG:NH2	2.31	0.45
1:A:898:LYS:HA	1:A:898:LYS:HD2	1.55	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:58:ASP:OD1	1:A:63:LYS:HE3	2.16	0.45
1:A:96:ILE:HG23	1:A:479:LYS:HD2	1.98	0.45
1:A:411[B]:SER:HB2	1:A:412:ASP:H	1.63	0.44
1:A:448:TRP:CG	1:A:454:ILE:HG13	2.52	0.44
1:A:584:ALA:HB3	1:A:586:ASN:ND2	2.32	0.44
1:A:155:ASN:HB3	7:A:5894:HOH:O	2.17	0.44
1:A:714:LYS:HB3	1:A:714:LYS:HE2	1.78	0.44
1:A:468:PHE:CZ	1:A:474:ILE:HA	2.53	0.43
1:A:723:ARG:NH1	1:A:723:ARG:CG	2.72	0.43
1:A:567:GLN:CD	7:A:5380:HOH:O	2.56	0.43
1:A:63:LYS:NZ	7:A:5293:HOH:O	2.50	0.43
1:A:666:GLU:HG3	1:A:667:ASP:N	2.32	0.43
1:A:661:LEU:HB2	1:A:664:TYR:HB3	2.01	0.43
1:A:480:THR:HG23	7:A:5328:HOH:O	2.18	0.43
1:A:53[A]:SER:O	1:A:54:PHE:HB2	2.19	0.43
1:A:934:PHE:CE2	1:A:936:GLU:HB2	2.54	0.42
1:A:770:ARG:HD2	1:A:770:ARG:HH11	1.70	0.42
1:A:280:LYS:HE2	1:A:301:VAL:CG2	2.50	0.42
1:A:47:GLU:CD	1:A:51:ARG:HE	2.24	0.41
1:A:761:PRO:O	1:A:762:SER:HB2	2.20	0.41
1:A:531:TRP:CD2	1:A:532:PRO:HA	2.54	0.41
1:A:990:PRO:O	1:A:991:GLU:C	2.59	0.41
1:A:936:GLU:OE1	1:A:936:GLU:N	2.54	0.41
1:A:228:ARG:NH2	5:A:5002[A]:MSN:H72	2.36	0.41
1:A:540:ARG:HB2	7:A:5588:HOH:O	2.21	0.40
1:A:394:GLN:O	1:A:394:GLN:HG2	2.22	0.40
1:A:206:PHE:HB2	1:A:418:TYR:CE1	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	1023/1045 (98%)	990 (97%)	27 (3%)	6 (1%)	28 6

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	991	GLU
1	A	95	TRP
1	A	665	PRO
1	A	683	GLY
1	A	204	ASP
1	A	549	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	912/929 (98%)	900 (99%)	12 (1%)	73 40

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	116	ARG
1	A	206	PHE
1	A	275	CYS
1	A	302	PRO
1	A	375	GLN
1	A	447[A]	SER
1	A	447[B]	SER
1	A	549	ILE
1	A	723	ARG
1	A	828	TYR
1	A	898	LYS
1	A	936	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	742	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 1 is monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	MSN	A	5002[A]	-	9,11,11	0.45	0	8,16,16	4.65	3 (37%)
5	MSN	A	5002[B]	-	9,11,11	0.45	0	8,16,16	4.65	3 (37%)
6	MPD	A	5003	-	7,7,7	0.77	0	9,10,10	0.55	0
2	NAG	A	5004	1	14,14,15	0.65	0	15,19,21	1.16	1 (6%)
4	PO4	A	5005	-	4,4,4	2.24	1 (25%)	6,6,6	0.89	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	MSN	A	5002[A]	-	-	0/2/22/22	0/1/1/1
5	MSN	A	5002[B]	-	-	0/2/22/22	0/1/1/1
6	MPD	A	5003	-	1/1/2/2	0/5/5/5	0/0/0/0
2	NAG	A	5004	1	-	0/6/23/26	0/1/1/1
4	PO4	A	5005	-	-	0/0/0/0	0/0/0/0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	5005	PO4	P-O1	4.28	1.59	1.50

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	5002[B]	MSN	C3-C2-C1	-10.64	95.74	106.30
5	A	5002[A]	MSN	C3-C2-C1	-10.64	95.74	106.30
2	A	5004	NAG	C4-C3-C2	-2.63	107.17	111.02
5	A	5002[B]	MSN	O2-C2-C3	3.65	123.52	111.83
5	A	5002[A]	MSN	O2-C2-C3	3.65	123.52	111.83
5	A	5002[B]	MSN	C2-C3-C4	6.41	112.99	102.64
5	A	5002[A]	MSN	C2-C3-C4	6.41	112.99	102.64

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
6	A	5003	MPD	C4

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	5002[A]	MSN	2	0
4	A	5005	PO4	2	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	1014/1045 (97%)	0.17	32 (3%)	48 49	6, 14, 28, 67	0

All (32) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	603	LEU	7.3
1	A	992	GLU	5.7
1	A	701	GLN	5.7
1	A	993	HIS	5.6
1	A	702	ASP	5.0
1	A	720	HIS	4.9
1	A	602	THR	4.7
1	A	78	HIS	4.4
1	A	600	HIS	4.4
1	A	991	GLU	4.3
1	A	536	VAL	4.1
1	A	721	GLY	4.1
1	A	638	ASP	3.6
1	A	538	ASP	3.5
1	A	682	ASN	3.2
1	A	990	PRO	3.2
1	A	534	SER	2.9
1	A	1044	SER	2.9
1	A	719	SER	2.9
1	A	1022	HIS	2.8
1	A	1024	ASP	2.7
1	A	742	GLN	2.6
1	A	57[A]	ILE	2.6
1	A	535	GLY	2.3
1	A	547	GLU	2.2
1	A	655	ASN	2.2
1	A	392	ALA	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	700	THR	2.1
1	A	606	THR	2.1
1	A	537	GLU	2.1
1	A	1026	MET	2.1
1	A	549	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
5	MSN	A	5002[B]	11/11	0.95	0.11	9.50	6,8,10,12	1
5	MSN	A	5002[A]	11/11	0.95	0.11	6.71	5,8,10,12	1
4	PO4	A	5005	5/5	0.95	0.13	1.75	28,29,31,32	0
6	MPD	A	5003	8/8	0.88	0.14	1.57	15,21,23,23	0
3	ZN	A	5001	1/1	1.00	0.05	-3.15	8,8,8,8	0
2	NAG	A	5004	14/15	0.73	0.36	-	35,41,43,43	0

6.5 Other polymers [i](#)

There are no such residues in this entry.