



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 15, 2017 – 10:21 AM EDT

PDB ID : 2F8Q
Title : An alkali thermostable F/10 xylanase from alkalophilic Bacillus sp. NG-27
Authors : Ramakumar, S.; Manikandan, K.; Bhardwaj, A.; Ghosh, A.; Reddy, V.S.
Deposited on : unknown
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.9-1692
EDS : rb-20030345
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20030345


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	4002 (2.20-2.20)
Clashscore	112137	4730 (2.20-2.20)
Ramachandran outliers	110173	4656 (2.20-2.20)
Sidechain outliers	110143	4657 (2.20-2.20)
RSRZ outliers	101464	4033 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	353	 83% 16% •
1	B	353	 4% 81% 19%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 6458 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called alkaline thermostable endoxylanase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	353	Total	C	N	O	S	0	0	0
			2850	1816	487	542	5			
1	B	353	Total	C	N	O	S	0	0	0
			2854	1818	487	544	5			

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Mg	0	0
			1	1		
2	A	1	Total	Mg	0	0
			1	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	391	Total	O	0	0
			391	391		
3	B	361	Total	O	0	0
			361	361		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: alkaline thermostable endoxylanase



- Molecule 1: alkaline thermostable endoxylanase



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	174.50 Å 54.74 Å 131.50 Å 90.00° 131.21° 90.00°	Depositor
Resolution (Å)	20.00 – 2.20 20.00 – 2.20	Depositor EDS
% Data completeness (in resolution range)	96.6 (20.00-2.20) 96.7 (20.00-2.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.97 (at 2.21 Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.197 , 0.235 0.196 , 0.200	Depositor DCC
R_{free} test set	2297 reflections (4.98%)	DCC
Wilson B-factor (Å ²)	25.4	Xtriage
Anisotropy	0.166	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 71.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.014 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6458	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.92% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.32	0/2930	0.58	0/3999
1	B	0.32	0/2934	0.58	0/4004
All	All	0.32	0/5864	0.58	0/8003

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2850	0	2657	46	0
1	B	2854	0	2661	54	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	391	0	0	4	0
3	B	361	0	0	3	0
All	All	6458	0	5318	100	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (100) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:316:ASN:HD22	1:B:316:ASN:H	1.20	0.88
1:B:31:ASN:HA	3:B:1132:HOH:O	1.76	0.84
1:A:316:ASN:HD22	1:A:316:ASN:H	1.26	0.81
1:A:316:ASN:HD22	1:A:316:ASN:N	1.83	0.76
1:B:316:ASN:HD22	1:B:316:ASN:N	1.84	0.74
1:B:123:LEU:O	1:B:127:GLU:HG3	1.89	0.73
1:B:231:ILE:HD11	1:B:294:LEU:HD21	1.70	0.72
1:A:277:ASP:HB2	3:A:924:HOH:O	1.92	0.69
1:B:2:GLN:HE21	1:B:2:GLN:HA	1.63	0.63
1:A:123:LEU:O	1:A:127:GLU:HG3	1.98	0.63
1:A:259:GLU:HG2	1:A:311:TRP:CE3	2.35	0.62
1:B:59:GLU:HG3	1:B:62:VAL:HG21	1.82	0.62
1:B:268:PRO:HB3	1:B:327:TYR:CD2	2.35	0.61
1:B:59:GLU:HG3	1:B:62:VAL:CG2	2.32	0.59
1:A:316:ASN:H	1:A:316:ASN:ND2	1.99	0.59
1:A:104:ASN:CB	3:A:1174:HOH:O	2.52	0.58
1:B:54:ILE:HA	1:B:94:PRO:HD3	1.86	0.57
1:B:109:GLU:HG2	1:B:114:LYS:HG3	1.88	0.55
1:A:228:GLN:HE21	1:A:230:HIS:CD2	2.25	0.55
1:A:130:ILE:O	1:A:134:VAL:HG23	2.06	0.55
1:B:2:GLN:HE22	1:B:350:TRP:HE1	1.55	0.54
1:A:111:ASN:HD22	1:A:114:LYS:H	1.54	0.54
1:A:316:ASN:N	1:A:316:ASN:ND2	2.56	0.54
1:B:131:LYS:HD3	1:B:131:LYS:C	2.28	0.54
1:B:316:ASN:ND2	1:B:316:ASN:H	1.96	0.54
1:B:292:ASN:O	1:B:296:GLU:HG3	2.08	0.53
1:B:148:ASN:HD22	1:B:195:ASN:HB3	1.73	0.53
1:A:321:ASP:O	1:A:325:ARG:HG3	2.07	0.53
1:B:200:GLU:HG2	1:B:241:ILE:HG12	1.90	0.53
1:A:315:ASP:HB2	1:A:341:PRO:HD3	1.91	0.53
1:B:94:PRO:HB2	1:B:96:TRP:CD1	2.43	0.52
1:A:2:GLN:N	3:A:934:HOH:O	2.42	0.52
1:A:148:ASN:HD22	1:A:195:ASN:HB3	1.75	0.52
1:B:228:GLN:HE21	1:B:230:HIS:CD2	2.29	0.51
1:A:292:ASN:O	1:A:296:GLU:HG3	2.12	0.50
1:B:84:PHE:CZ	1:B:133:VAL:HG11	2.46	0.50
1:A:94:PRO:HB2	1:A:96:TRP:CD1	2.46	0.50
1:A:2:GLN:HA	1:A:2:GLN:HE21	1.75	0.50
1:B:100:ASP:OD1	1:B:102:GLU:N	2.42	0.50
1:B:111:ASN:HD22	1:B:114:LYS:HB2	1.76	0.50
1:A:131:LYS:O	1:A:135:GLU:HG2	2.11	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:54:ILE:CA	1:B:94:PRO:HD3	2.42	0.50
1:B:228:GLN:HE21	1:B:230:HIS:HD2	1.60	0.49
1:B:111:ASN:HD21	1:B:113:ALA:HB3	1.77	0.49
1:A:340:ASP:HB2	1:A:341:PRO:CD	2.43	0.49
1:A:237:THR:OG1	1:A:240:GLU:HG3	2.13	0.49
1:B:283:ARG:NE	1:B:283:ARG:HA	2.27	0.48
1:A:94:PRO:HG2	1:A:97:PHE:CD2	2.48	0.48
1:A:111:ASN:ND2	1:A:114:LYS:H	2.11	0.48
1:B:267:TRP:HA	1:B:268:PRO:C	2.34	0.48
1:B:115:ARG:HG2	1:B:115:ARG:HH11	1.77	0.48
1:B:237:THR:OG1	1:B:240:GLU:HG3	2.14	0.48
1:B:111:ASN:ND2	1:B:114:LYS:H	2.12	0.47
1:B:340:ASP:HB2	1:B:341:PRO:CD	2.44	0.47
1:A:146:VAL:HG21	1:A:192:LEU:HB3	1.97	0.47
1:A:56:LEU:O	1:A:63:PHE:HA	2.15	0.47
1:B:193:PHE:CE2	1:B:256:GLN:HG3	2.50	0.47
1:B:2:GLN:HE21	1:B:3:PRO:HD2	1.78	0.47
1:A:195:ASN:HD22	1:A:226:GLY:HA3	1.79	0.46
1:A:340:ASP:HB2	1:A:341:PRO:HD2	1.97	0.46
1:A:84:PHE:CZ	1:A:133:VAL:HG11	2.50	0.46
1:B:2:GLN:N	3:B:1027:HOH:O	2.48	0.46
1:A:320:LEU:HD12	1:A:336:PRO:HB3	1.97	0.46
1:A:24:ALA:HA	1:A:46:VAL:HG13	1.97	0.46
1:A:228:GLN:HE21	1:A:230:HIS:HD2	1.62	0.45
1:A:111:ASN:HD21	1:A:113:ALA:HB3	1.82	0.45
1:B:56:LEU:O	1:B:63:PHE:HA	2.17	0.45
1:A:264:LEU:HD11	1:A:335:ALA:HB3	1.99	0.45
1:A:238:ILE:HG23	1:A:297:LEU:HD22	1.99	0.44
1:A:23:ALA:HB2	1:A:42:TYR:CE2	2.53	0.44
1:B:8:VAL:O	1:B:40:HIS:HE1	2.01	0.44
1:B:73:GLU:HB2	3:B:1083:HOH:O	2.16	0.44
1:B:195:ASN:HD22	1:B:226:GLY:HA3	1.82	0.44
1:A:283:ARG:HA	1:A:283:ARG:NE	2.32	0.44
1:B:109:GLU:HG2	1:B:114:LYS:CG	2.48	0.44
1:B:2:GLN:NE2	1:B:2:GLN:HA	2.29	0.44
1:B:111:ASN:O	1:B:114:LYS:N	2.51	0.44
1:B:238:ILE:HG23	1:B:297:LEU:HD22	2.00	0.43
1:B:119:LYS:O	1:B:123:LEU:HG	2.18	0.43
1:A:111:ASN:HD22	1:A:114:LYS:N	2.15	0.43
1:B:225:VAL:HG21	1:B:248:PHE:CD1	2.54	0.43
1:B:315:ASP:OD2	1:B:335:ALA:HB1	2.19	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:283:ARG:HA	1:B:283:ARG:HE	1.84	0.42
1:B:276:TYR:CZ	1:B:341:PRO:HD3	2.54	0.42
1:B:146:VAL:HG21	1:B:192:LEU:HB3	2.00	0.42
1:B:197:TYR:HB3	1:B:228:GLN:NE2	2.34	0.42
1:A:292:ASN:HB2	1:A:351:ARG:HB3	2.02	0.42
1:A:54:ILE:HA	1:A:94:PRO:HD3	2.01	0.42
1:B:24:ALA:HA	1:B:46:VAL:HG13	2.02	0.42
1:A:259:GLU:HG2	1:A:311:TRP:CZ3	2.55	0.41
1:A:8:VAL:O	1:A:40:HIS:HE1	2.03	0.41
1:A:2:GLN:HE22	1:A:350:TRP:HE1	1.68	0.41
1:B:23:ALA:HB2	1:B:42:TYR:CE2	2.55	0.41
1:A:191:LYS:HE3	3:A:945:HOH:O	2.20	0.40
1:A:111:ASN:HB3	1:A:114:LYS:HB3	2.03	0.40
1:A:331:VAL:O	1:A:331:VAL:HG23	2.22	0.40
1:A:205:ARG:HD3	1:A:247:MET:SD	2.61	0.40
1:B:131:LYS:HD3	1:B:132:THR:N	2.36	0.40
1:B:194:ILE:HG23	1:B:194:ILE:O	2.20	0.40
1:B:42:TYR:CD1	1:B:42:TYR:N	2.89	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	351/353 (99%)	344 (98%)	7 (2%)	0	100	100
1	B	351/353 (99%)	332 (95%)	19 (5%)	0	100	100
All	All	702/706 (99%)	676 (96%)	26 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	294/305 (96%)	288 (98%)	6 (2%)	60	74
1	B	295/305 (97%)	292 (99%)	3 (1%)	80	89
All	All	589/610 (97%)	580 (98%)	9 (2%)	70	82

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2	GLN
1	A	34	GLN
1	A	82	LEU
1	A	305	LEU
1	A	316	ASN
1	A	336	PRO
1	B	82	LEU
1	B	305	LEU
1	B	316	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (23) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	2	GLN
1	A	34	GLN
1	A	40	HIS
1	A	91	ASN
1	A	111	ASN
1	A	148	ASN
1	A	195	ASN
1	A	198	ASN
1	A	213	GLN
1	A	228	GLN
1	A	316	ASN
1	B	2	GLN
1	B	40	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	91	ASN
1	B	111	ASN
1	B	116	GLN
1	B	148	ASN
1	B	168	GLN
1	B	195	ASN
1	B	198	ASN
1	B	207	HIS
1	B	228	GLN
1	B	316	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	353/353 (100%)	-0.30	0 100 100	16, 27, 43, 50	15 (4%)
1	B	353/353 (100%)	-0.08	13 (3%) 42 40	14, 27, 46, 57	17 (4%)
All	All	706/706 (100%)	-0.19	13 (1%) 69 66	14, 27, 45, 57	32 (4%)

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	110	THR	4.1
1	B	101	GLU	3.7
1	B	108	GLU	3.5
1	B	111	ASN	3.2
1	B	102	GLU	3.1
1	B	331	VAL	2.8
1	B	113	ALA	2.8
1	B	100	ASP	2.8
1	B	103	GLY	2.4
1	B	115	ARG	2.1
1	B	155	THR	2.1
1	B	271	PRO	2.1
1	B	329	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	MG	B	802	1/1	0.99	0.06	-	11,11,11,11	0
2	MG	A	801	1/1	0.99	0.04	-	7,7,7,7	0

6.5 Other polymers [i](#)

There are no such residues in this entry.