



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 12:55 pm GMT

PDB ID : 5F8M  
Title : Enterovirus 71 Polymerase Elongation Complex (C3S4/5 Form)  
Authors : Shu, B.; Gong, P.  
Deposited on : 2015-12-09  
Resolution : 2.83 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.83 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	3466 (2.88-2.80)
Clashscore	112137	3975 (2.88-2.80)
Ramachandran outliers	110173	3902 (2.88-2.80)
Sidechain outliers	110143	3905 (2.88-2.80)
RSRZ outliers	101464	3501 (2.88-2.80)
RNA backbone	2435	1054 (3.18-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	468	<div> <div>2%</div> <div>79%</div> <div>19%</div> <div>..</div> </div>
2	B	35	<div> <div>3%</div> <div>6%</div> <div>23%</div> <div>6%</div> <div>66%</div> </div>
3	C	19	<div> <div>11%</div> <div>32%</div> <div>21%</div> <div>47%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	MG	A	1001	-	-	-	X
7	POP	A	1005	-	-	-	X

## 2 Entry composition

There are 9 unique types of molecules in this entry. The entry contains 4247 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Genome polypeptide.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	462	Total	C	N	O	S	0	0	0
			3681	2354	611	698	18			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	463	HIS	-	expression tag	UNP E5RPG2
A	464	HIS	-	expression tag	UNP E5RPG2
A	465	HIS	-	expression tag	UNP E5RPG2
A	466	HIS	-	expression tag	UNP E5RPG2
A	467	HIS	-	expression tag	UNP E5RPG2
A	468	HIS	-	expression tag	UNP E5RPG2

- Molecule 2 is a RNA chain called RNA (35-MER).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	12	Total	C	N	O	P	0	0	0
			248	111	38	87	12			

- Molecule 3 is a RNA chain called RNA (5'-R(\*UP\*GP\*UP\*UP\*CP\*GP\*AP\*CP\*GP\*AP\*GP\*AP\*GP\*AP\*GP\*AP\*CP\*CP\*U)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	10	Total	C	N	O	P	0	0	0
			217	97	43	67	10			

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	2	Total	Mg	0	0
			2	2		

- Molecule 5 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Zn	0	0
			1	1		

- Molecule 6 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 7 is PYROPHOSPHATE 2- (three-letter code: POP) (formula: H<sub>2</sub>O<sub>7</sub>P<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	O	P	0	0
			9	7	2		

- Molecule 8 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	70	Total	O	0	0
			70	70		
9	B	6	Total	O	0	0
			6	6		
9	C	2	Total	O	0	0
			2	2		



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	63.64Å 76.71Å 150.10Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	41.91 – 2.83 41.91 – 2.81	Depositor EDS
% Data completeness (in resolution range)	95.8 (41.91-2.83) 95.0 (41.91-2.81)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.76 (at 2.81Å)	Xtriage
Refinement program	PHENIX	Depositor
R, $R_{free}$	0.193 , 0.232 0.193 , 0.232	Depositor DCC
$R_{free}$ test set	876 reflections (5.03%)	DCC
Wilson B-factor (Å <sup>2</sup> )	64.7	Xtriage
Anisotropy	0.334	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 38.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.45$ , $\langle L^2 \rangle = 0.27$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	4247	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	63.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.13% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, MG, ZN, POP, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.46	0/3775	0.58	0/5124
2	B	0.98	0/274	1.26	2/423 (0.5%)
3	C	1.09	2/243 (0.8%)	1.49	4/377 (1.1%)
All	All	0.56	2/4292 (0.0%)	0.74	6/5924 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	702	U	N1-C2	6.15	1.44	1.38
3	C	693	A	N7-C5	5.28	1.42	1.39

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	702	U	N3-C2-O2	-7.04	117.27	122.20
3	C	702	U	N1-C2-O2	6.76	127.53	122.80
2	B	609	U	N3-C2-O2	-6.47	117.67	122.20
2	B	609	U	N1-C2-O2	6.19	127.13	122.80
3	C	701	C	N3-C2-O2	-5.85	117.81	121.90
3	C	702	U	C2-N1-C1'	5.37	124.15	117.70

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3681	0	3583	58	0
2	B	248	0	129	15	0
3	C	217	0	110	7	0
4	A	2	0	0	0	0
5	A	1	0	0	0	0
6	A	5	0	0	0	0
7	A	9	0	0	1	0
8	B	6	0	8	1	0
9	A	70	0	0	3	0
9	B	6	0	0	2	0
9	C	2	0	0	0	0
All	All	4247	0	3830	70	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (70) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:174:ARG:NH2	7:A:1005:POP:O3	2.15	0.78
1:A:17:ILE:HG21	1:A:155:SER:HB3	1.65	0.78
3:C:695:A:H2'	3:C:696:G:H5''	1.77	0.67
1:A:114:THR:HG21	2:B:601:G:OP2	1.96	0.65
2:B:602:G:N3	9:B:801:HOH:O	2.30	0.65
2:B:600:A:H2'	2:B:601:G:C8	2.36	0.61
1:A:2:GLU:HB2	1:A:64:GLY:HA2	1.84	0.58
1:A:79:LEU:HD13	1:A:255:GLU:HG3	1.87	0.57
1:A:114:THR:HG23	2:B:600:A:H5''	1.88	0.56
1:A:114:THR:HG21	2:B:601:G:P	2.46	0.55
2:B:603:U:O2'	8:B:701:GOL:O3	2.24	0.55
1:A:49:ARG:NH1	1:A:168:ILE:HD13	2.21	0.55
2:B:605:U:H2'	2:B:606:C:C6	2.42	0.55
1:A:272:THR:HG21	1:A:288:PRO:HG3	1.89	0.54
1:A:112:LEU:HB3	1:A:128:ARG:HD3	1.90	0.54
1:A:49:ARG:HH12	1:A:168:ILE:HD13	1.72	0.54
1:A:229:LEU:HD11	1:A:369:TRP:CE2	2.43	0.53
2:B:600:A:H2'	2:B:601:G:H8	1.73	0.53
1:A:24:LYS:HE2	2:B:599:C:N4	2.24	0.53
3:C:695:A:C2'	3:C:696:G:H5''	2.39	0.52
1:A:234:TYR:CD2	1:A:329:ASP:HB2	2.44	0.52

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:605:U:H2'	2:B:606:C:H6	1.75	0.52
1:A:442:VAL:O	1:A:446:LEU:HG	2.10	0.50
1:A:450:ASN:HB3	1:A:453:ASN:HD22	1.76	0.50
1:A:434:VAL:O	1:A:438:ARG:HG2	2.12	0.50
1:A:287:MET:O	1:A:289:SER:N	2.45	0.50
1:A:290:GLY:HA2	2:B:600:A:N3	2.27	0.49
1:A:377:ARG:NH2	9:A:1110:HOH:O	2.44	0.49
1:A:29:VAL:HG11	1:A:442:VAL:HG21	1.93	0.49
1:A:174:ARG:NH1	9:A:1103:HOH:O	2.30	0.49
1:A:244:VAL:HG13	1:A:245:TRP:N	2.27	0.49
2:B:607:U:H2'	2:B:608:C:C6	2.48	0.49
1:A:399:HIS:O	1:A:403:ARG:HG3	2.12	0.49
1:A:450:ASN:HB3	1:A:453:ASN:ND2	2.28	0.49
1:A:237:TYR:CG	1:A:329:ASP:HB3	2.48	0.48
1:A:238:ASP:OD1	1:A:239:ALA:N	2.47	0.48
1:A:68:TYR:O	1:A:247:ARG:HG3	2.13	0.48
1:A:222:PRO:HA	1:A:369:TRP:CZ2	2.49	0.47
1:A:155:SER:HB2	1:A:178:ALA:HB3	1.97	0.47
1:A:244:VAL:HG13	1:A:245:TRP:H	1.80	0.47
1:A:24:LYS:HE2	2:B:599:C:H42	1.80	0.46
1:A:390:HIS:CE1	1:A:458:TRP:CZ3	3.03	0.46
1:A:120:TYR:HB3	1:A:125:ILE:HB	1.98	0.46
1:A:114:THR:O	1:A:127:LYS:HE3	2.16	0.46
1:A:147:LYS:HD3	1:A:148:TYR:CE2	2.51	0.46
1:A:181:LEU:O	1:A:185:VAL:HG23	2.16	0.46
1:A:401:SER:OG	3:C:699:A:OP1	2.20	0.45
1:A:449:PRO:HB2	1:A:454:LEU:HD11	1.98	0.45
1:A:5:TRP:CE2	1:A:282:CYS:HB2	2.52	0.44
1:A:448:ILE:HA	1:A:449:PRO:HD3	1.84	0.44
1:A:238:ASP:HB3	3:C:702:U:O2'	2.18	0.44
3:C:696:G:H2'	3:C:697:A:O4'	2.18	0.44
3:C:697:A:H2'	3:C:698:G:C8	2.53	0.43
1:A:438:ARG:CZ	1:A:448:ILE:HD12	2.48	0.43
1:A:438:ARG:O	1:A:444:LYS:HE3	2.18	0.43
1:A:296:ILE:HG13	9:B:804:HOH:O	2.18	0.43
1:A:274:HIS:HB2	1:A:281:TYR:CZ	2.53	0.43
1:A:335:TYR:CG	1:A:336:PRO:HD2	2.54	0.43
1:A:210:VAL:HA	1:A:327:TYR:CE2	2.54	0.42
1:A:154:TYR:HB3	1:A:177:GLU:HB3	2.02	0.42
3:C:700:C:H2'	3:C:701:C:C6	2.54	0.42
1:A:326:ALA:HA	1:A:330:ASP:O	2.20	0.42

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:9:ASN:HB3	1:A:14:ARG:O	2.20	0.41
1:A:230:PHE:HB2	1:A:341:CYS:SG	2.61	0.41
1:A:455:ARG:O	1:A:459:LEU:HG	2.20	0.41
1:A:188:ARG:HD2	2:B:602:G:OP1	2.20	0.41
1:A:127:LYS:NZ	2:B:601:G:OP1	2.53	0.41
1:A:209:ALA:HB2	1:A:325:VAL:HG12	2.02	0.41
1:A:113:HIS:N	1:A:113:HIS:ND1	2.68	0.41
1:A:14:ARG:NH1	9:A:1108:HOH:O	2.54	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	460/468 (98%)	439 (95%)	19 (4%)	2 (0%)	38 69

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	289	SER
1	A	288	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	402/412 (98%)	387 (96%)	15 (4%)	39 71

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	89	ASP
1	A	98	GLU
1	A	113	HIS
1	A	114	THR
1	A	121	SER
1	A	139	SER
1	A	164	SER
1	A	165	ILE
1	A	169	LYS
1	A	212	CYS
1	A	218	TRP
1	A	261	GLU
1	A	332	LEU
1	A	359	ASP
1	A	435	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	B	11/35 (31%)	2 (18%)	0
3	C	9/19 (47%)	3 (33%)	0
All	All	20/54 (37%)	5 (25%)	0

All (5) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
2	B	602	G
2	B	608	C
3	C	694	G
3	C	695	A
3	C	696	G

There are no RNA pucker outliers to report.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 3 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
6	SO4	A	1004	-	4,4,4	0.26	0	6,6,6	0.33	0
7	POP	A	1005	4	8,8,8	1.15	1 (12%)	8,13,13	0.78	0
8	GOL	B	701	-	5,5,5	0.35	0	5,5,5	0.48	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	SO4	A	1004	-	-	0/0/0/0	0/0/0/0
7	POP	A	1005	4	-	0/6/6/6	0/0/0/0
8	GOL	B	701	-	-	0/4/4/4	0/0/0/0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	A	1005	POP	P2-O	2.72	1.64	1.60

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	A	1005	POP	1	0
8	B	701	GOL	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	462/468 (98%)	0.01	10 (2%) 62 54	45, 59, 83, 93	0
2	B	12/35 (34%)	0.77	1 (8%) 12 7	58, 63, 113, 118	0
3	C	10/19 (52%)	0.94	2 (20%) 1 1	62, 67, 114, 123	0
All	All	484/522 (92%)	0.04	13 (2%) 55 45	45, 60, 84, 123	0

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	C	693	A	4.2
2	B	610	C	3.9
1	A	131	LEU	3.9
3	C	694	G	3.1
1	A	52	VAL	3.0
1	A	18	ASN	2.9
1	A	50	LEU	2.9
1	A	277	ARG	2.5
1	A	295	SER	2.5
1	A	48	PRO	2.3
1	A	144	TYR	2.2
1	A	51	GLU	2.1
1	A	279	LYS	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.



## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
4	MG	A	1001	1/1	0.93	0.39	3.16	56,56,56,56	1
7	POP	A	1005	9/9	0.84	0.36	3.04	42,52,67,70	9
8	GOL	B	701	6/6	0.97	0.25	0.83	54,59,61,62	0
6	SO4	A	1004	5/5	0.97	0.20	0.18	49,50,52,53	5
5	ZN	A	1003	1/1	0.99	0.15	-0.74	39,39,39,39	1
4	MG	A	1002	1/1	0.96	0.33	-	51,51,51,51	1

## 6.5 Other polymers [i](#)

There are no such residues in this entry.