



Full wwPDB NMR Structure Validation Report ⓘ

Feb 12, 2017 – 06:02 pm GMT

PDB ID : 1F9X
Title : AVERAGE NMR SOLUTION STRUCTURE OF THE BIR-3 DOMAIN OF XIAP
Authors : Sun, C.; Cai, M.; Meadows, R.P.; Fesik, S.W.
Deposited on : 2000-07-11

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)
NmrClust : Kelley et al. (1996)
MolProbity : 4.02b-467
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : trunk28760
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR


The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	125131	11601
Ramachandran outliers	121729	10391
Sidechain outliers	121581	10367

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	120	

2 Ensemble composition and analysis ⓘ

This entry contains 1 models. Identification of well-defined residues and clustering analysis are not possible.

3 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 1828 atoms, of which 882 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called INHIBITOR OF APOPTOSIS PROTEIN XIAP.

Mol	Chain	Residues	Atoms						Trace
1	A	117	Total	C	H	N	O	S	0
			1827	597	882	162	180	6	

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	237	GLY	-	CLONING ARTIFACT	UNP P98170
A	238	SER	-	CLONING ARTIFACT	UNP P98170
A	239	HIS	-	CLONING ARTIFACT	UNP P98170
A	240	MET	-	CLONING ARTIFACT	UNP P98170

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms	
2	A	1	Total	Zn
			1	1

4 Residue-property plots [i](#)

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

• Molecule 1: INHIBITOR OF APOPTOSIS PROTEIN XIAP

Chain A: 



5 Refinement protocol and experimental data overview ⓘ

The models were refined using the following method: *simulated annealing*.

Of the ? calculated structures, 1 were deposited, based on the following criterion: ?.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CNX	structure solution	
CNX	refinement	

No chemical shift data was provided. No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality ⓘ

6.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section:
ZN

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	945	882	881	45
All	All	946	882	881	45

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 25.

All clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)
1:A:272:PHE:CD1	1:A:291:ALA:HB2	0.89	2.02
1:A:263:ALA:HB1	1:A:268:ARG:NH2	0.69	2.03
1:A:267:ALA:O	1:A:271:THR:HG23	0.65	1.91
1:A:321:ALA:HB3	1:A:342:ILE:HG21	0.63	1.71
1:A:276:ILE:HG22	1:A:296:ASP:HA	0.61	1.73
1:A:321:ALA:HB1	1:A:339:ILE:HG23	0.59	1.74
1:A:276:ILE:HD11	1:A:280:ASN:CB	0.57	2.29
1:A:339:ILE:N	1:A:339:ILE:HD13	0.57	2.14
1:A:292:LEU:HD21	1:A:299:LYS:HB3	0.57	1.76
1:A:276:ILE:HD11	1:A:280:ASN:HB2	0.56	1.77
1:A:344:LEU:O	1:A:345:THR:HG23	0.56	2.00
1:A:272:PHE:CE1	1:A:291:ALA:HB2	0.55	2.35
1:A:292:LEU:N	1:A:292:LEU:HD13	0.55	2.16

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)
1:A:331:LEU:HD11	1:A:336:GLN:HG3	0.54	1.79
1:A:260:PRO:HA	1:A:263:ALA:HB3	0.54	1.78
1:A:352:LEU:O	1:A:353:VAL:HG23	0.54	2.03
1:A:272:PHE:CZ	1:A:284:LEU:HD13	0.50	2.41
1:A:254:THR:HG22	1:A:257:PRO:HG2	0.50	1.83
1:A:290:TYR:CE1	1:A:292:LEU:CD1	0.50	2.95
1:A:284:LEU:HD22	1:A:289:PHE:HB3	0.49	1.85
1:A:297:LYS:HB3	1:A:308:THR:HG22	0.49	1.85
1:A:279:VAL:O	1:A:279:VAL:HG12	0.48	2.09
1:A:292:LEU:HD11	1:A:299:LYS:HB3	0.47	1.86
1:A:352:LEU:HD12	1:A:353:VAL:HB	0.47	1.87
1:A:321:ALA:CB	1:A:339:ILE:HG23	0.47	2.39
1:A:317:TRP:CZ3	1:A:330:LEU:CD2	0.45	2.99
1:A:289:PHE:CD2	1:A:298:VAL:CG2	0.45	3.00
1:A:263:ALA:HB1	1:A:268:ARG:HH21	0.45	1.70
1:A:317:TRP:CZ2	1:A:334:LYS:CG	0.45	2.99
1:A:306:GLY:O	1:A:323:TRP:CZ2	0.44	2.70
1:A:353:VAL:HG12	1:A:354:ARG:N	0.44	2.28
1:A:289:PHE:CD2	1:A:298:VAL:HG21	0.44	2.48
1:A:272:PHE:O	1:A:275:TRP:CE3	0.43	2.71
1:A:307:LEU:CD2	1:A:323:TRP:CD1	0.43	3.01
1:A:269:ILE:HD11	1:A:281:LYS:HA	0.43	1.89
1:A:262:MET:O	1:A:267:ALA:HB1	0.43	2.13
1:A:297:LYS:N	1:A:297:LYS:CD	0.43	2.82
1:A:352:LEU:HD12	1:A:352:LEU:C	0.43	2.34
1:A:317:TRP:CZ2	1:A:334:LYS:HG3	0.42	2.50
1:A:265:TYR:CD2	1:A:285:ALA:HB1	0.42	2.50
1:A:277:TYR:CD1	1:A:277:TYR:N	0.41	2.87
1:A:346:HIS:CE1	1:A:347:SER:OG	0.41	2.74
1:A:248:ARG:O	1:A:250:PHE:CD2	0.41	2.73
1:A:295:GLY:O	1:A:296:ASP:CB	0.41	2.69
1:A:313:SER:O	1:A:314:GLU:C	0.41	2.58

6.3 Torsion angles ⓘ

6.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	115/120 (96%)	73 (63%)	21 (18%)	21 (18%)	0	3
All	All	115/120 (96%)	73 (63%)	21 (18%)	21 (18%)	0	3

All 21 Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type
1	A	313	SER
1	A	353	VAL
1	A	248	ARG
1	A	243	ALA
1	A	281	LYS
1	A	344	LEU
1	A	328	LYS
1	A	277	TYR
1	A	279	VAL
1	A	264	ASP
1	A	345	THR
1	A	312	PRO
1	A	348	LEU
1	A	354	ARG
1	A	257	PRO
1	A	314	GLU
1	A	252	ASN
1	A	335	GLY
1	A	324	TYR
1	A	347	SER
1	A	276	ILE

6.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	101/103 (98%)	66 (65%)	35 (35%)	1	10
All	All	101/103 (98%)	66 (65%)	35 (35%)	1	10

All 35 residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type
1	A	354	ARG
1	A	311	LYS
1	A	253	SER
1	A	327	CYS
1	A	309	ASP
1	A	344	LEU
1	A	294	GLU
1	A	328	LYS
1	A	254	THR
1	A	353	VAL
1	A	299	LYS
1	A	336	GLN
1	A	284	LEU
1	A	303	CYS
1	A	352	LEU
1	A	248	ARG
1	A	317	TRP
1	A	345	THR
1	A	350	GLU
1	A	307	LEU
1	A	323	TRP
1	A	324	TYR
1	A	262	MET
1	A	347	SER
1	A	274	THR
1	A	280	ASN
1	A	290	TYR
1	A	302	HIS
1	A	334	LYS
1	A	356	THR
1	A	292	LEU
1	A	264	ASP
1	A	278	SER
1	A	286	ARG
1	A	246	SER

6.3.3 RNA ⓘ

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

No chemical shift data were provided