



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 12, 2017 – 08:19 pm GMT

PDB ID : 1FAJ
Title : INORGANIC PYROPHOSPHATASE
Authors : Kankare, J.A.; Salminen, T.; Goldman, A.
Deposited on : 1996-01-10
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	112137	1278 (2.16-2.16)
Ramachandran outliers	110173	1256 (2.16-2.16)
Sidechain outliers	110143	1255 (2.16-2.16)
RSRZ outliers	101464	1175 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	175	<div> <div>90%</div> <div>70%23% . .</div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 1408 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SOLUBLE INORGANIC PYROPHOSPHATASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	168	Total	C	N	O	S	0	0	0
			1327	857	213	252	5			


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	81	Total	O	0	0
			81	81		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: SOLUBLE INORGANIC PYROPHOSPHATASE



4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α , β , γ	111.50Å 111.50Å 76.50Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	8.00 – 2.15 7.98 – 2.15	Depositor EDS
% Data completeness (in resolution range)	99.9 (8.00-2.15) 29.4 (7.98-2.15)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	8.90 (at 2.14Å)	Xtriage
Refinement program	X-PLOR 3.1	Depositor
R, R_{free}	0.187 , (Not available) 0.533 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	39.4	Xtriage
Anisotropy	1.275	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	-1.81 , 86.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.53$, $\langle L^2 \rangle = 0.37$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.46	EDS
Total number of atoms	1408	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.51% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.87	0/1360	0.97	4/1849 (0.2%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	126	LEU	CA-CB-CG	6.60	130.47	115.30
1	A	103	ALA	N-CA-C	-6.01	94.77	111.00
1	A	113	LEU	CA-CB-CG	5.54	128.04	115.30
1	A	140	HIS	N-CA-C	5.10	124.77	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1327	0	1308	28	1
2	A	81	0	0	6	1
All	All	1408	0	1308	28	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:96:THR:HG22	1:A:100:GLY:O	1.74	0.86
1:A:9:LYS:HE2	1:A:86:ARG:NH2	2.05	0.72
1:A:118:ASP:O	1:A:121:LYS:HE2	1.90	0.71
1:A:111:SER:HB3	2:A:247:HOH:O	1.91	0.70
1:A:32:ILE:HD13	1:A:39:LEU:HD23	1.73	0.70
1:A:60:HIS:HE1	2:A:228:HOH:O	1.85	0.58
1:A:32:ILE:CD1	1:A:39:LEU:HD23	2.33	0.58
1:A:43:ARG:NH1	1:A:45:MET:HA	2.20	0.56
1:A:43:ARG:HH12	1:A:45:MET:HA	1.71	0.55
1:A:92:VAL:HG13	1:A:156:GLU:HB2	1.88	0.53
1:A:2:LEU:HD12	1:A:32:ILE:HD12	1.90	0.53
1:A:151:LYS:HD2	1:A:152:VAL:N	2.24	0.53
1:A:30:TYR:CD1	1:A:39:LEU:HD13	2.45	0.52
1:A:63:SER:HB2	2:A:235:HOH:O	2.08	0.52
1:A:88:ARG:HD3	2:A:195:HOH:O	2.10	0.51
1:A:117:TYR:HD2	1:A:120:ILE:HD12	1.78	0.47
1:A:64:LEU:C	1:A:66:GLY:H	2.18	0.47
1:A:8:GLY:HA3	1:A:15:ILE:HA	1.97	0.46
1:A:95:MET:HE1	1:A:138:PHE:CB	2.46	0.46
1:A:124:ASN:HA	1:A:131:LYS:NZ	2.31	0.45
1:A:2:LEU:CD1	1:A:32:ILE:HD12	2.46	0.45
1:A:95:MET:HE1	1:A:138:PHE:HB3	1.99	0.44
1:A:88:ARG:HB3	2:A:252:HOH:O	2.17	0.44
1:A:59:ASN:O	1:A:60:HIS:HB2	2.18	0.43
1:A:9:LYS:HE3	1:A:9:LYS:HB3	1.84	0.43
1:A:127:PRO:O	1:A:131:LYS:HG3	2.17	0.43
1:A:106:VAL:HG11	1:A:126:LEU:HD11	2.00	0.43
1:A:25:ALA:HB1	2:A:198:HOH:O	2.21	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:80:GLN:NE2	2:A:245:HOH:O[2_555]	2.07	0.13

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	162/175 (93%)	155 (96%)	7 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	145/150 (97%)	136 (94%)	9 (6%)	21	15

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	12	PRO
1	A	39	LEU
1	A	93	LEU
1	A	96	THR
1	A	101	GLU
1	A	102	ASP
1	A	124	ASN
1	A	141	TYR
1	A	145	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	4	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	168/175 (96%)	6.67	158 (94%) 0 0	15, 30, 60, 81	0

All (158) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	71	VAL	22.2
1	A	96	THR	21.7
1	A	63	SER	21.3
1	A	21	ILE	19.8
1	A	158	ALA	18.3
1	A	120	ILE	14.1
1	A	6	PRO	13.7
1	A	32	ILE	13.6
1	A	5	VAL	13.5
1	A	12	PRO	13.2
1	A	111	SER	13.1
1	A	136	HIS	12.9
1	A	125	ASP	11.9
1	A	113	LEU	11.6
1	A	17	VAL	11.5
1	A	60	HIS	11.4
1	A	87	CYS	11.4
1	A	95	MET	11.3
1	A	152	VAL	11.3
1	A	74	PRO	10.9
1	A	52	PRO	10.9
1	A	104	LYS	10.8
1	A	157	ASN	10.7
1	A	53	CYS	10.7
1	A	166	VAL	10.6
1	A	141	TYR	10.5
1	A	36	SER	10.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	22	PRO	10.0
1	A	129	LEU	9.8
1	A	29	LYS	9.3
1	A	149	TRP	9.3
1	A	18	VAL	9.2
1	A	170	GLU	9.1
1	A	14	ASP	9.1
1	A	54	ASN	9.0
1	A	107	ALA	9.0
1	A	154	GLY	9.0
1	A	49	MET	8.9
1	A	62	LEU	8.8
1	A	109	PRO	8.8
1	A	88	ARG	8.7
1	A	50	PHE	8.7
1	A	93	LEU	8.7
1	A	110	HIS	8.6
1	A	3	LEU	8.6
1	A	67	ASP	8.5
1	A	117	TYR	8.3
1	A	103	ALA	8.3
1	A	160	ALA	8.3
1	A	28	ILE	8.2
1	A	80	GLN	8.2
1	A	123	VAL	8.1
1	A	169	PHE	8.1
1	A	94	LYS	8.0
1	A	105	LEU	8.0
1	A	171	ARG	7.9
1	A	106	VAL	7.8
1	A	25	ALA	7.7
1	A	85	ILE	7.6
1	A	102	ASP	7.6
1	A	126	LEU	7.4
1	A	43	ARG	7.4
1	A	48	ALA	7.1
1	A	137	PHE	7.1
1	A	101	GLU	7.0
1	A	31	GLU	7.0
1	A	2	LEU	6.9
1	A	100	GLY	6.9
1	A	130	LEU	6.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	57	TYR	6.8
1	A	1	SER	6.6
1	A	140	HIS	6.5
1	A	172	ALA	6.4
1	A	76	PRO	6.4
1	A	119	HIS	6.4
1	A	161	ALA	6.4
1	A	135	ALA	6.3
1	A	165	ILE	6.2
1	A	168	SER	6.1
1	A	10	ASP	6.1
1	A	77	TYR	6.1
1	A	118	ASP	6.0
1	A	64	LEU	6.0
1	A	108	VAL	6.0
1	A	55	TYR	5.8
1	A	23	ALA	5.8
1	A	40	PHE	5.8
1	A	56	GLY	5.7
1	A	51	TYR	5.7
1	A	4	ASN	5.7
1	A	114	SER	5.6
1	A	19	ILE	5.5
1	A	16	TYR	5.5
1	A	37	GLY	5.5
1	A	86	ARG	5.4
1	A	30	TYR	5.3
1	A	65	ASP	5.3
1	A	33	ASP	5.3
1	A	97	ASP	5.2
1	A	68	PRO	5.2
1	A	92	VAL	5.2
1	A	150	VAL	5.1
1	A	159	GLU	5.1
1	A	115	LYS	5.1
1	A	20	GLU	4.9
1	A	27	PRO	4.9
1	A	15	ILE	4.9
1	A	121	LYS	4.8
1	A	162	LYS	4.7
1	A	167	ALA	4.7
1	A	124	ASN	4.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	47	THR	4.6
1	A	90	VAL	4.5
1	A	46	SER	4.5
1	A	11	LEU	4.4
1	A	145	GLU	4.4
1	A	156	GLU	4.4
1	A	44	PHE	4.1
1	A	66	GLY	4.1
1	A	133	GLN	4.1
1	A	138	PHE	4.1
1	A	122	ASP	4.0
1	A	143	ASP	3.9
1	A	134	ILE	3.9
1	A	89	PRO	3.8
1	A	112	LYS	3.8
1	A	131	LYS	3.7
1	A	26	ASP	3.7
1	A	42	ASP	3.7
1	A	39	LEU	3.7
1	A	24	ASN	3.7
1	A	58	ILE	3.6
1	A	132	ALA	3.6
1	A	163	ALA	3.5
1	A	144	LEU	3.4
1	A	173	LYS	3.3
1	A	72	LEU	3.3
1	A	45	MET	3.2
1	A	41	VAL	3.2
1	A	155	TRP	3.2
1	A	59	ASN	3.2
1	A	79	LEU	3.2
1	A	8	GLY	3.1
1	A	75	THR	3.0
1	A	151	LYS	2.7
1	A	164	GLU	2.7
1	A	128	GLU	2.7
1	A	35	GLU	2.7
1	A	153	GLU	2.7
1	A	91	GLY	2.6
1	A	34	LYS	2.6
1	A	139	GLU	2.6
1	A	70	ASP	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	9	LYS	2.5
1	A	84	VAL	2.5
1	A	116	GLU	2.5
1	A	82	GLY	2.3
1	A	78	PRO	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.