



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 14, 2017 – 05:49 am GMT

PDB ID : 3FBE
Title : Crystal structure of the Mimivirus NDK N62L-R107G double mutant complexed with GDP
Authors : Jeudy, S.; Lartigue, A.; Claverie, J.M.; Abergel, C.
Deposited on : 2008-11-19
Resolution : 2.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	3166 (2.40-2.40)
Clashscore	112137	3674 (2.40-2.40)
Ramachandran outliers	110173	3616 (2.40-2.40)
Sidechain outliers	110143	3617 (2.40-2.40)
RSRZ outliers	101464	3195 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	142	<div> <div>4%</div> <div> <div></div> <div>75%</div> <div>18%</div> <div>6%</div> </div> </div>
1	B	142	<div> <div></div> <div> <div>73%</div> <div>16%</div> <div>• 9%</div> </div> </div>
1	C	142	<div> <div>%</div> <div> <div></div> <div>75%</div> <div>15%</div> <div>• • 7%</div> </div> </div>
1	D	142	<div> <div>%</div> <div> <div></div> <div>71%</div> <div>18%</div> <div>• 7%</div> </div> </div>
1	E	142	<div> <div>6%</div> <div> <div></div> <div>70%</div> <div>19%</div> <div>• 7%</div> </div> </div>
1	F	142	<div> <div>4%</div> <div> <div></div> <div>64%</div> <div>24%</div> <div>• 8%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GDP	A	138	-	-	-	X
3	GDP	B	138	-	-	-	X
3	GDP	C	138	-	-	-	X
3	GDP	D	138	-	-	-	X
3	GDP	E	138	-	-	-	X
3	GDP	F	138	-	-	-	X

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 6610 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nucleoside diphosphate kinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	133	Total	C	N	O	S	0	0	0
			1058	671	180	203	4			
1	B	129	Total	C	N	O	S	0	0	0
			1033	657	176	196	4			
1	C	132	Total	C	N	O	S	0	0	0
			1053	668	179	202	4			
1	D	132	Total	C	N	O	S	0	0	0
			1053	668	179	202	4			
1	E	132	Total	C	N	O	S	0	0	0
			1053	668	179	202	4			
1	F	130	Total	C	N	O	S	0	0	0
			1037	659	177	197	4			

There are 48 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-4	TYR	-	EXPRESSION TAG	UNP Q5UQL3
A	-3	LYS	-	EXPRESSION TAG	UNP Q5UQL3
A	-2	LYS	-	EXPRESSION TAG	UNP Q5UQL3
A	-1	ALA	-	EXPRESSION TAG	UNP Q5UQL3
A	0	GLY	-	EXPRESSION TAG	UNP Q5UQL3
A	1	LEU	-	EXPRESSION TAG	UNP Q5UQL3
A	62	LEU	ASN	ENGINEERED	UNP Q5UQL3
A	107	GLY	ARG	ENGINEERED	UNP Q5UQL3
B	-4	TYR	-	EXPRESSION TAG	UNP Q5UQL3
B	-3	LYS	-	EXPRESSION TAG	UNP Q5UQL3
B	-2	LYS	-	EXPRESSION TAG	UNP Q5UQL3
B	-1	ALA	-	EXPRESSION TAG	UNP Q5UQL3
B	0	GLY	-	EXPRESSION TAG	UNP Q5UQL3
B	1	LEU	-	EXPRESSION TAG	UNP Q5UQL3
B	62	LEU	ASN	ENGINEERED	UNP Q5UQL3
B	107	GLY	ARG	ENGINEERED	UNP Q5UQL3
C	-4	TYR	-	EXPRESSION TAG	UNP Q5UQL3

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	-3	LYS	-	EXPRESSION TAG	UNP Q5UQL3
C	-2	LYS	-	EXPRESSION TAG	UNP Q5UQL3
C	-1	ALA	-	EXPRESSION TAG	UNP Q5UQL3
C	0	GLY	-	EXPRESSION TAG	UNP Q5UQL3
C	1	LEU	-	EXPRESSION TAG	UNP Q5UQL3
C	62	LEU	ASN	ENGINEERED	UNP Q5UQL3
C	107	GLY	ARG	ENGINEERED	UNP Q5UQL3
D	-4	TYR	-	EXPRESSION TAG	UNP Q5UQL3
D	-3	LYS	-	EXPRESSION TAG	UNP Q5UQL3
D	-2	LYS	-	EXPRESSION TAG	UNP Q5UQL3
D	-1	ALA	-	EXPRESSION TAG	UNP Q5UQL3
D	0	GLY	-	EXPRESSION TAG	UNP Q5UQL3
D	1	LEU	-	EXPRESSION TAG	UNP Q5UQL3
D	62	LEU	ASN	ENGINEERED	UNP Q5UQL3
D	107	GLY	ARG	ENGINEERED	UNP Q5UQL3
E	-4	TYR	-	EXPRESSION TAG	UNP Q5UQL3
E	-3	LYS	-	EXPRESSION TAG	UNP Q5UQL3
E	-2	LYS	-	EXPRESSION TAG	UNP Q5UQL3
E	-1	ALA	-	EXPRESSION TAG	UNP Q5UQL3
E	0	GLY	-	EXPRESSION TAG	UNP Q5UQL3
E	1	LEU	-	EXPRESSION TAG	UNP Q5UQL3
E	62	LEU	ASN	ENGINEERED	UNP Q5UQL3
E	107	GLY	ARG	ENGINEERED	UNP Q5UQL3
F	-4	TYR	-	EXPRESSION TAG	UNP Q5UQL3
F	-3	LYS	-	EXPRESSION TAG	UNP Q5UQL3
F	-2	LYS	-	EXPRESSION TAG	UNP Q5UQL3
F	-1	ALA	-	EXPRESSION TAG	UNP Q5UQL3
F	0	GLY	-	EXPRESSION TAG	UNP Q5UQL3
F	1	LEU	-	EXPRESSION TAG	UNP Q5UQL3
F	62	LEU	ASN	ENGINEERED	UNP Q5UQL3
F	107	GLY	ARG	ENGINEERED	UNP Q5UQL3

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Mg 1 1	0	0
2	D	1	Total Mg 1 1	0	0
2	C	1	Total Mg 1 1	0	0
2	F	1	Total Mg 1 1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	E	1	Total	Mg	0	0
			1	1		

- Molecule 3 is GUANOSINE-5'-DIPHOSPHATE (three-letter code: GDP) (formula: $C_{10}H_{15}N_5O_{11}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			28	10	5	11	2		
3	B	1	Total	C	N	O	P	0	0
			28	10	5	11	2		
3	C	1	Total	C	N	O	P	0	0
			28	10	5	11	2		
3	D	1	Total	C	N	O	P	0	0
			28	10	5	11	2		
3	E	1	Total	C	N	O	P	0	0
			28	10	5	11	2		
3	F	1	Total	C	N	O	P	0	0
			28	10	5	11	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	30	Total	O	0	0
			30	30		
4	B	17	Total	O	0	0
			17	17		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	C	25	Total 25	O 25	0	0
4	D	25	Total 25	O 25	0	0
4	E	28	Total 28	O 28	0	0
4	F	25	Total 25	O 25	0	0



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	79.24Å 152.90Å 183.78Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	28.53 – 2.40 28.53 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.8 (28.53-2.40) 99.8 (28.53-2.40)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.52 (at 2.39Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.190 , 0.230 0.185 , 0.222	Depositor DCC
R_{free} test set	2212 reflections (5.04%)	DCC
Wilson B-factor (Å ²)	41.0	Xtriage
Anisotropy	0.071	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 40.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6610	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.18% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GDP, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.33	0/1078	0.54	0/1455
1	B	0.37	0/1053	0.54	0/1421
1	C	0.36	0/1073	0.53	0/1448
1	D	0.32	0/1073	0.50	0/1448
1	E	0.35	0/1073	0.55	0/1448
1	F	0.34	0/1057	0.53	0/1426
All	All	0.35	0/6407	0.53	0/8646

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1058	0	1055	14	0
1	B	1033	0	1034	20	0
1	C	1053	0	1050	19	0
1	D	1053	0	1050	19	0
1	E	1053	0	1050	20	0
1	F	1037	0	1037	30	0
2	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	C	1	0	0	0	0
2	D	1	0	0	0	0
2	E	1	0	0	0	0
2	F	1	0	0	0	0
3	A	28	0	12	6	0
3	B	28	0	12	7	0
3	C	28	0	12	3	0
3	D	28	0	12	5	0
3	E	28	0	12	6	0
3	F	28	0	12	2	0
4	A	30	0	0	1	0
4	B	17	0	0	0	0
4	C	25	0	0	0	0
4	D	25	0	0	0	0
4	E	28	0	0	1	0
4	F	25	0	0	1	0
All	All	6610	0	6348	131	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 131 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:87:ARG:HG2	1:F:87:ARG:HH11	1.08	1.09
1:B:53:HIS:HE1	3:B:138:GDP:H2'	1.18	1.04
1:B:53:HIS:CE1	3:B:138:GDP:H2'	2.02	0.93
3:A:138:GDP:H5'	3:A:138:GDP:H8	1.41	0.85
1:F:87:ARG:CG	1:F:87:ARG:HH11	1.91	0.84

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	131/142 (92%)	126 (96%)	5 (4%)	0	100	100
1	B	127/142 (89%)	123 (97%)	4 (3%)	0	100	100
1	C	130/142 (92%)	125 (96%)	4 (3%)	1 (1%)	22	33
1	D	130/142 (92%)	126 (97%)	3 (2%)	1 (1%)	22	33
1	E	130/142 (92%)	123 (95%)	6 (5%)	1 (1%)	22	33
1	F	128/142 (90%)	124 (97%)	4 (3%)	0	100	100
All	All	776/852 (91%)	747 (96%)	26 (3%)	3 (0%)	38	54

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	130	GLU
1	C	130	GLU
1	E	130	GLU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	117/126 (93%)	111 (95%)	6 (5%)	28	44
1	B	115/126 (91%)	108 (94%)	7 (6%)	22	34
1	C	117/126 (93%)	111 (95%)	6 (5%)	28	44
1	D	117/126 (93%)	109 (93%)	8 (7%)	18	29
1	E	117/126 (93%)	106 (91%)	11 (9%)	10	15
1	F	115/126 (91%)	107 (93%)	8 (7%)	18	28
All	All	698/756 (92%)	652 (93%)	46 (7%)	19	30

5 of 46 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	5	LEU
1	D	94	THR
1	F	86	ARG
1	D	29	ASN
1	D	67	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	89	GLN
1	F	48	GLN
1	E	48	GLN
1	C	29	ASN
1	E	29	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 5 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GDP	A	138	-	25,30,30	1.21	2 (8%)	26,47,47	2.18	7 (26%)
3	GDP	B	138	-	25,30,30	1.25	2 (8%)	26,47,47	2.06	7 (26%)
3	GDP	C	138	-	25,30,30	1.26	3 (12%)	26,47,47	2.12	6 (23%)
3	GDP	D	138	-	25,30,30	1.28	2 (8%)	26,47,47	2.09	6 (23%)
3	GDP	E	138	-	25,30,30	1.24	2 (8%)	26,47,47	2.33	8 (30%)
3	GDP	F	138	-	25,30,30	1.25	2 (8%)	26,47,47	2.21	6 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GDP	A	138	-	-	0/12/32/32	0/3/3/3
3	GDP	B	138	-	-	0/12/32/32	0/3/3/3
3	GDP	C	138	-	-	0/12/32/32	0/3/3/3
3	GDP	D	138	-	-	0/12/32/32	0/3/3/3
3	GDP	E	138	-	-	0/12/32/32	0/3/3/3
3	GDP	F	138	-	-	0/12/32/32	0/3/3/3

The worst 5 of 13 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	138	GDP	PB-O3A	2.13	1.63	1.60
3	E	138	GDP	C5-C4	2.95	1.47	1.40
3	F	138	GDP	C5-C4	2.99	1.47	1.40
3	A	138	GDP	C5-C4	3.08	1.47	1.40
3	B	138	GDP	C5-C4	3.19	1.47	1.40

The worst 5 of 40 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	E	138	GDP	C6-C5-C4	-5.36	115.51	120.84
3	A	138	GDP	C6-C5-C4	-4.73	116.14	120.84
3	F	138	GDP	C6-C5-C4	-4.65	116.22	120.84
3	D	138	GDP	C6-C5-C4	-4.45	116.42	120.84
3	C	138	GDP	C6-C5-C4	-4.21	116.66	120.84

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6 monomers are involved in 29 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	138	GDP	6	0
3	B	138	GDP	7	0
3	C	138	GDP	3	0
3	D	138	GDP	5	0
3	E	138	GDP	6	0
3	F	138	GDP	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	133/142 (93%)	-0.30	5 (3%) 41 40	28, 39, 66, 80	0
1	B	129/142 (90%)	-0.53	0 100 100	25, 37, 56, 69	0
1	C	132/142 (92%)	-0.40	1 (0%) 86 84	28, 38, 57, 68	0
1	D	132/142 (92%)	-0.21	2 (1%) 74 72	32, 48, 66, 84	0
1	E	132/142 (92%)	-0.14	8 (6%) 22 20	28, 39, 67, 86	0
1	F	130/142 (91%)	-0.23	6 (4%) 33 31	30, 43, 64, 80	0
All	All	788/852 (92%)	-0.30	22 (2%) 53 51	25, 41, 64, 86	0

The worst 5 of 22 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	131	THR	4.3
1	E	130	GLU	4.0
1	A	-1	ALA	3.5
1	A	55	GLU	3.3
1	D	131	THR	2.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	GDP	B	138	28/28	0.44	0.50	7.20	100,108,118,123	0
3	GDP	E	138	28/28	0.78	0.36	6.28	69,76,96,109	0
3	GDP	D	138	28/28	0.78	0.31	3.76	57,73,94,108	0
3	GDP	C	138	28/28	0.79	0.31	3.60	62,72,92,94	0
3	GDP	A	138	28/28	0.83	0.31	3.52	57,73,88,101	0
3	GDP	F	138	28/28	0.76	0.35	3.48	71,82,105,114	0
2	MG	D	163	1/1	0.89	0.14	-0.65	59,59,59,59	0
2	MG	F	165	1/1	0.84	0.14	-	77,77,77,77	0
2	MG	A	160	1/1	0.94	0.11	-	54,54,54,54	0
2	MG	C	162	1/1	0.77	0.26	-	70,70,70,70	0
2	MG	E	164	1/1	0.53	0.35	-	78,78,78,78	0

6.5 Other polymers [i](#)

There are no such residues in this entry.