



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 02:35 am GMT

PDB ID : 1FCV  
Title : CRYSTAL STRUCTURE OF BEE VENOM HYALURONIDASE IN COM-  
PLEX WITH HYALURONIC ACID TETRAMER  
Authors : Markovic-Housley, Z.; Miglierini, G.; Soldatova, L.; Rizkallah, P.J.; Mueller,  
U.; Schirmer, T.  
Deposited on : 2000-07-19  
Resolution : 2.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	3491 (2.70-2.62)
Clashscore	112137	1026 (2.68-2.64)
Ramachandran outliers	110173	1010 (2.68-2.64)
Sidechain outliers	110143	1010 (2.68-2.64)
RSRZ outliers	101464	3511 (2.70-2.62)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	350	<div> <div>2%</div> <div>69%</div> <div>18%</div> <div>• • 7%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GCU	A	401	X	-	-	-
2	NAG	A	402	-	-	-	X

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GCU	A	403	X	-	-	-
2	NAG	A	404	X	-	-	X

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2866 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HYALURONOGLUCOSAMINIDASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	324	Total	C	N	O	S	45	0	0
			2681	1717	470	482	12			

- Molecule 2 is a polymer of unknown type called SUGAR (5-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	5	Total	C	N	O	0	0
			53	28	2	23		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	132	Total	O	0	0
			132	132		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: HYALURONOGLUCOSAMINIDASE



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	71.01Å 71.01Å 151.33Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	8.00 – 2.65 20.50 – 2.65	Depositor EDS
% Data completeness (in resolution range)	(Not available) (8.00-2.65) 99.8 (20.50-2.65)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.36 (at 2.63Å)	Xtriage
Refinement program	REFMAC	Depositor
R, $R_{free}$	0.163 , 0.252 0.162 , 0.232	Depositor DCC
$R_{free}$ test set	1340 reflections (11.15%)	DCC
Wilson B-factor (Å <sup>2</sup> )	24.3	Xtriage
Anisotropy	0.392	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 81.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.042 for -h,-k,l	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	2866	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.05% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GCU, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.59	2/2753 (0.1%)	1.48	35/3726 (0.9%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
2	A	3	0
All	All	3	2

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	332	ALA	C-N	-11.12	1.08	1.34
1	A	10	GLU	CB-CG	7.84	1.67	1.52

All (35) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	332	ALA	O-C-N	-18.15	93.66	122.70
1	A	250	ARG	NE-CZ-NH2	13.96	127.28	120.30
1	A	244	ARG	NE-CZ-NH1	-13.74	113.43	120.30
1	A	332	ALA	CA-C-N	10.88	141.14	117.20
1	A	317	ARG	NE-CZ-NH2	-10.63	114.99	120.30
1	A	116	ARG	NE-CZ-NH2	-9.66	115.47	120.30
1	A	250	ARG	NE-CZ-NH1	-9.07	115.76	120.30
1	A	160	TYR	CB-CG-CD2	-8.82	115.71	121.00
1	A	229	ARG	NE-CZ-NH2	-8.60	116.00	120.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	28	ARG	NE-CZ-NH1	8.43	124.52	120.30
1	A	47	ARG	NE-CZ-NH2	-8.11	116.25	120.30
1	A	155	ARG	NE-CZ-NH1	8.00	124.30	120.30
1	A	123	TRP	CA-CB-CG	7.71	128.35	113.70
1	A	244	ARG	NH1-CZ-NH2	7.70	127.87	119.40
1	A	28	ARG	CD-NE-CZ	7.62	134.28	123.60
1	A	256	THR	N-CA-CB	-7.07	96.87	110.30
1	A	273	ARG	NE-CZ-NH1	-7.07	116.77	120.30
1	A	280	ARG	CG-CD-NE	7.00	126.51	111.80
1	A	176	ARG	CD-NE-CZ	6.81	133.13	123.60
1	A	44	ASP	CB-CG-OD2	6.80	124.42	118.30
1	A	176	ARG	NE-CZ-NH1	-6.57	117.02	120.30
1	A	183	TYR	CB-CG-CD2	-6.45	117.13	121.00
1	A	288	ARG	CA-CB-CG	6.05	126.72	113.40
1	A	155	ARG	NE-CZ-NH2	-6.04	117.28	120.30
1	A	305	ASP	CB-CG-OD1	5.75	123.47	118.30
1	A	93	ASP	CB-CG-OD1	-5.68	113.19	118.30
1	A	148	ARG	CD-NE-CZ	5.57	131.40	123.60
1	A	236	GLU	CG-CD-OE1	-5.55	107.21	118.30
1	A	93	ASP	CB-CG-OD2	5.54	123.29	118.30
1	A	272	ASP	CB-CG-OD2	5.52	123.27	118.30
1	A	236	GLU	CG-CD-OE2	5.52	129.34	118.30
1	A	47	ARG	NH1-CZ-NH2	5.36	125.29	119.40
1	A	66	ASP	CB-CG-OD2	5.16	122.94	118.30
1	A	115	TRP	CB-CG-CD2	5.14	133.28	126.60
1	A	269	LYS	CA-CB-CG	5.01	124.42	113.40

All (3) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	A	401	GCU	C1
2	A	403	GCU	C1
2	A	404	NAG	C1

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	101	ASP	Mainchain
1	A	66	ASP	Mainchain



## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2681	0	2625	46	1
2	A	53	0	37	0	0
3	A	132	0	0	4	0
All	All	2866	0	2662	46	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (46) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:193:THR:HG22	1:A:196:GLN:H	1.30	0.94
1:A:75:ASN:HD21	1:A:83:ASN:H	1.12	0.89
1:A:80:GLN:HE21	1:A:80:GLN:H	0.91	0.88
1:A:80:GLN:NE2	1:A:80:GLN:H	1.73	0.87
1:A:80:GLN:HE21	1:A:80:GLN:N	1.74	0.84
1:A:59:MET:H	1:A:87:HIS:HD2	1.23	0.83
1:A:64:LEU:HB2	1:A:72:VAL:HG23	1.67	0.76
1:A:246:LYS:HE3	1:A:293:LEU:HD21	1.66	0.75
1:A:75:ASN:ND2	1:A:83:ASN:H	1.87	0.72
1:A:88:LEU:O	1:A:92:ARG:HG3	1.96	0.66
1:A:66:ASP:HB2	1:A:69:GLY:H	1.61	0.64
1:A:59:MET:H	1:A:87:HIS:CD2	2.13	0.61
1:A:190:TYR:O	1:A:196:GLN:HB3	2.02	0.60
1:A:87:HIS:HE1	1:A:168:THR:OG1	1.87	0.58
1:A:180:ASN:HD22	1:A:220:ASP:HB3	1.68	0.57
1:A:64:LEU:HB2	1:A:72:VAL:CG2	2.34	0.57
1:A:289:LYS:HE2	1:A:289:LYS:HA	1.86	0.57
1:A:180:ASN:HD22	1:A:220:ASP:CB	2.19	0.56
1:A:59:MET:N	1:A:87:HIS:HD2	1.99	0.56
1:A:67:PRO:HD2	3:A:531:HOH:O	2.05	0.56
1:A:180:ASN:ND2	1:A:220:ASP:HB3	2.21	0.55
1:A:196:GLN:HB2	1:A:202:GLU:HB2	1.89	0.54
1:A:193:THR:HG23	1:A:194:PRO:HD2	1.90	0.53
1:A:47:ARG:NH2	1:A:98:GLN:HE22	2.05	0.53

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:217:GLU:OE1	1:A:256:THR:HB	2.09	0.53
1:A:11:PHE:HE2	1:A:332:ALA:HB1	1.74	0.53
1:A:250:ARG:HH12	1:A:251:ILE:CD1	2.22	0.53
1:A:310:LYS:HB2	3:A:466:HOH:O	2.10	0.51
1:A:31:GLU:HB2	1:A:35:LYS:HG3	1.92	0.51
1:A:280:ARG:HB2	1:A:323:GLU:OE1	2.13	0.48
1:A:190:TYR:HB3	1:A:193:THR:HG21	1.95	0.48
1:A:250:ARG:HH11	1:A:250:ARG:HB3	1.79	0.48
1:A:193:THR:HG23	1:A:195:ASN:H	1.79	0.47
1:A:250:ARG:HH12	1:A:251:ILE:HD11	1.79	0.46
1:A:16:ASN:HA	3:A:418:HOH:O	2.14	0.46
1:A:98:GLN:HA	1:A:98:GLN:HE21	1.82	0.45
1:A:247:GLU:O	1:A:251:ILE:HG12	2.17	0.45
1:A:323:GLU:HG3	3:A:492:HOH:O	2.17	0.44
1:A:108:GLY:O	1:A:181:TRP:HA	2.18	0.43
1:A:319:TYR:O	1:A:320:LEU:C	2.58	0.43
1:A:240:LEU:C	1:A:240:LEU:HD23	2.40	0.42
1:A:331:ILE:O	1:A:332:ALA:C	2.57	0.42
1:A:12:ASN:O	1:A:297:GLY:HA3	2.20	0.41
1:A:331:ILE:HG22	1:A:332:ALA:N	2.36	0.41
1:A:20:PHE:HA	1:A:23:HIS:CE1	2.56	0.40
1:A:264:PRO:HD2	1:A:297:GLY:O	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:236:GLU:OE1	1:A:261:LYS:NZ[4_455]	2.09	0.11

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	322/350 (92%)	307 (95%)	13 (4%)	2 (1%)	28	43

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	331	ILE
1	A	273	ARG

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	284/309 (92%)	265 (93%)	19 (7%)	19	30

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	30	GLU
1	A	35	LYS
1	A	75	ASN
1	A	80	GLN
1	A	85	THR
1	A	98	GLN
1	A	105	PRO
1	A	123	TRP
1	A	180	ASN
1	A	193	THR
1	A	218	SER
1	A	222	LEU
1	A	250	ARG
1	A	256	THR
1	A	259	ARG
1	A	265	TYR
1	A	273	ARG
1	A	278	LEU
1	A	310	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	75	ASN
1	A	80	GLN
1	A	87	HIS
1	A	98	GLN
1	A	121	GLN
1	A	162	GLN
1	A	180	ASN
1	A	254	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

Of 5 carbohydrates modelled in this entry, 1 is modelled with single atom - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	GCU	A	401	2	9,12,13	0.64	0	13,17,19	1.47	2 (15%)
2	NAG	A	402	2	14,14,15	0.59	0	15,19,21	1.25	1 (6%)
2	GCU	A	403	2	9,12,13	1.27	1 (11%)	13,17,19	1.94	1 (7%)
2	NAG	A	404	2	14,14,15	1.00	1 (7%)	15,19,21	2.39	4 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical

component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GCU	A	401	2	1/1/5/6	0/0/21/24	0/1/1/1
2	NAG	A	402	2	-	0/6/23/26	0/1/1/1
2	GCU	A	403	2	1/1/5/6	0/0/21/24	0/1/1/1
2	NAG	A	404	2	1/1/5/7	0/6/23/26	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	403	GCU	O5-C5	-2.78	1.40	1.43
2	A	404	NAG	O5-C1	-2.37	1.39	1.43

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	GCU	O5-C1-C2	-3.15	105.86	110.79
2	A	402	NAG	O3-C3-C4	-2.81	104.24	110.36
2	A	401	GCU	C1-C2-C3	2.57	112.92	109.65
2	A	404	NAG	C2-N2-C7	2.62	126.77	122.94
2	A	404	NAG	C1-C2-N2	3.45	116.37	110.49
2	A	404	NAG	C1-O5-C5	4.25	118.03	112.17
2	A	404	NAG	O5-C1-C2	6.32	120.27	111.47
2	A	403	GCU	C1-O5-C5	6.36	123.63	112.17

All (3) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	A	401	GCU	C1
2	A	404	NAG	C1
2	A	403	GCU	C1

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.6 Ligand geometry ⓘ

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	323/350 (92%)	-0.61	6 (1%) 67 66	15, 28, 60, 96	8 (2%)

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	144	TRP	2.9
1	A	194	PRO	2.7
1	A	147	GLN	2.5
1	A	195	ASN	2.4
1	A	68	ASN	2.1
1	A	17	VAL	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	NAG	A	404	14/15	0.92	0.24	3.74	22,38,54,58	0
2	NAG	A	402	14/15	0.93	0.18	3.06	25,33,41,45	0
2	GCU	A	403	12/13	0.97	0.16	1.85	25,29,35,37	0
2	GCU	A	405	1/13	0.87	0.47	-	51,51,51,51	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	GCU	A	401	12/13	0.91	0.24	-	43,50,54,54	0

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.