



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 11:50 am GMT

PDB ID : 3FCE  
Title : Crystal Structure of Bacillus cereus D-alanyl Carrier Protein Ligase DltA in Complex with ATP: Implications for Adenylation Mechanism  
Authors : Osman, K.T.; Du, L.; He, Y.; Luo, Y.  
Deposited on : 2008-11-21  
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	5047 (1.90-1.90)
Clashscore	112137	5731 (1.90-1.90)
Ramachandran outliers	110173	5669 (1.90-1.90)
Sidechain outliers	110143	5670 (1.90-1.90)
RSRZ outliers	101464	5100 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	512	<div> <div>4%</div> <div>72%</div> <div>22%</div> <div>• •</div> </div>

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4399 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called D-alanine--poly(phosphoribitol) ligase subunit 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	500	Total	C	N	O	S	0	0	0
			3943	2531	640	749	23			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	505	LEU	-	EXPRESSION TAG	UNP Q81G39
A	506	GLU	-	EXPRESSION TAG	UNP Q81G39
A	507	HIS	-	EXPRESSION TAG	UNP Q81G39
A	508	HIS	-	EXPRESSION TAG	UNP Q81G39
A	509	HIS	-	EXPRESSION TAG	UNP Q81G39
A	510	HIS	-	EXPRESSION TAG	UNP Q81G39
A	511	HIS	-	EXPRESSION TAG	UNP Q81G39
A	512	HIS	-	EXPRESSION TAG	UNP Q81G39

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Ca	0	0
			1	1		

- Molecule 3 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula: C<sub>10</sub>H<sub>16</sub>N<sub>5</sub>O<sub>13</sub>P<sub>3</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			31	10	5	13	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	424	Total	O	0	0
			424	424		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: D-alanine--poly(phosphoribitol) ligase subunit 1



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	61.40Å 87.40Å 57.40Å 90.00° 114.00° 90.00°	Depositor
Resolution (Å)	20.00 – 1.90 34.47 – 1.88	Depositor EDS
% Data completeness (in resolution range)	87.6 (20.00-1.90) 85.0 (34.47-1.88)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	0.06	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.19 (at 1.88Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.207 , 0.252 0.212 , 0.254	Depositor DCC
$R_{free}$ test set	1922 reflections (5.00%)	DCC
Wilson B-factor (Å <sup>2</sup> )	16.3	Xtriage
Anisotropy	0.177	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 49.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	4399	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.64% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, ATP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.32	0/4034	0.59	0/5469

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3943	0	3922	112	0
2	A	1	0	0	0	0
3	A	31	0	12	0	0
4	A	424	0	0	14	0
All	All	4399	0	3934	112	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

All (112) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:409:MET:SD	1:A:474:MET:HG2	2.05	0.97
1:A:279:ARG:HD2	1:A:313:LEU:HD13	1.53	0.91
1:A:415:GLU:HG2	1:A:429:ILE:HD13	1.64	0.79
1:A:227:LEU:HD22	1:A:231:LEU:HD21	1.66	0.78
1:A:27:ILE:HD12	1:A:66:ILE:HD11	1.66	0.78
1:A:471:PRO:HD2	1:A:474:MET:SD	2.26	0.76
1:A:477:ARG:HD2	1:A:478:LYS:HE3	1.69	0.74
1:A:111:THR:HG22	1:A:112:ASP:OD1	1.89	0.72
1:A:216:ILE:HG23	1:A:220:MET:SD	2.30	0.72
1:A:287:LYS:HE3	1:A:287:LYS:HA	1.75	0.69
1:A:47:TYR:HB3	1:A:50:ASP:HB3	1.76	0.68
1:A:228:PHE:CE2	1:A:252:GLU:HG3	2.29	0.68
1:A:276:GLU:HB2	4:A:1059:HOH:O	1.93	0.68
1:A:227:LEU:HD22	1:A:231:LEU:CD2	2.24	0.67
1:A:428:VAL:HG11	1:A:498:LEU:CD2	2.25	0.66
1:A:364:GLU:O	1:A:368:LYS:HG2	1.95	0.66
1:A:108:VAL:HG12	4:A:978:HOH:O	1.94	0.66
1:A:415:GLU:CG	1:A:429:ILE:HD13	2.25	0.66
1:A:264:LYS:HD3	1:A:265:THR:OG1	1.97	0.65
1:A:76:HIS:HE1	4:A:916:HOH:O	1.80	0.64
1:A:52:SER:O	1:A:76:HIS:HD2	1.80	0.64
1:A:497:LYS:O	1:A:501:GLU:HG2	1.99	0.62
1:A:403:LYS:NZ	1:A:403:LYS:HB2	2.12	0.62
1:A:334:LYS:HZ2	1:A:338:THR:HB	1.65	0.62
1:A:41:HIS:ND1	1:A:130:HIS:HD2	1.98	0.61
1:A:189:LEU:HB2	1:A:236:ILE:HD13	1.83	0.61
1:A:39:LEU:HA	1:A:126:ILE:HD13	1.83	0.61
1:A:136:ASN:HB3	1:A:139:HIS:CD2	2.37	0.60
1:A:402:ILE:CG1	1:A:409:MET:HG3	2.31	0.60
1:A:440:TYR:CE1	1:A:478:LYS:HD3	2.36	0.59
1:A:85:ILE:HG23	1:A:89:ARG:HE	1.66	0.59
1:A:403:LYS:HG3	4:A:1021:HOH:O	2.04	0.58
1:A:428:VAL:HG11	1:A:498:LEU:HD21	1.85	0.58
1:A:433:LYS:HE2	1:A:438:TYR:CZ	2.38	0.58
1:A:336:ASP:C	1:A:338:THR:H	2.06	0.57
1:A:466:LEU:HG	1:A:470:LEU:HD12	1.85	0.57
1:A:85:ILE:HD11	1:A:195:SER:OG	2.04	0.57
1:A:437:LYS:HG3	4:A:1021:HOH:O	2.05	0.57
1:A:336:ASP:OD1	1:A:337:GLY:N	2.38	0.57
1:A:2:LYS:HB2	1:A:5:GLU:HG3	1.87	0.56
1:A:8:GLU:CG	1:A:33:LYS:HE2	2.36	0.56
1:A:428:VAL:HG11	1:A:498:LEU:HD23	1.88	0.55

Continued on next page...



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:2:LYS:O	1:A:6:GLN:HG2	2.07	0.55
1:A:419:ARG:HD2	4:A:1071:HOH:O	2.06	0.54
1:A:428:VAL:CG1	1:A:444:VAL:HB	2.38	0.54
1:A:336:ASP:O	1:A:338:THR:N	2.40	0.53
1:A:14:THR:OG1	1:A:17:GLN:HB2	2.08	0.53
1:A:14:THR:HG21	4:A:963:HOH:O	2.09	0.53
1:A:428:VAL:HG13	1:A:444:VAL:HB	1.91	0.53
1:A:250:LEU:HD13	1:A:280:LYS:HE3	1.90	0.52
1:A:334:LYS:HZ1	1:A:338:THR:HG21	1.74	0.52
1:A:152:THR:HG23	1:A:152:THR:O	2.09	0.52
1:A:291:MET:HE2	1:A:307:HIS:HA	1.91	0.52
1:A:430:VAL:HG11	1:A:502:VAL:HG21	1.91	0.52
1:A:433:LYS:HE2	1:A:438:TYR:CE2	2.44	0.52
1:A:8:GLU:HG2	1:A:33:LYS:HE2	1.92	0.51
1:A:403:LYS:HZ2	1:A:403:LYS:HB2	1.74	0.51
1:A:290:ILE:O	1:A:291:MET:HE2	2.10	0.51
1:A:227:LEU:CD2	1:A:231:LEU:HD21	2.38	0.51
1:A:445:VAL:HG22	1:A:481:TYR:CD2	2.46	0.51
1:A:489:PRO:HG2	4:A:831:HOH:O	2.11	0.51
1:A:14:THR:O	1:A:14:THR:HG23	2.12	0.50
1:A:9:LYS:O	1:A:13:GLU:HG2	2.12	0.49
1:A:220:MET:HG2	1:A:227:LEU:HA	1.92	0.49
1:A:403:LYS:HA	1:A:407:TYR:O	2.12	0.49
1:A:499:LEU:O	1:A:502:VAL:HG22	2.13	0.49
1:A:498:LEU:O	1:A:502:VAL:HG13	2.12	0.49
1:A:289:THR:HG23	1:A:307:HIS:CE1	2.48	0.48
1:A:391:LEU:HD13	1:A:393:PHE:CZ	2.49	0.48
1:A:234:SER:HB2	1:A:236:ILE:HG13	1.96	0.48
1:A:27:ILE:CD1	1:A:66:ILE:HD11	2.42	0.47
1:A:416:HIS:HD2	4:A:860:HOH:O	1.97	0.47
1:A:57:TYR:HB3	1:A:104:SER:HB3	1.97	0.47
1:A:108:VAL:HG13	1:A:108:VAL:O	2.15	0.46
1:A:460:SER:O	1:A:464:LYS:HG3	2.15	0.46
1:A:89:ARG:HA	1:A:92:ARG:HH11	1.80	0.46
1:A:199:SER:O	1:A:203:ILE:HG13	2.16	0.46
1:A:492:LYS:HE2	4:A:930:HOH:O	2.15	0.46
1:A:402:ILE:HD11	1:A:414:ILE:CD1	2.46	0.45
1:A:95:GLU:HG2	1:A:114:PRO:HG3	1.97	0.45
1:A:88:ASP:O	1:A:92:ARG:HG3	2.16	0.45
1:A:245:PHE:O	1:A:248:MET:HG3	2.16	0.45
1:A:467:ASN:HA	1:A:475:ILE:HD11	1.97	0.45

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:478:LYS:HE3	4:A:1031:HOH:O	2.17	0.45
1:A:402:ILE:HG13	1:A:409:MET:CG	2.46	0.44
1:A:102:LEU:HD23	1:A:102:LEU:C	2.38	0.44
1:A:442:LEU:HD11	1:A:480:MET:HG3	1.99	0.44
1:A:97:SER:C	1:A:99:ALA:H	2.20	0.44
1:A:227:LEU:O	1:A:231:LEU:CD2	2.66	0.44
1:A:320:PRO:HB3	1:A:393:PHE:CE2	2.52	0.44
1:A:371:THR:CG2	1:A:378:ALA:HB3	2.48	0.43
1:A:411:LEU:HD23	1:A:411:LEU:C	2.38	0.43
1:A:402:ILE:HG22	1:A:438:TYR:CE2	2.53	0.43
1:A:57:TYR:CE2	1:A:108:VAL:HA	2.54	0.43
1:A:336:ASP:C	1:A:338:THR:N	2.72	0.43
1:A:129:THR:HG22	1:A:130:HIS:ND1	2.34	0.42
1:A:223:ARG:HA	1:A:224:PRO:HD2	1.88	0.42
1:A:248:MET:HG3	1:A:249:CYS:N	2.35	0.42
1:A:111:THR:O	1:A:112:ASP:C	2.58	0.42
1:A:41:HIS:CD2	1:A:133:ASN:HB3	2.54	0.42
1:A:501:GLU:HG2	1:A:501:GLU:H	1.63	0.42
1:A:334:LYS:NZ	1:A:338:THR:HG21	2.35	0.41
1:A:363:PRO:O	1:A:367:GLU:OE2	2.38	0.41
1:A:300:THR:C	1:A:301:VAL:HG23	2.41	0.41
1:A:431:PRO:HG2	4:A:846:HOH:O	2.21	0.41
1:A:291:MET:CE	1:A:307:HIS:HA	2.49	0.41
1:A:380:LYS:HG3	4:A:1092:HOH:O	2.20	0.41
1:A:179:ASP:HB3	4:A:1203:HOH:O	2.20	0.40
1:A:39:LEU:HD22	1:A:43:ILE:HG13	2.03	0.40
1:A:417:HIS:HD2	1:A:469:ARG:HH21	1.69	0.40
1:A:409:MET:HB2	1:A:474:MET:CE	2.51	0.40
1:A:160:LYS:HE2	1:A:359:TYR:CZ	2.57	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	496/512 (97%)	474 (96%)	17 (3%)	5 (1%)	18 7

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	153	SER
1	A	112	ASP
1	A	335	GLU
1	A	337	GLY
1	A	301	VAL

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	434/446 (97%)	417 (96%)	17 (4%)	37 26

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	8	GLU
1	A	39	LEU
1	A	191	GLN
1	A	227	LEU
1	A	248	MET
1	A	276	GLU
1	A	279	ARG
1	A	287	LYS
1	A	291	MET
1	A	307	HIS
1	A	351	VAL
1	A	364	GLU
1	A	391	LEU
1	A	403	LYS
1	A	445	VAL
1	A	478	LYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	498	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	76	HIS
1	A	130	HIS
1	A	133	ASN
1	A	139	HIS
1	A	146	ASN
1	A	190	ASN
1	A	191	GLN
1	A	237	GLN
1	A	307	HIS
1	A	315	GLN
1	A	389	ASN
1	A	395	ASN
1	A	401	GLN
1	A	416	HIS
1	A	417	HIS
1	A	467	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	ATP	A	711	2	27,33,33	2.43	9 (33%)	25,52,52	1.75	2 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ATP	A	711	2	-	0/18/38/38	0/3/3/3

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	711	ATP	C5'-C4'	-5.65	1.33	1.51
3	A	711	ATP	PG-O3G	-2.32	1.45	1.54
3	A	711	ATP	C8-N7	-2.17	1.30	1.34
3	A	711	ATP	O5'-C5'	2.20	1.53	1.44
3	A	711	ATP	C2-N3	2.29	1.36	1.32
3	A	711	ATP	O4'-C1'	2.58	1.44	1.41
3	A	711	ATP	C2-N1	4.38	1.42	1.33
3	A	711	ATP	C4-N3	5.00	1.42	1.35
3	A	711	ATP	O2'-C2'	5.06	1.54	1.43

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	711	ATP	N3-C2-N1	-7.87	122.00	128.86
3	A	711	ATP	O3'-C3'-C2'	2.38	119.45	111.83

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	500/512 (97%)	0.12	23 (4%)	33 37	8, 20, 41, 55	0

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	365	LEU	4.5
1	A	111	THR	4.3
1	A	364	GLU	4.2
1	A	112	ASP	3.9
1	A	336	ASP	3.8
1	A	408	ARG	3.7
1	A	154	GLY	3.6
1	A	367	GLU	3.4
1	A	402	ILE	3.4
1	A	92	ARG	3.3
1	A	503	THR	3.3
1	A	109	THR	3.2
1	A	335	GLU	3.2
1	A	434	LYS	3.2
1	A	337	GLY	3.2
1	A	407	TYR	3.1
1	A	108	VAL	3.1
1	A	363	PRO	2.9
1	A	368	LYS	2.8
1	A	437	LYS	2.6
1	A	152	THR	2.5
1	A	153	SER	2.3
1	A	105	ALA	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	ATP	A	711	31/31	0.94	0.13	-0.09	11,14,40,41	0
2	CA	A	710	1/1	0.98	0.05	-	33,33,33,33	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.