



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 14, 2017 – 11:07 am GMT

PDB ID : 1FEC  
Title : UNLIGANDED CRITHIDIA FASCICULATA TRYPANOTHIONE REDUC-  
TASE AT 1.7 ANGSTROM RESOLUTION  
Authors : Strickland, C.; Karplus, P.  
Deposited on : 1995-07-12  
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	3453 (1.70-1.70)
Clashscore	112137	3876 (1.70-1.70)
Ramachandran outliers	110173	3815 (1.70-1.70)
Sidechain outliers	110143	3815 (1.70-1.70)
RSRZ outliers	101464	3491 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	490	<div> <div>%</div> <div> <div></div> <div>82%</div> <div>16%</div> <div>..</div> </div> </div>
1	B	490	<div> <div>3%</div> <div> <div></div> <div>88%</div> <div>11%</div> <div>..</div> </div> </div>

## 2 Entry composition i

There are 3 unique types of molecules in this entry. The entry contains 8547 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called TRYPANOTHIONE REDUCTASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	485	Total	C	N	O	S	0	11	0
			3743	2353	648	718	24			
1	B	485	Total	C	N	O	S	0	4	0
			3710	2337	642	709	22			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	296	GLU	ASP	CONFLICT	UNP P39040
A	478	GLU	GLN	CONFLICT	UNP P39040
B	296	GLU	ASP	CONFLICT	UNP P39040
B	478	GLU	GLN	CONFLICT	UNP P39040

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: C<sub>27</sub>H<sub>33</sub>N<sub>9</sub>O<sub>15</sub>P<sub>2</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	495	Total	O	0	0
			495	495		
3	B	493	Total	O	0	0
			493	493		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: TRYPANOTHIONE REDUCTASE



#### • Molecule 1: TRYPANOTHIONE REDUCTASE



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	57.80Å 160.70Å 60.00Å 90.00° 99.40° 90.00°	Depositor
Resolution (Å)	8.00 – 1.70 46.50 – 1.60	Depositor EDS
% Data completeness (in resolution range)	(Not available) (8.00-1.70) 84.8 (46.50-1.60)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.46 (at 1.60Å)	Xtriage
Refinement program	X-PLOR	Depositor
R, $R_{free}$	0.204 , 0.256 0.189 , 0.241	Depositor DCC
$R_{free}$ test set	5211 reflections (4.98%)	DCC
Wilson B-factor (Å <sup>2</sup> )	14.3	Xtriage
Anisotropy	0.424	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 82.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.027 for l,-k,h	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	8547	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.49% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.76	0/3846	0.83	0/5206
1	B	0.76	0/3799	0.82	2/5147 (0.0%)
All	All	0.76	0/7645	0.83	2/10353 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	326	ASP	CB-CG-OD2	-5.95	112.94	118.30
1	B	417	ASN	N-CA-C	-5.31	96.65	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3743	0	3671	51	0
1	B	3710	0	3655	37	0
2	A	53	0	31	0	0
2	B	53	0	31	1	0
3	A	495	0	0	7	2
3	B	493	0	0	10	2
All	All	8547	0	7388	88	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (88) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:51[B]:CYS:HB2	1:B:56:CYS:SG	2.11	0.89
1:A:51[B]:CYS:SG	3:B:698:HOH:O	2.26	0.87
1:A:196:GLY:O	1:A:200:ILE:HD13	1.81	0.80
3:A:678:HOH:O	1:B:51[B]:CYS:SG	2.45	0.75
1:B:466:GLU:HG2	3:B:957:HOH:O	1.91	0.71
1:A:3:ALA:O	1:A:29:ARG:HD3	1.91	0.70
1:B:423[B]:VAL:HG23	1:B:444:LEU:HD22	1.73	0.69
1:A:138:GLU:HG2	1:A:147:LEU:HD21	1.75	0.67
1:B:423[B]:VAL:HG22	1:B:448:ALA:O	1.96	0.65
1:B:20:TRP:O	1:B:24:SER:HB3	1.96	0.65
1:A:249:ARG:HB3	1:A:252:GLU:HG3	1.79	0.64
1:A:253:ASN:O	1:A:270:GLU:HG3	1.97	0.63
1:A:315:LYS:HD2	1:A:319:ASP:HA	1.83	0.60
1:B:445:LYS:HG2	3:B:980:HOH:O	2.01	0.60
1:A:129:GLN:HB3	1:A:133:THR:HB	1.83	0.60
1:B:43:HIS:HD2	3:B:703:HOH:O	1.85	0.60
1:B:220:TYR:CE2	1:B:222:GLY:HA3	2.37	0.59
1:B:36:GLN:OE1	1:B:42:PRO:HD2	2.01	0.59
1:A:51[B]:CYS:HB2	1:A:56:CYS:SG	2.43	0.58
1:A:51[B]:CYS:HB2	1:A:334:THR:HG1	1.69	0.58
1:A:51[B]:CYS:SG	1:A:56:CYS:SG	3.02	0.57
1:B:294:GLN:O	1:B:297:LYS:HB2	2.04	0.57
1:B:384:LYS:HE2	3:B:925:HOH:O	2.04	0.57
1:B:486:ILE:O	1:B:486:ILE:HG13	2.05	0.56
1:B:200:ILE:CD1	1:B:218:LEU:HD21	2.36	0.56
1:B:200:ILE:HD12	1:B:218:LEU:HD21	1.88	0.55
1:A:394:SER:HB3	1:A:411:MET:HG2	1.89	0.55
1:A:391:TYR:CZ	1:A:473:PRO:HG3	2.42	0.55
1:B:131:ASN:HB3	3:B:882:HOH:O	2.06	0.55
1:A:389:ALA:HB3	1:A:416:THR:OG1	2.07	0.55
1:B:7:VAL:HG22	1:B:31:ALA:HB3	1.88	0.54
1:A:38:HIS:CD2	3:A:664:HOH:O	2.61	0.53
1:A:51[B]:CYS:CB	1:A:56:CYS:SG	2.98	0.52
1:A:304:LYS:HD2	1:A:304:LYS:O	2.10	0.52
1:A:387:GLN:HG2	1:A:478:GLU:OE2	2.10	0.51
1:A:28:LYS:HE3	1:A:349:PHE:HD1	1.75	0.51
1:A:395:PHE:CE1	1:A:466:GLU:HG2	2.46	0.51

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:392:GLU:HB2	3:B:746:HOH:O	2.11	0.50
1:B:423[B]:VAL:CG2	1:B:444:LEU:HD22	2.41	0.49
1:B:27:LYS:HD3	1:B:27:LYS:N	2.27	0.49
1:A:88:ARG:HD2	3:A:694:HOH:O	2.13	0.48
1:B:362:ALA:HB2	1:B:441:ALA:HB2	1.95	0.48
1:B:43:HIS:HE1	3:B:791:HOH:O	1.96	0.48
1:A:395:PHE:HE1	1:A:466:GLU:HG2	1.78	0.47
1:A:346:ASP:HA	1:A:350:ALA:HB3	1.96	0.47
1:A:28:LYS:HE3	1:A:349:PHE:CD1	2.50	0.47
1:A:51[B]:CYS:HB3	3:A:683:HOH:O	2.13	0.47
1:B:366:PHE:HE2	3:B:798:HOH:O	1.97	0.47
1:A:59:LYS:HD2	1:A:60:LYS:N	2.30	0.46
1:B:417:ASN:ND2	1:B:419:ALA:H	2.14	0.46
1:A:315:LYS:HG2	1:A:316:THR:O	2.15	0.46
1:B:88:ARG:HH21	1:B:212:ARG:NH1	2.14	0.46
1:A:43:HIS:O	1:A:44:TYR:HB2	2.17	0.45
1:B:41:PRO:HG3	1:B:44:TYR:CE1	2.51	0.45
1:A:388:VAL:HB	1:A:477:TYR:HB2	1.99	0.45
1:B:431:ASP:O	1:B:432:SER:HB2	2.17	0.45
1:A:460:HIS:HA	1:A:461:PRO:HA	1.80	0.45
1:A:354:ARG:HD2	3:A:911:HOH:O	2.17	0.44
1:B:33:ILE:HG22	2:B:499:FAD:H2A	1.99	0.44
1:A:350:ALA:O	1:A:352:LYS:HG3	2.16	0.44
1:A:74:GLU:HB3	1:A:403:SER:HB2	2.00	0.44
1:A:200:ILE:CD1	1:A:200:ILE:N	2.81	0.44
1:A:89:PRO:HD3	3:A:855:HOH:O	2.18	0.44
1:A:223:ASP:HA	1:A:251:HIS:NE2	2.33	0.43
1:A:191:LEU:HD23	1:A:280:VAL:HG13	1.99	0.43
1:A:58:PRO:HB3	1:A:98:LYS:HD3	2.00	0.43
1:A:394:SER:HA	1:A:410:PHE:O	2.18	0.43
1:B:33:ILE:HA	1:B:122:HIS:O	2.19	0.43
1:A:247:ASN:HA	3:A:725:HOH:O	2.17	0.43
1:A:362:ALA:HB2	1:A:441:ALA:HB2	2.01	0.42
1:A:156:LEU:HD11	1:A:324:ILE:HG23	2.00	0.42
1:A:220:TYR:O	1:A:251:HIS:N	2.52	0.42
1:A:223:ASP:OD1	1:A:223:ASP:N	2.53	0.42
1:B:389:ALA:HB3	1:B:416:THR:OG1	2.19	0.42
1:B:476:PHE:CG	1:B:486:ILE:HG12	2.54	0.41
1:B:221:ARG:HG2	3:B:582:HOH:O	2.20	0.41
1:A:112:MET:HE1	1:A:113:PHE:CE1	2.56	0.41
1:A:125:PHE:CD1	1:A:139:SER:HA	2.56	0.41

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:160:GLY:HA2	1:B:326:ASP:HB2	2.02	0.41
1:A:159:THR:HG21	1:A:293:LEU:HD21	2.03	0.41
1:A:166:LEU:N	1:A:171:ASP:OD2	2.54	0.41
1:B:220:TYR:CE2	1:B:222:GLY:CA	3.04	0.41
1:B:386:ASP:OD1	1:B:479:LYS:HG2	2.21	0.40
1:A:169:GLU:OE1	1:A:256:LYS:HE2	2.20	0.40
1:A:478:GLU:O	1:A:479:LYS:C	2.59	0.40
1:B:383:LYS:HD3	1:B:383:LYS:HA	1.81	0.40
1:B:391:TYR:CZ	1:B:473:PRO:HG3	2.56	0.40
1:A:68:TYR:HA	1:A:71:THR:OG1	2.21	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:632:HOH:O	3:B:623:HOH:O[1_455]	2.09	0.11
3:A:553:HOH:O	3:B:697:HOH:O[1_556]	2.19	0.01

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	494/490 (101%)	471 (95%)	23 (5%)	0	100	100
1	B	487/490 (99%)	473 (97%)	13 (3%)	1 (0%)	51	31
All	All	981/980 (100%)	944 (96%)	36 (4%)	1 (0%)	55	34

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	54	VAL

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	396/393 (101%)	384 (97%)	12 (3%)	46	25
1	B	392/393 (100%)	383 (98%)	9 (2%)	56	36
All	All	788/786 (100%)	767 (97%)	21 (3%)	50	30

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	56	CYS
1	A	92	LYS
1	A	119	LEU
1	A	184	ASP
1	A	200	ILE
1	A	223	ASP
1	A	247	ASN
1	A	287	VAL
1	A	304	LYS
1	A	369	PRO
1	A	384	LYS
1	A	481	LYS
1	B	56	CYS
1	B	59	LYS
1	B	98	LYS
1	B	133	THR
1	B	221	ARG
1	B	256	LYS
1	B	354	ARG
1	B	417	ASN
1	B	484	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	26	HIS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	207	ASN
1	A	247	ASN
1	B	38	HIS
1	B	43	HIS
1	B	123	GLN
1	B	207	ASN
1	B	251	HIS
1	B	417	ASN
1	B	418	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	FAD	A	499	-	51,58,58	1.84	10 (19%)	54,89,89	1.49	3 (5%)
2	FAD	B	499	-	51,58,58	1.92	15 (29%)	54,89,89	1.40	3 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	A	499	-	-	0/28/50/50	0/6/6/6
2	FAD	B	499	-	-	0/28/50/50	0/6/6/6

All (25) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	499	FAD	C8A-N7A	-3.50	1.28	1.34
2	B	499	FAD	PA-O2A	-3.41	1.38	1.55
2	B	499	FAD	C4A-N3A	-3.35	1.30	1.35
2	B	499	FAD	C8M-C8	-3.04	1.45	1.51
2	A	499	FAD	C2'-C3'	-2.82	1.47	1.53
2	B	499	FAD	C2-N3	2.05	1.42	1.38
2	A	499	FAD	C9-C9A	2.07	1.45	1.40
2	B	499	FAD	C1'-N10	2.09	1.50	1.48
2	A	499	FAD	C9-C8	2.16	1.43	1.37
2	A	499	FAD	C4'-C3'	2.28	1.58	1.53
2	B	499	FAD	C7M-C7	2.41	1.55	1.51
2	B	499	FAD	C9A-N10	2.42	1.41	1.38
2	B	499	FAD	O4'-C4'	2.43	1.48	1.43
2	B	499	FAD	O3'-C3'	2.58	1.48	1.43
2	B	499	FAD	C9-C9A	2.59	1.46	1.40
2	B	499	FAD	C10-N1	2.90	1.37	1.33
2	A	499	FAD	C5X-N5	2.91	1.39	1.35
2	A	499	FAD	C8-C7	2.93	1.48	1.41
2	A	499	FAD	C2B-C3B	3.09	1.61	1.53
2	A	499	FAD	C4-N3	3.56	1.39	1.33
2	B	499	FAD	C2B-C3B	3.82	1.63	1.53
2	B	499	FAD	O2B-C2B	3.87	1.51	1.43
2	B	499	FAD	C5X-N5	3.99	1.41	1.35
2	A	499	FAD	C10-N1	4.56	1.39	1.33
2	A	499	FAD	O4B-C1B	5.88	1.49	1.41

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	499	FAD	C4X-C4-N3	-3.88	117.96	123.48
2	B	499	FAD	C4X-C4-N3	-3.57	118.41	123.48
2	B	499	FAD	C4X-C10-N10	-2.07	119.08	120.52
2	A	499	FAD	C1'-N10-C9A	3.23	121.31	118.35

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	499	FAD	C4-N3-C2	7.23	121.48	115.16
2	A	499	FAD	C4-N3-C2	7.76	121.94	115.16

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	499	FAD	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	485/490 (98%)	-0.31	4 (0%) 86 88	7, 19, 48, 79	0
1	B	485/490 (98%)	-0.32	14 (2%) 52 58	7, 17, 38, 68	0
All	All	970/980 (98%)	-0.31	18 (1%) 67 72	7, 18, 46, 79	0

All (18) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	486	ILE	6.3
1	B	117	GLU	3.5
1	B	119	LEU	3.1
1	A	212[A]	ARG	3.0
1	B	479	LYS	2.9
1	B	114	ALA	2.8
1	A	485	LYS	2.4
1	B	111	GLY	2.4
1	B	25	LEU	2.4
1	B	115	ASP	2.4
1	A	166	LEU	2.3
1	B	23	ALA	2.3
1	B	22	ALA	2.3
1	B	27	LYS	2.3
1	B	3	ALA	2.2
1	A	168	ILE	2.2
1	B	350	ALA	2.1
1	B	349	PHE	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	FAD	B	499	53/53	0.98	0.08	0.13	7,12,17,20	0
2	FAD	A	499	53/53	0.98	0.07	-0.35	6,14,18,19	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.