



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 09:05 am GMT

PDB ID : 4FGK  
Title : Oxidized quinone reductase 2 in complex with chloroquine  
Authors : Leung, K.K.; Shilton, B.H.  
Deposited on : 2012-06-04  
Resolution : 1.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	1307 (1.40-1.40)
Clashscore	112137	1411 (1.40-1.40)
Ramachandran outliers	110173	1373 (1.40-1.40)
Sidechain outliers	110143	1372 (1.40-1.40)
RSRZ outliers	101464	1315 (1.40-1.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	233	<div> <div>8%</div> <div> <div></div> <div>92%</div> <div>6% •</div> </div> </div>
1	B	233	<div> <div>3%</div> <div> <div></div> <div>96%</div> <div>• •</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	OTX	A	302[A]	-	-	-	X
3	OTX	A	302[B]	-	-	-	X
3	OTX	A	304	-	-	-	X

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 8091 atoms, of which 3622 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Ribosyldihyronicotinamide dehydrogenase [quinone].

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	230	Total	C	H	N	O	S	0	14	0
			3677	1239	1750	320	359	9			
1	B	230	Total	C	H	N	O	S	0	5	0
			3641	1194	1784	310	344	9			

There are 6 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	EXPRESSION TAG	UNP P16083
A	-1	ALA	-	EXPRESSION TAG	UNP P16083
A	46	PHE	LEU	SEE REMARK 999	UNP P16083
B	-2	GLY	-	EXPRESSION TAG	UNP P16083
B	-1	ALA	-	EXPRESSION TAG	UNP P16083
B	46	PHE	LEU	SEE REMARK 999	UNP P16083

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula:  $C_{27}H_{33}N_9O_{15}P_2$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	H	N	O	P	
			84	27	31	9	15	2	0
2	B	1	Total	C	H	N	O	P	
			84	27	31	9	15	2	0

- Molecule 3 is (4S)-N 4 -(7-CHLOROQUINOLIN-4-YL)-N 1 ,N 1 -DIETHYLPENTANE-1,4 -DIAMINE (three-letter code: OTX) (formula: C<sub>18</sub>H<sub>26</sub>ClN<sub>3</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	Cl	N			
			44	36	2	6		0	1
3	A	1	Total	C	Cl	H	N		
			48	18	1	26	3	0	0

- Molecule 4 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Zn		
			1	1	0	0
4	A	1	Total	Zn		
			1	1	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	221	Total	O		
			221	221	0	0

Continued on next page...


*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	290	Total 290	O 290	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Ribosyldihydronicotinamide dehydrogenase [quinone]



- Molecule 1: Ribosyldihydronicotinamide dehydrogenase [quinone]



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	56.37Å 83.11Å 106.58Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	32.67 – 1.40 32.67 – 1.40	Depositor EDS
% Data completeness (in resolution range)	99.5 (32.67-1.40) 99.5 (32.67-1.40)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	0.07	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.08 (at 1.40Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.7.3_928)	Depositor
R, $R_{free}$	0.133 , 0.174 0.132 , 0.172	Depositor DCC
$R_{free}$ test set	1998 reflections (2.03%)	DCC
Wilson B-factor (Å <sup>2</sup> )	13.0	Xtriage
Anisotropy	0.586	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.45 , 52.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.98	EDS
Total number of atoms	8091	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.03% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, FAD, OTX

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.37	0/1980	0.59	0/2684
1	B	0.36	0/1907	0.57	0/2588
All	All	0.37	0/3887	0.58	0/5272

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1927	1750	1870	25	1
1	B	1857	1784	1811	6	0
2	A	53	31	30	0	0
2	B	53	31	31	1	0
3	A	66	26	78	11	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	221	0	0	4	1
5	B	290	0	0	4	0
All	All	4469	3622	3820	39	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 5.

All (39) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:127[B]:ASP:OD2	5:A:621:HOH:O	1.95	0.85
3:A:304:OTX:CL	5:B:689:HOH:O	2.35	0.80
1:A:72:HIS:CG	1:A:122[B]:GLN:HG3	2.23	0.73
1:A:131[B]:PHE:CB	1:A:178:PHE:CZ	2.80	0.65
1:B:27:ASP:OD1	5:B:554:HOH:O	2.15	0.63
1:A:127[B]:ASP:N	1:A:130[B]:GLY:O	2.25	0.59
3:A:302[B]:OTX:H181	1:B:193:GLU:OE1	2.02	0.58
1:A:169:TRP:HB3	1:A:170:PRO:HD3	1.85	0.58
1:A:131[B]:PHE:HB2	1:A:178:PHE:CZ	2.39	0.57
1:A:122[B]:GLN:HB3	1:A:126:PHE:CZ	2.39	0.57
1:A:127[B]:ASP:O	1:A:131[B]:PHE:HD1	1.89	0.55
1:A:131[B]:PHE:CD2	1:A:178:PHE:CE1	2.96	0.53
3:A:304:OTX:H61	5:B:689:HOH:O	2.09	0.53
1:A:131[B]:PHE:CG	1:A:178:PHE:CZ	2.96	0.53
1:A:72:HIS:HB2	1:A:122[B]:GLN:CG	2.38	0.53
1:A:127[B]:ASP:HB3	1:A:129[B]:PRO:O	2.10	0.52
1:A:24:VAL:HG11	1:A:208:SER:HB2	1.91	0.52
1:A:131[B]:PHE:HB2	1:A:178:PHE:CE1	2.45	0.52
1:A:132[B]:TYR:CD1	1:A:178:PHE:HA	2.47	0.50
1:A:131[A]:PHE:CE1	3:A:302[A]:OTX:H162	2.47	0.49
1:A:122[B]:GLN:HG2	5:A:473:HOH:O	2.13	0.47
3:A:302[B]:OTX:H182	3:A:302[B]:OTX:H122	1.47	0.46
1:B:169:TRP:HB3	1:B:170:PRO:HD3	1.97	0.45
1:A:72:HIS:CB	1:A:122[B]:GLN:HG3	2.47	0.44
1:A:129[B]:PRO:HA	1:A:131[B]:PHE:CE1	2.52	0.44
1:A:129[B]:PRO:HA	1:A:131[B]:PHE:HE1	1.82	0.44
3:A:302[B]:OTX:C11	3:A:302[B]:OTX:H142	2.49	0.42
1:A:133[B]:ASP:OD1	5:A:584:HOH:O	2.21	0.42
1:A:24:VAL:CG1	1:A:208:SER:HB2	2.49	0.42
1:A:72:HIS:HB2	1:A:122[B]:GLN:HG3	2.01	0.42
3:A:302[A]:OTX:H81	1:B:105:TRP:CZ3	2.55	0.41
3:A:304:OTX:H121	3:A:304:OTX:H162	1.85	0.41
1:A:132[B]:TYR:CZ	1:A:178:PHE:HD1	2.38	0.41
3:A:302[B]:OTX:H181	1:B:193:GLU:CD	2.41	0.41
3:A:302[B]:OTX:H153	3:A:302[B]:OTX:H162	1.68	0.41
3:A:302[B]:OTX:H172	3:A:302[B]:OTX:H132	1.79	0.41
5:A:440:HOH:O	2:B:302:FAD:HM72	2.19	0.41
1:A:132[B]:TYR:CE1	1:A:178:PHE:HD1	2.39	0.41

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:127:ASP:OD1	5:B:451:HOH:O	2.22	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:218:GLU:OE1	5:A:621:HOH:O[4_445]	2.01	0.19

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	241/233 (103%)	232 (96%)	9 (4%)	0	100	100
1	B	233/233 (100%)	224 (96%)	9 (4%)	0	100	100
All	All	474/466 (102%)	456 (96%)	18 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	204/195 (105%)	204 (100%)	0	100	100
1	B	198/195 (102%)	197 (100%)	1 (0%)	91	77

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
All	All	402/390 (103%)	401 (100%)	1 (0%)	94	84

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	230	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 7 ligands modelled in this entry, 2 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	FAD	A	301	-	50,58,58	1.89	12 (24%)	53,89,89	1.96	8 (15%)
3	0TX	A	302[A]	-	22,23,23	0.99	2 (9%)	28,30,30	1.37	5 (17%)
3	0TX	A	302[B]	-	22,23,23	1.08	2 (9%)	28,30,30	1.28	4 (14%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	0TX	A	304	-	22,23,23	1.11	2 (9%)	28,30,30	1.31	4 (14%)
2	FAD	B	302	-	50,58,58	1.92	12 (24%)	53,89,89	1.90	9 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	A	301	-	-	0/28/50/50	0/6/6/6
3	0TX	A	302[A]	-	-	0/14/14/14	0/2/2/2
3	0TX	A	302[B]	-	-	0/14/14/14	0/2/2/2
3	0TX	A	304	-	-	0/14/14/14	0/2/2/2
2	FAD	B	302	-	-	0/28/50/50	0/6/6/6

All (30) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	301	FAD	C2B-C3B	-3.85	1.43	1.53
2	B	302	FAD	C2B-C3B	-3.67	1.43	1.53
2	A	301	FAD	O4'-C4'	-2.89	1.37	1.43
2	B	302	FAD	O4'-C4'	-2.80	1.37	1.43
3	A	302[B]	0TX	C9-N1	-2.74	1.32	1.37
2	A	301	FAD	O3B-C3B	-2.68	1.36	1.43
2	A	301	FAD	C2B-C1B	-2.66	1.49	1.53
2	B	302	FAD	O3B-C3B	-2.53	1.37	1.43
3	A	304	0TX	C9-N1	-2.51	1.33	1.37
3	A	302[A]	0TX	C9-N1	-2.45	1.33	1.37
2	B	302	FAD	C9A-C5X	-2.36	1.37	1.42
2	A	301	FAD	C9A-C5X	-2.25	1.37	1.42
2	A	301	FAD	C9A-N10	-2.19	1.35	1.38
2	B	302	FAD	C9A-N10	-2.15	1.35	1.38
2	B	302	FAD	C6A-N6A	2.02	1.42	1.34
2	A	301	FAD	C6A-N6A	2.10	1.42	1.34
2	B	302	FAD	C1'-N10	2.43	1.50	1.48
2	A	301	FAD	C2-N1	2.55	1.43	1.38
2	B	302	FAD	C2-N1	2.57	1.43	1.38
2	A	301	FAD	C4X-N5	2.65	1.37	1.33
2	B	302	FAD	C4X-N5	2.84	1.37	1.33
3	A	302[A]	0TX	C3-N2	2.95	1.45	1.37
3	A	302[B]	0TX	C3-N2	3.09	1.46	1.37
2	A	301	FAD	C1'-N10	3.26	1.51	1.48

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	304	OTX	C3-N2	3.42	1.46	1.37
2	B	302	FAD	C4-N3	3.47	1.41	1.36
2	A	301	FAD	C4-C4X	3.69	1.47	1.41
2	B	302	FAD	C4-C4X	3.76	1.47	1.41
2	A	301	FAD	C10-N1	7.64	1.44	1.33
2	B	302	FAD	C10-N1	7.91	1.44	1.33

All (30) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	301	FAD	N3A-C2A-N1A	-9.49	120.59	128.86
2	B	302	FAD	N3A-C2A-N1A	-8.16	121.75	128.86
2	B	302	FAD	C4B-O4B-C1B	-4.21	105.29	109.77
2	A	301	FAD	C4B-O4B-C1B	-3.73	105.80	109.77
2	A	301	FAD	C1B-N9A-C4A	-3.49	120.60	126.64
2	B	302	FAD	C1B-N9A-C4A	-2.93	121.57	126.64
3	A	302[A]	OTX	C4-C9-N1	-2.76	119.94	122.89
2	A	301	FAD	C4X-C10-N10	-2.74	118.62	120.52
3	A	304	OTX	C4-C9-N1	-2.60	120.10	122.89
2	B	302	FAD	C4X-C10-N10	-2.37	118.87	120.52
2	A	301	FAD	N3-C2-N1	-2.36	117.34	122.16
3	A	302[A]	OTX	C11-C12-C13	-2.33	106.47	113.51
3	A	302[B]	OTX	C12-C11-C10	-2.32	108.88	113.78
2	B	302	FAD	N3-C2-N1	-2.30	117.47	122.16
3	A	302[B]	OTX	C4-C9-N1	-2.17	120.56	122.89
3	A	302[A]	OTX	C2-C1-N1	-2.08	121.35	124.58
3	A	302[B]	OTX	C8-C9-N1	2.29	120.42	117.97
2	A	301	FAD	C1'-N10-C9A	2.30	120.45	118.35
3	A	304	OTX	C3-C4-C9	2.55	119.48	118.04
3	A	302[B]	OTX	C1-N1-C9	2.68	121.14	116.89
3	A	304	OTX	C8-C9-N1	2.69	120.85	117.97
3	A	304	OTX	C1-N1-C9	2.80	121.34	116.89
2	A	301	FAD	C5X-C9A-N10	2.87	119.79	117.66
2	B	302	FAD	C4X-N5-C5X	2.99	119.92	116.76
2	B	302	FAD	C1'-N10-C9A	3.01	121.11	118.35
2	B	302	FAD	C5X-C9A-N10	3.12	119.98	117.66
3	A	302[A]	OTX	C8-C9-N1	3.36	121.56	117.97
3	A	302[A]	OTX	C1-N1-C9	3.53	122.49	116.89
2	A	301	FAD	C2-N1-C10	5.14	119.85	114.90
2	B	302	FAD	C2-N1-C10	5.22	119.93	114.90

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	302[A]	0TX	2	0
3	A	302[B]	0TX	6	0
3	A	304	0TX	3	0
2	B	302	FAD	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	230/233 (98%)	-0.02	18 (7%) 14 13	6, 19, 36, 51	0
1	B	230/233 (98%)	-0.38	8 (3%) 44 44	7, 13, 33, 49	0
All	All	460/466 (98%)	-0.20	26 (5%) 24 23	6, 15, 35, 51	0

All (26) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	64	VAL	4.8
1	A	198	GLU	4.7
1	B	1	ALA	4.2
1	A	63	GLU	3.7
1	A	197	GLU	3.6
1	B	198	GLU	3.4
1	A	57	GLY	3.2
1	B	197	GLU	3.2
1	A	196	SER	3.1
1	A	1[A]	ALA	3.0
1	A	201	LYS	2.9
1	A	62	PRO	2.8
1	A	230	GLN	2.8
1	B	129	PRO	2.8
1	B	2	GLY	2.7
1	A	59	LEU	2.7
1	B	159[A]	GLY	2.6
1	A	195	ALA	2.5
1	A	158	THR	2.5
1	A	58	THR	2.3
1	A	128[A]	ILE	2.2
1	A	61	ASN	2.2
1	A	135[A]	GLY	2.1
1	B	196	SER	2.1

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	158[A]	THR	2.1
1	A	194	ILE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	OTX	A	302[A]	22/22	0.91	0.21	3.24	8,14,25,27	22
3	OTX	A	302[B]	22/22	0.91	0.21	2.61	9,22,33,34	22
3	OTX	A	304	22/22	0.93	0.18	2.53	26,47,57,57	26
2	FAD	A	301	53/53	0.97	0.08	-0.16	8,15,31,38	0
2	FAD	B	302	53/53	0.97	0.07	-0.29	10,15,31,39	0
4	ZN	A	303	1/1	0.99	0.05	-1.26	22,22,22,22	0
4	ZN	B	301	1/1	1.00	0.03	-2.50	11,11,11,11	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.