



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 15, 2017 – 02:02 am GMT

PDB ID : 5FGJ
Title : Structure of tetrameric rat phenylalanine hydroxylase, residues 1-453
Authors : Taylor, A.B.; Roberts, K.M.; Fitzpatrick, P.F.
Deposited on : 2015-12-20
Resolution : 3.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	1026 (3.74-3.46)
Clashscore	112137	1036 (3.70-3.50)
Ramachandran outliers	110173	1030 (3.72-3.48)
Sidechain outliers	110143	1030 (3.72-3.48)
RSRZ outliers	101464	1051 (3.74-3.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	453	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 13%, green 81%, grey 6%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> % 81% 13% 6% </div> </div>
1	B	453	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, yellow 16%, green 77%, grey 6%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> % 77% 16% 6% </div> </div>
1	C	453	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 14%, green 79%, grey 6%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> % 79% 14% 6% </div> </div>
1	D	453	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 17%, green 75%, grey 7%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> % 75% 17% 7% </div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 13812 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phenylalanine-4-hydroxylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	428	Total	C	N	O	S	0	0	0
			3476	2223	588	655	10			
1	B	426	Total	C	N	O	S	0	0	0
			3458	2212	584	652	10			
1	C	424	Total	C	N	O	S	0	0	0
			3446	2203	584	649	10			
1	D	422	Total	C	N	O	S	0	0	0
			3427	2193	580	644	10			

- Molecule 2 is FE (III) ION (three-letter code: FE) (formula: Fe).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Fe	0	0
			1	1		
2	A	1	Total	Fe	0	0
			1	1		
2	D	1	Total	Fe	0	0
			1	1		
2	C	1	Total	Fe	0	0
			1	1		


- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mg	0	0
			1	1		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Phenylalanine-4-hydroxylase



• Molecule 1: Phenylalanine-4-hydroxylase





• Molecule 1: Phenylalanine-4-hydroxylase





● Molecule 1: Phenylalanine-4-hydroxylase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	96.77Å 102.69Å 202.97Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	101.48 – 3.60 101.48 – 3.60	Depositor EDS
% Data completeness (in resolution range)	99.9 (101.48-3.60) 99.9 (101.48-3.60)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.39	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.12 (at 3.58Å)	Xtriage
Refinement program	PHENIX (1.10_2155: ???)	Depositor
R, R_{free}	0.268 , 0.296 0.270 , 0.296	Depositor DCC
R_{free} test set	2007 reflections (8.32%)	DCC
Wilson B-factor (Å ²)	76.0	Xtriage
Anisotropy	0.324	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , -2.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.43$, $\langle L^2 \rangle = 0.26$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.86	EDS
Total number of atoms	13812	wwPDB-VP
Average B, all atoms (Å ²)	76.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 11.63% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, FE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.25	0/3560	0.43	0/4820
1	B	0.25	0/3542	0.45	0/4797
1	C	0.25	0/3530	0.44	1/4779 (0.0%)
1	D	0.25	0/3511	0.43	0/4755
All	All	0.25	0/14143	0.44	1/19151 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	277	TYR	C-N-CA	5.94	136.55	121.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3476	0	3404	37	1
1	B	3458	0	3383	45	1
1	C	3446	0	3371	44	0
1	D	3427	0	3355	52	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	1	0	0	0	0
All	All	13812	0	13513	166	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 166 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:270:ARG:HH11	1:D:278:THR:HG21	1.46	0.81
1:B:71:ARG:HG3	1:C:297:ARG:HG2	1.66	0.77
1:A:54:LEU:O	1:A:58:ASN:ND2	2.19	0.75
1:C:129:ASP:OD1	1:C:243:ARG:NH2	2.19	0.75
1:D:192:ARG:NH1	1:D:221:GLU:OE2	2.22	0.73

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:159:LYS:CD	1:B:163:ASP:OD2[2_554]	2.11	0.09

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	424/453 (94%)	407 (96%)	16 (4%)	1 (0%)	51	85
1	B	422/453 (93%)	407 (96%)	15 (4%)	0	100	100
1	C	420/453 (93%)	402 (96%)	16 (4%)	2 (0%)	32	73

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	D	418/453 (92%)	401 (96%)	17 (4%)	0	100	100
All	All	1684/1812 (93%)	1617 (96%)	64 (4%)	3 (0%)	51	85

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	278	THR
1	C	21	GLU
1	A	21	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	379/399 (95%)	377 (100%)	2 (0%)	91	97
1	B	377/399 (94%)	373 (99%)	4 (1%)	78	91
1	C	376/399 (94%)	373 (99%)	3 (1%)	85	94
1	D	374/399 (94%)	371 (99%)	3 (1%)	85	94
All	All	1506/1596 (94%)	1494 (99%)	12 (1%)	85	94

5 of 12 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	270	ARG
1	C	30	ASN
1	D	30	ASN
1	B	143	ASP
1	C	424	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 5 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	428/453 (94%)	-0.04	4 (0%) 84 72	62, 75, 91, 93	0
1	B	426/453 (94%)	-0.19	1 (0%) 94 91	63, 72, 82, 89	0
1	C	424/453 (93%)	-0.08	3 (0%) 87 78	61, 74, 90, 96	0
1	D	422/453 (93%)	0.10	5 (1%) 79 66	71, 83, 95, 100	0
All	All	1700/1812 (93%)	-0.05	13 (0%) 86 75	61, 76, 92, 100	0

The worst 5 of 13 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	23	SER	3.6
1	D	30	ASN	3.4
1	D	136	LEU	3.3
1	A	31	GLN	2.7
1	D	53	ARG	2.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	FE	C	501	1/1	0.79	0.24	-0.02	61,61,61,61	0
2	FE	A	501	1/1	0.95	0.21	-0.78	67,67,67,67	0
2	FE	B	501	1/1	0.94	0.18	-0.93	62,62,62,62	0
2	FE	D	501	1/1	0.90	0.22	-0.96	78,78,78,78	0
3	MG	A	502	1/1	0.85	0.34	-	58,58,58,58	0

6.5 Other polymers [i](#)

There are no such residues in this entry.