



# Full wwPDB NMR Structure Validation Report ⓘ

Feb 12, 2017 – 09:52 pm GMT

PDB ID : 2FHO  
Title : NMR solution structure of the human spliceosomal protein complex p14-SF3b155  
Authors : Kuwasako, K.; Dohmae, N.; Inoue, M.; Shirouzu, M.; Guntert, P.; Seraphin, B.; Muto, Y.; Yokoyama, S.; RIKEN Structural Genomics/Proteomics Initiative (RSGI)  
Deposited on : 2005-12-26

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange	:	Kirchner and Güntert (2011)
NmrClust	:	Kelley et al. (1996)
MolProbity	:	4.02b-467
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
ShiftChecker	:	trunk28760
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*SOLUTION NMR*

The overall completeness of chemical shifts assignment is 93%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	125131	11601
Ramachandran outliers	121729	10391
Sidechain outliers	121581	10367

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	47	
2	B	87	

## 2 Ensemble composition and analysis

This entry contains 20 models. Model 18 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:404-A:418, B:17-B:93 (92)	0.39	18

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters and 4 single-model clusters were found.

Cluster number	Models
1	2, 3, 5, 6, 9, 10, 13, 16, 17, 18, 20
2	4, 7, 11, 15, 19
Single-model clusters	1; 8; 12; 14

### 3 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 1996 atoms, of which 893 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called spliceosomal protein SF3b155.

Mol	Chain	Residues	Atoms						Trace
1	A	47	Total	C	H	N	O	S	0
			681	248	293	65	74	1	

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	378	GLY	-	CLONING ARTIFACT	UNP O75533

- Molecule 2 is a protein called spliceosomal protein p14.

Mol	Chain	Residues	Atoms						Trace
2	B	87	Total	C	H	N	O	S	0
			1315	459	600	125	128	3	

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	7	GLY	-	CLONING ARTIFACT	UNP Q9Y3B4

## 4 Residue-property plots [i](#)


### 4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: spliceosomal protein SF3b155



- Molecule 2: spliceosomal protein p14




### 4.2 Scores per residue for each member of the ensemble


Colouring as in section 4.1 above.

#### 4.2.1 Score per residue for model 1

- Molecule 1: spliceosomal protein SF3b155




- Molecule 2: spliceosomal protein p14




### 4.2.2 Score per residue for model 2

- Molecule 1: spliceosomal protein SF3b155




- Molecule 2: spliceosomal protein p14




### 4.2.3 Score per residue for model 3

- Molecule 1: spliceosomal protein SF3b155




- Molecule 2: spliceosomal protein p14




### 4.2.4 Score per residue for model 4

- Molecule 1: spliceosomal protein SF3b155




- Molecule 2: spliceosomal protein p14




### 4.2.5 Score per residue for model 5

- Molecule 1: spliceosomal protein SF3b155




- Molecule 2: spliceosomal protein p14




### 4.2.6 Score per residue for model 6

- Molecule 1: spliceosomal protein SF3b155




- Molecule 2: spliceosomal protein p14




### 4.2.7 Score per residue for model 7

- Molecule 1: spliceosomal protein SF3b155




- Molecule 2: spliceosomal protein p14




### 4.2.8 Score per residue for model 8

- Molecule 1: spliceosomal protein SF3b155




- Molecule 2: spliceosomal protein p14




### 4.2.9 Score per residue for model 9

- Molecule 1: spliceosomal protein SF3b155




- Molecule 2: spliceosomal protein p14




### 4.2.10 Score per residue for model 10

- Molecule 1: spliceosomal protein SF3b155




- Molecule 2: spliceosomal protein p14






### 4.2.11 Score per residue for model 11

- Molecule 1: spliceosomal protein SF3b155




- Molecule 2: spliceosomal protein p14




### 4.2.12 Score per residue for model 12

- Molecule 1: spliceosomal protein SF3b155




- Molecule 2: spliceosomal protein p14




### 4.2.13 Score per residue for model 13

- Molecule 1: spliceosomal protein SF3b155




- Molecule 2: spliceosomal protein p14




#### 4.2.14 Score per residue for model 14

- Molecule 1: spliceosomal protein SF3b155




- Molecule 2: spliceosomal protein p14




#### 4.2.15 Score per residue for model 15

- Molecule 1: spliceosomal protein SF3b155




- Molecule 2: spliceosomal protein p14




#### 4.2.16 Score per residue for model 16

- Molecule 1: spliceosomal protein SF3b155



- Molecule 2: spliceosomal protein p14




### 4.2.17 Score per residue for model 17

- Molecule 1: spliceosomal protein SF3b155




- Molecule 2: spliceosomal protein p14




### 4.2.18 Score per residue for model 18 (medoid)

- Molecule 1: spliceosomal protein SF3b155



- Molecule 2: spliceosomal protein p14




### 4.2.19 Score per residue for model 19

- Molecule 1: spliceosomal protein SF3b155




- Molecule 2: spliceosomal protein p14




4.2.20 Score per residue for model 20

- Molecule 1: spliceosomal protein SF3b155



- Molecule 2: spliceosomal protein p14



## 5 Refinement protocol and experimental data overview

The models were refined using the following method: *torsion angle dynamics, RESTRAINED ENERGY MINIMIZATION*.

Of the 100 calculated structures, 20 were deposited, based on the following criterion: *structures with the least restraint violations, structures with the lowest energy, target function*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CYANA	structure solution	2.1
OPALP	structure solution	1.4
CYANA	refinement	2.1
OPALP	refinement	1.4

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	BMRB entry 11366, BMRB entry 11367
Number of chemical shift lists	2
Total number of shifts	2003
Number of shifts mapped to atoms	2003
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	93%

No validations of the models with respect to experimental NMR restraints is performed at this time.

## 6 Model quality

### 6.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.69±0.01	0±0/123 (0.0±0.0%)	0.90±0.05	0±0/170 (0.0±0.1%)
2	B	0.71±0.01	0±0/652 (0.0±0.0%)	0.96±0.03	1±1/885 (0.1±0.1%)
All	All	0.70	0/15500 (0.0%)	0.95	12/21100 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	0.1±0.2
2	B	0.0±0.0	0.7±0.9
All	All	0	15

There are no bond-length outliers.

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
2	B	46	ARG	NE-CZ-NH2	-6.13	117.24	120.30	6	1
2	B	85	ARG	NE-CZ-NH2	-6.06	117.27	120.30	5	3
2	B	57	ARG	NE-CZ-NH2	-6.02	117.29	120.30	6	1
2	B	49	ARG	NE-CZ-NH2	-5.87	117.36	120.30	3	2
2	B	24	ARG	NE-CZ-NH2	-5.81	117.39	120.30	17	1
2	B	91	TYR	CB-CG-CD2	-5.60	117.64	121.00	5	1
2	B	49	ARG	NE-CZ-NH1	5.21	122.90	120.30	3	1
2	B	57	ARG	CD-NE-CZ	5.13	130.78	123.60	12	1
1	A	405	ASP	CB-CG-OD1	5.08	122.88	118.30	20	1

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
2	B	64	TYR	Sidechain	4
2	B	92	TYR	Sidechain	2
2	B	46	ARG	Sidechain	2
2	B	57	ARG	Sidechain	2
2	B	85	ARG	Sidechain	1
1	A	412	TYR	Sidechain	1
2	B	49	ARG	Sidechain	1
2	B	86	TYR	Sidechain	1
2	B	22	TYR	Sidechain	1

## 6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	117	92	117	0±0
2	B	637	535	625	0±0
All	All	15080	12540	14840	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
2:B:17:VAL:HG13	2:B:91:TYR:CE1	0.48	2.43	17	1
1:A:409:PRO:HG2	1:A:412:TYR:CE1	0.41	2.51	2	1
2:B:17:VAL:HG11	2:B:71:LYS:HG2	0.41	1.93	6	1

## 6.3 Torsion angles [i](#)

### 6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	15/47 (32%)	15±1 (98±4%)	0±1 (2±4%)	0±0 (0±0%)	100	100
2	B	76/87 (87%)	72±1 (95±2%)	4±2 (5±2%)	0±0 (0±0%)	58	87
All	All	1820/2680 (68%)	1742 (96%)	77 (4%)	1 (0%)	58	87

All 1 unique Ramachandran outliers are listed below.

Mol	Chain	Res	Type	Models (Total)
2	B	83	CYS	1

### 6.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	13/41 (32%)	13±0 (98±3%)	0±0 (2±3%)	72	96
2	B	68/76 (89%)	65±2 (96±3%)	3±2 (4±3%)	40	84
All	All	1620/2340 (69%)	1560 (96%)	60 (4%)	43	86

All 23 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
2	B	19	ARG	6
2	B	52	ASN	6
2	B	76	HIS	5
2	B	46	ARG	5
1	A	413	LYS	4
2	B	41	LYS	3
2	B	66	ASP	3
2	B	29	LYS	3
2	B	75	ASP	3
2	B	69	ASP	3
2	B	71	LYS	3
2	B	74	CYS	2
2	B	53	THR	2
2	B	49	ARG	2
2	B	81	ASN	2
2	B	18	ASN	1

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Models (Total)
2	B	31	THR	1
2	B	67	ILE	1
2	B	37	ASP	1
2	B	47	GLN	1
2	B	78	SER	1
2	B	72	ASN	1
2	B	57	ARG	1

### 6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 6.7 Other polymers [i](#)

There are no such molecules in this entry.

### 6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 93% for the well-defined parts and 86% for the entire structure.

### 7.1 Chemical shift list 1

File name: BMRB entry 11366

Chemical shift list name: *assigned\_chem\_shift\_list\_1*

#### 7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1432
Number of shifts mapped to atoms	1432
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

#### 7.1.2 Chemical shift referencing

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction $\pm$ precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	112	$-0.23 \pm 0.15$	None needed ( $< 0.5$ ppm)
$^{13}\text{C}_\beta$	105	$-0.01 \pm 0.18$	None needed ( $< 0.5$ ppm)
$^{13}\text{C}'$	99	$0.09 \pm 0.15$	None needed ( $< 0.5$ ppm)
$^{15}\text{N}$	100	$0.23 \pm 0.33$	None needed ( $< 0.5$ ppm)

#### 7.1.3 Completeness of resonance assignments

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 93%, i.e. 1089 atoms were assigned a chemical shift out of a possible 1176. 0 out of 15 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^1\text{H}$	$^{13}\text{C}$	$^{15}\text{N}$
Backbone	440/446 (99%)	177/177 (100%)	178/184 (97%)	85/85 (100%)
Sidechain	529/607 (87%)	326/357 (91%)	192/220 (87%)	11/30 (37%)

*Continued on next page...*

Continued from previous page...

	Total	<sup>1</sup> H	<sup>13</sup> C	<sup>15</sup> N
Aromatic	120/123 (98%)	62/64 (97%)	58/58 (100%)	0/1 (0%)
Overall	1089/1176 (93%)	565/598 (94%)	428/462 (93%)	96/116 (83%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 74%, i.e. 1290 atoms were assigned a chemical shift out of a possible 1734. 0 out of 19 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	<sup>1</sup> H	<sup>13</sup> C	<sup>15</sup> N
Backbone	522/646 (81%)	211/256 (82%)	211/268 (79%)	100/122 (82%)
Sidechain	640/933 (69%)	393/551 (71%)	235/330 (71%)	12/52 (23%)
Aromatic	128/155 (83%)	66/80 (82%)	62/72 (86%)	0/3 (0%)
Overall	1290/1734 (74%)	670/887 (76%)	508/670 (76%)	112/177 (63%)


#### 7.1.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.


#### 7.1.5 Random Coil Index (RCI) plots [i](#)

The images below report *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:



Random coil index (RCI) for chain B:



## 7.2 Chemical shift list 2

File name: BMRB entry 11367

Chemical shift list name: *assigned\_chem\_shift\_list\_1*

### 7.2.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	571
Number of shifts mapped to atoms	571
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

### 7.2.2 Chemical shift referencing [i](#)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction $\pm$ precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	46	$1.13 \pm 0.23$	Should be applied
$^{13}\text{C}_\beta$	43	$-0.05 \pm 0.11$	None needed ( $< 0.5$ ppm)
$^{13}\text{C}'$	41	$0.93 \pm 0.18$	Should be applied
$^{15}\text{N}$	37	$-0.92 \pm 0.13$	Should be applied

### 7.2.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 15%, i.e. 178 atoms were assigned a chemical shift out of a possible 1176. 0 out of 15 assigned methyl groups (LEU and VAL) were assigned

stereospecifically.

	Total	$^1\text{H}$	$^{13}\text{C}$	$^{15}\text{N}$
Backbone	65/446 (15%)	26/177 (15%)	28/184 (15%)	11/85 (13%)
Sidechain	97/607 (16%)	61/357 (17%)	36/220 (16%)	0/30 (0%)
Aromatic	16/123 (13%)	8/64 (12%)	8/58 (14%)	0/1 (0%)
Overall	178/1176 (15%)	95/598 (16%)	72/462 (16%)	11/116 (9%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 30%, i.e. 528 atoms were assigned a chemical shift out of a possible 1734. 0 out of 19 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^1\text{H}$	$^{13}\text{C}$	$^{15}\text{N}$
Backbone	207/646 (32%)	83/256 (32%)	87/268 (32%)	37/122 (30%)
Sidechain	277/933 (30%)	176/551 (32%)	98/330 (30%)	3/52 (6%)
Aromatic	44/155 (28%)	22/80 (28%)	20/72 (28%)	2/3 (67%)
Overall	528/1734 (30%)	281/887 (32%)	205/670 (31%)	42/177 (24%)

## 7.2.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

## 7.2.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:

