



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 12, 2017 – 08:58 pm GMT

PDB ID : 2FHZ
Title : Molecular Basis of Inhibition of the Ribonuclease Activity in Colicin E5 by Its
Cognate Immunity Protein
Authors : Lin, Y.L.; Huang, R.H.
Deposited on : 2005-12-27
Resolution : 1.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	1078 (1.18-1.10)
Clashscore	112137	1123 (1.18-1.10)
Ramachandran outliers	110173	1074 (1.18-1.10)
Sidechain outliers	110143	1071 (1.18-1.10)
RSRZ outliers	101464	1082 (1.18-1.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	109	<div> <div>6%</div> <div> <div></div> <div>92%</div> <div>6%</div> <div></div> </div> <div></div> </div>
2	B	108	<div> <div>9%</div> <div> <div></div> <div>81%</div> <div>5%</div> <div>14%</div> </div> <div></div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 1821 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Colicin-E5 immunity protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	106	Total	C	N	O	S	0	0	0
			842	541	136	161	4			

There are 26 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	INITIATING METHIONINE	UNP P13476
A	2	THR	-	CLONING ARTIFACT	UNP P13476
A	3	ASN	-	CLONING ARTIFACT	UNP P13476
A	4	LYS	-	CLONING ARTIFACT	UNP P13476
A	5	LEU	-	CLONING ARTIFACT	UNP P13476
A	6	PHE	-	CLONING ARTIFACT	UNP P13476
A	7	GLU	-	CLONING ARTIFACT	UNP P13476
A	8	HIS	-	CLONING ARTIFACT	UNP P13476
A	9	THR	-	CLONING ARTIFACT	UNP P13476
A	10	VAL	-	CLONING ARTIFACT	UNP P13476
A	11	LEU	-	CLONING ARTIFACT	UNP P13476
A	12	TYR	-	CLONING ARTIFACT	UNP P13476
A	13	ASP	-	CLONING ARTIFACT	UNP P13476
A	14	SER	-	CLONING ARTIFACT	UNP P13476
A	15	GLY	-	CLONING ARTIFACT	UNP P13476
A	16	ASP	-	CLONING ARTIFACT	UNP P13476
A	17	ALA	-	CLONING ARTIFACT	UNP P13476
A	18	PHE	-	CLONING ARTIFACT	UNP P13476
A	19	PHE	-	CLONING ARTIFACT	UNP P13476
A	20	GLU	-	CLONING ARTIFACT	UNP P13476
A	21	LEU	-	CLONING ARTIFACT	UNP P13476
A	22	LYS	-	CLONING ARTIFACT	UNP P13476
A	23	GLY	-	CLONING ARTIFACT	UNP P13476
A	24	ASN	-	CLONING ARTIFACT	UNP P13476
A	25	ALA	-	CLONING ARTIFACT	UNP P13476
A	26	SER	-	CLONING ARTIFACT	UNP P13476

- Molecule 2 is a protein called Colicin-E5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	93	Total	C	N	O	S	0	0	0
			729	449	131	148	1			


There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	1	MET	-	INITIATING METHIONINE	UNP P18000

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	137	Total	O	0	0
			137	137		
3	B	113	Total	O	0	0
			113	113		

- Molecule 1: Colicin-E5 immunity protein



-
- | Category | Count |
|----------|-------|
| MET | 1 |
| LEU | 1 |
| ALA | 1 |
| LYS | 1 |
| ASN | 1 |
| LYS | 1 |
| GLY | 1 |
| ILE | 1 |
| PRO | 1 |
| GLY | 1 |
| L12 | 1 |
| K13 | 1 |
| T28 | 1 |
| E29 | 1 |
| D30 | 1 |
| K33 | 1 |
| N34 | 1 |
| K47 | 1 |
| K51 | 1 |
| Y67 | 1 |
| V74 | 1 |
| G103 | 1 |
| N104 | 1 |
| LYS | 1 |
| ASN | 1 |
| ASP | 1 |
| GLN | 1 |

4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	61.79Å 73.64Å 110.11Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	25.00 – 1.15 23.96 – 1.15	Depositor EDS
% Data completeness (in resolution range)	92.7 (25.00-1.15) 96.1 (23.96-1.15)	Depositor EDS
R_{merge}	0.03	Depositor
R_{sym}	0.03	Depositor
$\langle I/\sigma(I) \rangle$ ¹	7.44 (at 1.15Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.195 , 0.200 0.196 , 0.201	Depositor DCC
R_{free} test set	6985 reflections (8.14%)	DCC
Wilson B-factor (Å ²)	9.0	Xtriage
Anisotropy	0.101	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 45.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	1821	wwPDB-VP
Average B, all atoms (Å ²)	12.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.86% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.30	0/862	0.56	0/1162
2	B	0.28	0/745	0.60	0/1011
All	All	0.29	0/1607	0.57	0/2173

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	842	0	823	4	0
2	B	729	0	702	4	0
3	A	137	0	0	1	0
3	B	113	0	0	0	0
All	All	1821	0	1525	8	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (8) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:40:GLU:HG3	1:A:44:LYS:HE2	1.76	0.67

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:35:ILE:HD12	1:A:89:ILE:HG23	1.83	0.60
2:B:104:ASN:N	2:B:104:ASN:HD22	2.07	0.52
2:B:47:LYS:HD3	2:B:47:LYS:C	2.33	0.49
1:A:40:GLU:CG	1:A:44:LYS:HE2	2.44	0.47
2:B:29:GLU:CD	2:B:29:GLU:H	2.18	0.46
1:A:55:HIS:HD2	3:A:121:HOH:O	1.97	0.45
2:B:67:TYR:HB2	2:B:74:VAL:CG1	2.49	0.43

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	104/109 (95%)	102 (98%)	2 (2%)	0	100	100
2	B	91/108 (84%)	87 (96%)	4 (4%)	0	100	100
All	All	195/217 (90%)	189 (97%)	6 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	89/92 (97%)	88 (99%)	1 (1%)	78	40

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
2	B	81/93 (87%)	79 (98%)	2 (2%)	53 11
All	All	170/185 (92%)	167 (98%)	3 (2%)	64 22

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	71	TYR
2	B	51	LYS
2	B	104	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	39	ASN
1	A	55	HIS
1	A	85	ASN
2	B	38	ASN
2	B	101	GLN
2	B	104	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	106/109 (97%)	0.71	7 (6%) 19 20	6, 9, 16, 22	0
2	B	93/108 (86%)	0.78	10 (10%) 6 9	6, 9, 19, 25	0
All	All	199/217 (91%)	0.74	17 (8%) 11 14	6, 9, 18, 25	0

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	29	GLU	4.8
2	B	104	ASN	4.6
1	A	79	SER	4.4
2	B	30	ASP	4.2
1	A	109	MET	4.1
1	A	20	GLU	3.6
2	B	33	LYS	3.3
2	B	34	ASN	3.2
2	B	12	LEU	3.2
2	B	103	GLY	3.2
1	A	51	ILE	2.9
1	A	5	LEU	2.5
2	B	28	THR	2.5
2	B	13	LYS	2.4
2	B	51	LYS	2.3
1	A	58	ASN	2.1
1	A	105	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.